arXiv:1304.7760v1 [hep-th] 29 Apr 2013

Implications of conformal invariance in momentum space

Adam Bzowski[a,d], Paul McFadden[c] and Kostas Skenderis[a,b,d,e]

[a] *Mathematical Sciences, University of Southampton, UK.*
[b] *STAG research centre, University of Southampton, UK.*
[c] *Perimeter Institute for Theoretical Physics, Waterloo, Canada.*
[d] *Korteweg-de Vries Institute for Mathematics, Amsterdam, Netherlands.*
[e] *Institute of Physics, Amsterdam, Netherlands.*

E-mail: a.w.bzowski@uva.nl, pmcfadden@perimeterinstitute.ca, k.skenderis@soton.ac.uk

Abstract

We present a comprehensive analysis of the implications of conformal invariance for 3-point functions of the stress-energy tensor, conserved currents and scalar operators in general dimension and *in momentum space*. Our starting point is a novel and very effective decomposition of tensor correlators which reduces their computation to that of a number of scalar form factors. For example, the most general 3-point function of a conserved and traceless stress-energy tensor is determined by only five form factors. Dilatations and special conformal Ward identities then impose additional conditions on these form factors. The special conformal Ward identities become a set of first and second order differential equations, whose general solution is given in terms of integrals involving a product of three Bessel functions ('triple-K integrals'). All in all, the correlators are completely determined up to a number of constants, in agreement with well-known position space results. We develop systematic methods for explicitly evaluating the triple-K integrals. In odd dimensions they are given in terms of elementary functions while in even dimensions the results involve dilogarithms. In some cases, the triple-K integrals diverge and subtractions are necessary and we show how such subtractions are related to conformal anomalies.

This paper contains two parts that can be read independently of each other. In the first part, we explain the method that leads to the solution for the correlators in terms of triple-K integrals and how to evaluate these integrals, while the second part contains a self-contained presentation of all results. Readers interested only in results may directly consult the second part of the paper.

Contents

1 Introduction and summary of results **2**

2 3-point function of scalar operators **9**
2.1 From position to momentum space . . . 9
2.2 Conformal Ward identities . . . 10
2.3 Uniqueness of the solution . . . 11
2.4 Region of validity and anomalies . . . 13

3 Decomposition of tensors **14**
3.1 Representations of tensor structures . . . 14
3.2 Decomposition of $\langle t^{\mu_1\nu_1} t^{\mu_2\nu_2} t^{\mu_3\nu_3} \rangle$. . . 16
3.3 Finding the form factors . . . 18
3.4 Example . . . 19

4 Conformal Ward identities in momentum space **20**
4.1 From position space to momentum space . . . 20
4.2 Dilatation Ward identity . . . 21
4.3 Special conformal Ward identities . . . 22
4.4 Transverse Ward identities . . . 25
4.5 Trace Ward identities . . . 28

5 Solutions to conformal Ward identities **28**
5.1 Triple-K integrals . . . 29
5.2 Solutions to the primary conformal Ward identities . . . 31
5.3 Solutions to the secondary conformal Ward identities . . . 34

6 Evaluation of triple-K integrals **42**
6.1 Reduction scheme . . . 42
6.2 The case $d = 4$. . . 43
6.3 Integrals in even dimensions $d \geq 4$. . . 45
6.4 Evaluation of the master integral . . . 46

7 Worked example: $\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3} \rangle$ **47**
7.1 Primary conformal Ward identities . . . 48
7.2 Evaluation of secondary conformal Ward identities . . . 48
7.3 Solutions to secondary conformal Ward identities . . . 50
7.4 General form of $\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3} \rangle$ in $d = 3$. . . 51
7.5 Free fermions in $d = 3$. . . 53
7.6 General form of $\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3} \rangle$ in $d = 4$. . . 54
7.7 Free fermions in $d = 4$. . . 54

8 Divergences and anomalies in $d = 4$ **55**
8.1 Counterterms and anomalies . . . 56
8.2 $\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3} \rangle$. . . 57
8.3 $\langle T^{\mu_1\nu_1} T^{\mu_2\nu_2} T^{\mu_3\nu_3} \rangle$. . . 58

9 Extensions **59**
9.1 Helicity formalism . . . 59
9.2 Higher-point correlation functions . . . 61

List of results **63**
1 Definitions . . . 63
2 $\langle \mathcal{O}\mathcal{O}\mathcal{O}\rangle$. . . 65
3 $\langle J^{\mu_1}\mathcal{O}\mathcal{O}\rangle$. . . 65
4 $\langle J^{\mu_1}J^{\mu_2}\mathcal{O}\rangle$. . . 67
5 $\langle J^{\mu_1}J^{\mu_2}J^{\mu_3}\rangle$. . . 70
6 $\langle T^{\mu_1\nu_1}\mathcal{O}\mathcal{O}\rangle$. . . 72
7 $\langle T^{\mu_1\nu_1}J^{\mu_2}\mathcal{O}\rangle$. . . 74
8 $\langle T^{\mu_1\nu_1}J^{\mu_2}J^{\mu_3}\rangle$. . . 75
9 $\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}\mathcal{O}\rangle$. . . 79
10 $\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}J^{\mu_3}\rangle$. . . 83
11 $\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}T^{\mu_3\nu_3}\rangle$. . . 85

Appendix **92**
A.1 Decomposition of $\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}T^{\mu_3\nu_3}\rangle$ in non-conformal case . . . 92
A.2 Degeneracy of the tensor structure of $\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}T^{\mu_3\nu_3}\rangle$ in $d=3$. . . 94
A.3 Triple-K integrals by Fourier transform . . . 95
A.4 Properties of triple-K integrals . . . 98
A.5 Appell's F_4 function . . . 99
A.6 Master integral . . . 102
A.7 Triviality of $\langle T^{\mu_1\nu_1}J^{\mu_2}\mathcal{O}\rangle$. . . 103
A.8 Form of scaling anomalies . . . 104
A.9 Identities with projectors . . . 106

1 Introduction and summary of results

It is well known that conformal invariance imposes strong constraints on correlation functions. In particular, 2- and 3-point functions of the stress-energy tensor, conserved currents and scalar primary operators are completely determined up to a few constants. The result for the 3-point function of scalar primary operators already appeared in [1], while the 3-point function of currents for $d = 4$ was determined a few years later in [2]. A complete analysis of all such 3-point functions, and in general dimension, was carried out in [3, 4]; for a sample of more recent work on this topic see also [5, 6, 7, 8, 9, 10, 11]. All of these papers obtain their results through the use position space techniques.

The purpose of this paper is to present the analogous set of results in momentum space. In principle, the results in momentum space can be obtained from those in position space by Fourier transform. Typically, however, the position space expressions (in the form often quoted) are only valid at separated points, and do not possess a Fourier transform prior to renormalisation.[1] Even after renormalising, it is technically rather difficult to carry out explicitly the Fourier transforms, see for example [13]. Here we will present a complete analysis from first principles of the constraints due to conformal symmetry directly in momentum space. We believe such an analysis gives considerably more insight into the results and is interesting in its own right.

A momentum space analysis is natural from the perspective of Feynman diagram computations, which are usually performed in momentum space. Furthermore, a number of recent works have exemplified the need for CFT results in momentum space. Our original motivation for analysing this question was the requirement for these results in our work on holographic cosmology [14, 15, 16, 17, 18], and similar applications of the conformal/de Sitter symmetry in

[1] For example, the 2-point function $\langle \mathcal{O}(\boldsymbol{x})\mathcal{O}(0)\rangle \sim 1/|\boldsymbol{x}|^{2\Delta}$ of a scalar operator $\mathcal{O}$ of dimension Δ does not possess a Fourier transform due to short distance singularities when $\Delta = d/2 + k,\ k = 0, 1, 2, \ldots$. Only after renormalisation (using, for example, differential renormalisation [12]) can one Fourier transform this correlator to momentum space.

cosmology have been discussed in [19, 20, 21, 22, 23, 24, 25]. Other recent works that contain explicit computations of CFT correlation functions in momentum space include [26, 27, 28, 13, 29]. Our results may also be useful in the context of work on an a-theorem in diverse dimensions, see [30] for a relevant discussion in $d = 4$.

There are two main issues that complicate the analysis of the implications of conformal invariance in momentum space. While conformal transformations act naturally in position space, they lead to differential operators in momentum space. Dilatations, $\delta x^\mu = \lambda x^\mu$, being linear in x^μ lead to a Ward identity that is a first-order differential equation, and as such, it is easy to solve in complete generality. Special conformal transformations however are non-linear, $\delta x^\mu = b^\mu x^2 - 2x^\mu b\cdot x$, so after Fourier transform we obtain a Ward identity that is a second-order differential equation, which makes the analysis more complicated.

The second main issue is the complicated tensorial decomposition required for correlators involving vectors and tensors. We will return to this below, but let us begin by illustrating the first issue, focusing on the case of the 3-point function of scalar operators $\mathcal{O}_j$ of dimension Δ_j, $\langle \mathcal{O}_1(\boldsymbol{p}_1)\mathcal{O}_2(\boldsymbol{p}_2)\mathcal{O}_3(\boldsymbol{p}_3)\rangle$. We will discuss this computation in detail in section 2, so here we simply summarise the main points. As usual, translational invariance implies that we can pull out a momentum-conserving delta function,

$$\langle \mathcal{O}_1(\boldsymbol{p}_1)\mathcal{O}_2(\boldsymbol{p}_2)\mathcal{O}_3(\boldsymbol{p}_3)\rangle = (2\pi)^d \delta(\boldsymbol{p}_1+\boldsymbol{p}_2+\boldsymbol{p}_3)\langle\!\langle \mathcal{O}_1(\boldsymbol{p}_1)\mathcal{O}_2(\boldsymbol{p}_2)\mathcal{O}_3(\boldsymbol{p}_3)\rangle\!\rangle, \tag{1.1}$$

thereby defining the reduced matrix element which we denote with double brackets. Lorentz invariance then implies that $\langle\!\langle \mathcal{O}_1(\boldsymbol{p}_1)\mathcal{O}_2(\boldsymbol{p}_2)\mathcal{O}_3(\boldsymbol{p}_3)\rangle\!\rangle$ is only a function of the magnitude of the momenta $p_j = |\boldsymbol{p}_j|$, while dilatation invariance implies that it is a homogeneous function of total degree $(\Delta_t - 2d)$, where $\Delta_t = \sum \Delta_j$.

Finally, we impose invariance under special conformal transformations. The corresponding Ward identities are second-order differential equations which can be manipulated into the form

$$0 = \mathrm{K}_{12}\langle\!\langle \mathcal{O}_1(\boldsymbol{p}_1)\mathcal{O}_2(\boldsymbol{p}_2)\mathcal{O}_3(\boldsymbol{p}_3)\rangle\!\rangle = \mathrm{K}_{23}\langle\!\langle \mathcal{O}_1(\boldsymbol{p}_1)\mathcal{O}_2(\boldsymbol{p}_2)\mathcal{O}_3(\boldsymbol{p}_3)\rangle\!\rangle, \tag{1.2}$$

where

$$\mathrm{K}_{ij} = \mathrm{K}_i - \mathrm{K}_j, \qquad \mathrm{K}_j = \frac{\partial^2}{\partial p_j^2} + \frac{d+1-2\Delta_j}{p_j}\frac{\partial}{\partial p_j}, \qquad i,j = 1,2,3. \tag{1.3}$$

This system of differential equations is precisely that defining Appell's F_4 generalised hypergeometric function of two variables. There are four linearly independent solutions of these equations but three of them have unphysical singularities at certain values of the momenta leaving one physically acceptable solution. This solution has the following integral representation, which we will refer to as a *triple-K integral*:[2]

$$\begin{aligned}&\langle\!\langle \mathcal{O}_1(\boldsymbol{p}_1)\mathcal{O}_2(\boldsymbol{p}_2)\mathcal{O}_3(\boldsymbol{p}_3)\rangle\!\rangle \\ &\quad = C_{123} p_1^{\Delta_1-\frac{d}{2}} p_2^{\Delta_2-\frac{d}{2}} p_3^{\Delta_3-\frac{d}{2}} \int_0^\infty \mathrm{d}x\, x^{\frac{d}{2}-1} K_{\Delta_1-\frac{d}{2}}(p_1 x) K_{\Delta_2-\frac{d}{2}}(p_2 x) K_{\Delta_3-\frac{d}{2}}(p_3 x),\end{aligned} \tag{1.4}$$

where $K_\nu(p)$ is a modified Bessel function of the second kind (or Bessel K function, for short) and C_{123} is an overall undetermined constant. We thus arrive at the conclusion that scalar 3-point functions are uniquely determined up to one constant.

This result is still formal, however, since the integral in (1.4) may not converge. Depending on the conformal dimensions involved there are three cases: (i) the integral converges; (ii) the integral diverges but it can be defined via analytic continuation in the spacetime dimension and the conformal dimensions Δ_i; (iii) the integral diverges logarithmically and explicit subtractions

[2] This expression can also obtained by direct Fourier transform of the well-known position space result, see *e.g.*, [31].

are necessary. In the last case, after renomalisation, the correlators exhibit anomalous scaling transformations: the theory suffers from conformal anomalies. This is analogous to the discussion of 2-point functions (see footnote 1): renormalising the 2-point functions results in conformal anomalies, see, *e.g.*, the discussion in [32].

We now turn to discuss tensorial correlation functions, such as those involving stress-energy tensors and conserved currents. Lorentz invariance implies that the tensor structure will be carried by tensors constructed from the momenta p^μ and the metric $\delta_{\mu\nu}$ (throughout this paper we work with Euclidean signature). The standard procedure consists of writing down all possible such independent tensor structures and expressing the correlators as a sum of these structures, each multiplied by scalar form factor. In the case of correlators involving conserved currents and/or stress-energy tensors one then imposes the restrictions enforced by conservation (and tracelessness of the stress-energy tensor in the case of CFTs). Recent works discussing such a tensor decomposition include [28, 30, 27, 26, 13]. This methodology is in principle straightforward, but an inefficient parametrisation can produce unwieldy expressions. Here we present a new parametrisation that appears to yield a minimal number of form factors.

Before proceeding with this, let us briefly discuss the transverse and trace Ward identities (also known as the diffeomorphism and Weyl Ward identities, respectively). The fact that classically a current or stress-energy tensor is conserved implies that n-point functions involving insertions of $\partial_\alpha J^\alpha$ or $\partial_\alpha T^{\alpha\beta}$ are semi-local (*i.e.*, at least two points are coincident) and can be expressed in terms of lower-point functions without such insertions. Similarly, the trace Ward identity implies that correlation functions with insertions of the trace of the stress-energy tensor are also semi-local and are related to lower-point functions. The first step in our analysis is to implement these Ward identities. We do this by providing reconstruction formulae that yield the full 3-point functions involving stress-energy tensors/currents/scalar operators starting from expressions that are exactly conserved/traceless. These 3-point functions automatically satisfy the transverse and trace Ward identities.

To determine the general form of correlators consistent with the transverse and trace Ward identities, it thus suffices to start from an expression that is exactly conserved/traceless in all relevant indices. Such an expression may be obtained by means of projection operators. Recall that in momentum space the operator

$$\pi^\mu_\alpha(\boldsymbol{p}) = \delta^\mu_\alpha - \frac{p^\mu p_\alpha}{p^2} \tag{1.5}$$

is a projector onto tensors transverse to $\boldsymbol{p}$, *i.e.*, $p_\mu \pi^\mu_\alpha(\boldsymbol{p}) = 0$. Similarly, in d dimensions, the operator

$$\Pi^{\mu\nu}_{\alpha\beta}(\boldsymbol{p}) = \frac{1}{2}\left(\pi^\mu_\alpha(\boldsymbol{p})\pi^\nu_\beta(\boldsymbol{p}) + \pi^\mu_\beta(\boldsymbol{p})\pi^\nu_\alpha(\boldsymbol{p})\right) - \frac{1}{d-1}\pi^{\mu\nu}(\boldsymbol{p})\pi_{\alpha\beta}(\boldsymbol{p}) \tag{1.6}$$

is a projector onto transverse to $\boldsymbol{p}$, traceless, symmetric tensors of rank two.

To illustrate our discussion we will use as an example the 3-point function of the stress-energy tensor, $\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)T^{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle$. This is the most complicated case, but also perhaps the most interesting one. In the main text we will explain the method using the simpler example of $\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle$, and in part II we present the corresponding results for all 3-point functions.

To obtain the most general 3-point function of the stress-energy tensor satisfying the transverse and trace Ward identities it suffices to start from the following transverse-traceless expression,

$$\langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1)t^{\mu_2\nu_2}(\boldsymbol{p}_2)t^{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\!\rangle = \Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1)\Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)\Pi^{\mu_3\nu_3}_{\alpha_3\beta_3}(\boldsymbol{p}_3)X^{\alpha_1\beta_1\alpha_2\beta_2\alpha_3\beta_3}, \tag{1.7}$$

where $X^{\alpha_1\beta_1\alpha_2\beta_2\alpha_3\beta_3}$ is a rank six tensor built from the momenta and the metric. The complete 3-point function may then be obtained using the reconstruction formula in (R.11.4).

At this point it seems that we have not gained much since we traded a rank six tensor, the left-hand side of (1.7), with another rank six tensor, $X^{\alpha_1\beta_1\alpha_2\beta_2\alpha_3\beta_3}$. This is deceptive, however,

as the explicit projection operators annihilate many of the possible terms that can appear in $X^{\alpha_1\beta_1\alpha_2\beta_2\alpha_3\beta_3}$. To obtain the most economic parametrisation we would still like to impose one more requirement. The 3-point function is invariant under permutations of the labels:

$$\langle\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1)t^{\mu_2\nu_2}(\boldsymbol{p}_2)t^{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\rangle = \langle\langle t^{\mu_{\sigma(1)}\nu_{\sigma(1)}}(\boldsymbol{p}_{\sigma(1)})t^{\mu_{\sigma(2)}\nu_{\sigma(2)}}(\boldsymbol{p}_{\sigma(2)})t^{\mu_{\sigma(3)}\nu_{\sigma(3)}}(\boldsymbol{p}_{\sigma(3)})\rangle\rangle, \tag{1.8}$$

where σ denotes any element of the permutation group S_3 of the set $\{1,2,3\}$. We would like to find a parametrisation making this invariance as manifest as possible.

Recall that the reduced matrix elements are multiplied by a momentum-conserving delta function, as in (1.1). Often one uses momentum conservation to solve for one momentum in terms of the others, say $\boldsymbol{p}_3 = -(\boldsymbol{p}_1+\boldsymbol{p}_2)$, and then the right-hand side of (1.7) contains only p_1 and p_2. In doing so, however, one obscures the relation (1.8). Here, we will instead choose the independent momenta differently for different Lorentz indices:

$$\boldsymbol{p}_1, \boldsymbol{p}_2 \text{ for } \mu_1,\nu_1;\ \boldsymbol{p}_2, \boldsymbol{p}_3 \text{ for } \mu_2,\nu_2 \text{ and } \boldsymbol{p}_3, \boldsymbol{p}_1 \text{ for } \mu_3,\nu_3. \tag{1.9}$$

With this choice, it is straightforward to show that $X^{\alpha_1\beta_1\alpha_2\beta_2\alpha_3\beta_3}$ is determined by five form factors:

$$\begin{aligned}
X^{\alpha_1\beta_1\alpha_2\beta_2\alpha_3\beta_3} &= A_1 p_2^{\alpha_1} p_2^{\beta_1} p_3^{\alpha_2} p_3^{\beta_2} p_1^{\alpha_3} p_1^{\beta_3} \\
&\quad + A_2 \delta^{\beta_1\beta_2} p_2^{\alpha_1} p_3^{\alpha_2} p_1^{\alpha_3} p_1^{\beta_3} + A_2(p_1 \leftrightarrow p_3)\delta^{\beta_2\beta_3} p_2^{\alpha_1} p_2^{\beta_1} p_3^{\alpha_2} p_1^{\alpha_3} \\
&\qquad + A_2(p_2 \leftrightarrow p_3)\delta^{\beta_1\beta_3} p_2^{\alpha_1} p_3^{\alpha_2} p_3^{\beta_2} p_1^{\alpha_3} \\
&\quad + A_3 \delta^{\alpha_1\alpha_2}\delta^{\beta_1\beta_2} p_1^{\alpha_3} p_1^{\beta_3} + A_3(p_1 \leftrightarrow p_3)\delta^{\alpha_2\alpha_3}\delta^{\beta_2\beta_3} p_2^{\alpha_1} p_2^{\beta_1} \\
&\qquad + A_3(p_2 \leftrightarrow p_3)\delta^{\alpha_1\alpha_3}\delta^{\beta_1\beta_3} p_3^{\alpha_2} p_3^{\beta_2} \\
&\quad + A_4 \delta^{\alpha_1\alpha_3}\delta^{\alpha_2\beta_3} p_2^{\beta_1} p_3^{\beta_2} + A_4(p_1 \leftrightarrow p_3)\delta^{\alpha_1\alpha_3}\delta^{\alpha_2\beta_1} p_3^{\beta_2} p_1^{\beta_3} \\
&\qquad + A_4(p_2 \leftrightarrow p_3)\delta^{\alpha_1\alpha_2}\delta^{\alpha_3\beta_2} p_2^{\beta_1} p_1^{\beta_3} \\
&\quad + A_5 \delta^{\alpha_1\beta_2}\delta^{\alpha_2\beta_3}\delta^{\alpha_3\beta_1}\,,
\end{aligned} \tag{1.10}$$

where the A_i are functions of three variables, $A_i(p_1,p_2,p_3)$, and $A_2(p_1 \leftrightarrow p_3)$ denotes the same function but with p_1 interchanged with p_3, *i.e.*, $A_2(p_1 \leftrightarrow p_3) = A_2(p_3,p_2,p_1)$, etc.

In this paper we assume that the underlying theory is parity-invariant. Additional parity-violating terms can appear in the tensorial decomposition of the various correlators and it would be interesting to incorporate them in our analysis. We leave this for future work.

The form factors A_1 and A_5 are S_3-invariant,

$$A_j(p_1,p_2,p_3) = A_j(p_{\sigma(1)},p_{\sigma(2)},p_{\sigma(3)}), \quad j \in \{1,5\}, \tag{1.11}$$

while the remaining ones are symmetric under $p_1 \leftrightarrow p_2$, *i.e.*, they satisfy

$$A_j(p_2,p_1,p_3) = A_j(p_1,p_2,p_3), \qquad j \in \{2,3,4\}. \tag{1.12}$$

Under the action of S_3 the first and last terms are invariant on their own, while the three terms with A_2 are mapped to each other and similar for A_3 and A_4.

To illustrate the mechanics behind the decomposition (1.10) let us explain why there no terms with either $p_1^{\alpha_1}$ or $p_3^{\alpha_1}$. First, note that the index α_1 is linked with the indices μ_1 and ν_1 via (1.7) and from (1.9) follows that we should only use $p_1^{\alpha_1}$ and $p_2^{\alpha_1}$ when working out the possible terms (a possible $p_3^{\alpha_1}$ is converted into $-(p_1^{\alpha_1}+p_2^{\alpha_1})$ using momentum conservation). However, terms containing p^{α_1} vanish due to project operator in (1.7). Repeating this argument for the other indices leads to (1.10).

Thus we find that the most general 3-point function of the stress-energy tensor in any dimension satisfying the transverse and trace Ward identities is specified by five form factors.

If we relax the condition of Weyl invariance (*i.e.*, if we consider a stress-energy tensor with non-vanishing trace) then the number of form factors becomes ten, see appendix A.1. These results should be compared with those of [30] in $d = 4$. There, the expansion of the 3-point function for the conformal case (traceless $T_{\mu\nu}$) is done in terms of 13 tensors, of which eight are transverse traceless, two are transverse but not traceless, and three are neither transverse nor traceless. Our reconstruction formula (R.11.4) eliminates the need to construct a basis for the non-transverse-traceless part of the correlator, while the difference between the number of transverse traceless tensors is due to the fact that in [30] the expression is manifestly invariant under $\boldsymbol{p}_1 \leftrightarrow \boldsymbol{p}_2$ and not under the full S_3 permutation group. In the non-conformal case, [30] uses 22 tensor structures while we need only ten.

One may be concerned that the form factors so defined are difficult to extract from explicit computations of correlators, which directly give the entire correlator rather than the transverse traceless piece with the projection operators extracted. It turns out this is not the case. The form factors can be simply extracted by looking at the coefficients of certain tensor structures in the full correlator. For example, in the case of the 3-point function of the stress-energy tensor, one can extract the form factors from the following coefficients of the complete 3-point functions $\langle\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)T^{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\rangle$:

$$A_1 = \text{coefficient of } p_2^{\mu_1} p_2^{\nu_1} p_3^{\mu_2} p_3^{\nu_2} p_1^{\mu_3} p_1^{\nu_3}, \tag{1.13}$$

$$A_2 = 4 \cdot \text{coefficient of } \delta^{\nu_1\nu_2} p_2^{\mu_1} p_3^{\mu_2} p_1^{\mu_3} p_1^{\nu_3}, \tag{1.14}$$

$$A_3 = 2 \cdot \text{coefficient of } \delta^{\mu_1\mu_2}\delta^{\nu_1\nu_2} p_1^{\mu_3} p_1^{\nu_3}, \tag{1.15}$$

$$A_4 = 8 \cdot \text{coefficient of } \delta^{\mu_1\mu_3}\delta^{\mu_2\nu_3} p_2^{\nu_1} p_3^{\nu_2}, \tag{1.16}$$

$$A_5 = 8 \cdot \text{coefficient of } \delta^{\mu_1\nu_2}\delta^{\mu_2\nu_3}\delta^{\mu_3\nu_1}\,. \tag{1.17}$$

We are now ready to impose the dilatation and conformal Ward identities. Let us first define the tensorial dimension N_j of A_j to be equal to the number of momenta in the tensor structure that this form factor multiplies. We use the convention that A_j are ordered according to the tensorial dimension, with the form factor of highest tensorial dimension being the first one, A_1, etc. In the example above, $N_1 = 6, N_2 = 4, N_3 = N_4 = 2, N_5 = 0$. The dilatation Ward identity then implies that the form factors are homogeneous functions of the momenta of degree

$$\deg(A_j) = \Delta_t - 2d - N_j, \tag{1.18}$$

where Δ_t is the sum of the conformal dimensions of the three operators.

All that remains to be discussed are the special conformal Ward identities (CWIs). These split into a set of second-order differential equations, which we call primary CWIs, and a set of first-order partial differential equations, which we call secondary CWIs. The primary CWIs are very similar to the conformal Ward identity found in the case of scalar operators, (1.3). In particular, A_1, the term of highest tensorial dimension always satisfies (1.3), while the terms with lower tensorial dimension satisfy similar equations with linear inhomogeneous terms on the right-hand side. In the case of the 3-point function of $T_{\mu\nu}$, these read:

$$\begin{aligned}
&\mathrm{K}_{12}\, A_1 = 0, && \mathrm{K}_{13}\, A_1 = 0,\\
&\mathrm{K}_{12}\, A_2 = 0, && \mathrm{K}_{13}\, A_2 = 8A_1,\\
&\mathrm{K}_{12}\, A_3 = 0, && \mathrm{K}_{13}\, A_3 = 2A_2,\\
&\mathrm{K}_{12}\, A_4 = 4\left[A_2(p_1 \leftrightarrow p_3) - A_2(p_2 \leftrightarrow p_3)\right], && \mathrm{K}_{13}\, A_4 = -4A_2(p_2 \leftrightarrow p_3),\\
&\mathrm{K}_{12}\, A_5 = 2\left[A_4(p_2 \leftrightarrow p_3) - A_4(p_1 \leftrightarrow p_3)\right], && \mathrm{K}_{13}\, A_5 = 2\left[A_4 - A_4(p_1 \leftrightarrow p_3)\right].
\end{aligned} \tag{1.19}$$

It turns out that each pair of primary CWIs has a unique solution up to one arbitrary constant, and that these solutions can be expressed in terms of triple-K integrals, similar to the solution in (1.4). We will call these integration constants the primary constants. It follows that the 3-point functions are determined up to a number of constants that are at most equal to the number of form factors.

Finally, we need to impose the secondary CWIs. These are first order partial differential equations that depend in particular on the specific 2-point functions that appear in the transverse Ward identities. (As discussed above, the transverse Ward identities relate 3-point functions involving the divergence of the stress-energy tensor/conserved current to 2-point functions.) When inserting the solutions of the primary CWI to the secondary CWIs these become algebraic relations among the primary constants and the normalisation of the operators (through the coefficient of the 2-point functions). Solving these relations we obtain the final number of constants that determine the 3-point function. In the case of the 3-point function of the stress-energy tensor we find that the final answer depends on three constants, which may be taken to be two of the primary constants and the coefficient of the 2-point function. This agrees exactly with the position space analysis in [3].

As in the case of scalar operators, the triple-K integrals sometimes diverge and must be regularised and renormalised. This procedure may lead to conformal anomalies and we discuss in detail the $d = 4$ case. This has the following implications. Firstly, some of the primary constants are related to the anomaly coefficients. In $d = 4$ there are two anomaly coefficients, the c and a coefficients. The c-coefficient is directly related with the normalisation of the 2-point function of the stress-energy tensor, while the a coefficient is equal to a linear combination of the two primary constants and the normalisation of the 2-point function. It follows that the anomalies alone completely fix the 3-point function of the stress-energy tensor up to one constant. The same is true for the 3-point function of the stress-energy tensor and two symmetry currents: this three point function is completely fixed by the corresponding anomaly coefficient up to one constant. A second effect of the conformal anomaly is that specific ultra-local terms[3] in the 3-point functions are fixed by the anomaly and as such they are not scheme dependent. This is due to the fact that the counterterms that remove the infinities may also necessarily contain a finite part as well.

In our discussion so far we showed how to obtain the 3-point functions in terms of triple-K integrals. This is the most economic form of the solution but it is less explicit than the position space result where the 3-point functions are given in terms of elementary functions. Furthermore, in some cases the triple-K integrals diverge so without subtractions this is only a formal solution. In order to obtain completely explicit results we need to evaluate the triple-K integrals (and extract the finite part after regularisation when the integral diverges) and we show how to do this. In odd dimensions the Bessel K functions that appear in the triple-K integrals reduce to elementary functions and one can compute the integrals by standard methods. In even dimensions the evaluation is less trivial. In this case we develop a reduction scheme, generalising Davydychev's recursion relations [33], and show that all integrals can be reduced to a single master integral, which we compute explicitly. The answer involves logarithms and dilogarithms.

We will now discuss in detail the structure of the paper. We split this paper in two parts. In the first part we explain the method we use, while in the second part we give a complete list of all results. The second part is completely self-contained and can be used without reference to the first part of the paper. This second part starts with a collection of all basic definitions and a summary of conventions, followed by a list of all results for all 3-point functions involving the stress-energy tensor, conserved currents and scalar operators. For each such correlators we list: 1) the relevant trace and transverse Ward identities, 2) the reconstruction formula that yield the complete correlator from its transverse(-traceless) part, 3) the tensorial decomposition of the transverse(-traceless) part, 4) how to extract the form factors from the complete correlator, 5) the

[3]Ultra-local terms are those with support at coincident points of all three operators, $x_1 = x_2 = x_3$. In momentum space these terms are analytic in all momenta. Terms that have support at coincident points of two but not all three operators are called semi-local. These terms are analytic in two of the three momenta in momentum space. Ultra-local terms are generally scheme-dependent since such terms can be modified by adding finite local terms in the action. The main exception to this rule is when such terms are fixed by Ward identities and anomalies, such as the case we discuss here.

primary CWIs and their solution in terms of triple-K integrals, 6) the secondary Ward identities and the relations they impose on the integration constants of the primary Ward identities (the primary constants), 7) evaluation of the triple-K integrals in dimensions $d = 3, 4, 5$. Special effort was made so that each section describing any given correlator can be read in isolation so the reader interested only in the results for a specific correlator may directly consult that section.

In the first part we strive to keep a balance between describing the method and computations in enough detail so that all subtleties entering the derivation of the results in Part II are covered, while still keeping the level of technicalities to a minimum. We start in section 2 by discussing in detail the case of the 3-point function of scalar operators. This case serves as a good warm-up exercise, as one has to face many of the subtleties of the conformal Ward identities in momentum space without having to deal with complications due to the tensorial structure of other correlators. Here we see that the solution of the special conformal Ward identities may be expressed in terms of the Appell F_4 function, and discuss its representation in terms of triple-K integrals. Issues of renormalisation and anomalies already show up and are discussed here, again in a simplifying context. This is also a case where the answer can be easily obtained by a Fourier transform so we can directly check our results.

In section 3, we discuss the tensorial decomposition of correlation functions. We illustrate our approach using the case of $\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}\mathcal{O}\rangle$. We also compare in this section our decomposition with previous state-of-the-art results in [30]. In section 4, we derive the form of the conformal Ward identities in momentum space. In particular, we show that the special conformal Ward identities split into the primary and secondary conformal Ward identities.

Section 5 is devoted to the solution of the CWIs. We show first that the primary CWIs can be solved using triple-K integrals and we then substitute these into the secondary CWIs. This results in a number of relations between the primary constants and the normalisation of the 2-point functions, as described earlier. In this section we pay attention to the issues of regularisation and we discuss how to deal with special cases (when the dimension of the scalar operators takes special values, etc.).

In section 6, we discuss how to evaluate the triple-K integrals. We explain how to reduce the computation of integrals to a single master integral and spell out the reduction scheme for the $d = 4$ case, then compute the master integral. The results in this section may be useful in a wider context, and this section can be read independently from the rest of the paper.

In section 7 we work out completely the case of $\langle T^{\mu_1\nu_1}J^{\mu_2}J^{\mu_3}\rangle$, where J^μ is a conserved current. This example is more complex than the $\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}\mathcal{O}\rangle$ case used to illustrate the method in earlier sections, yet simpler than correlators with more stress-energy tensors. We also discuss the complete evaluation of all integrals in $d = 3$ and $d = 4$ and present a concrete model, free fermions, where these correlators can be explicitly computed by standard Feynman diagrams.

In section 8 we return to the issues of anomalies, and focusing on the $d = 4$ case, we discuss the form and implications of the conformal anomalies for the $\langle T^{\mu_1\nu_1}J^{\mu_2}J^{\mu_3}\rangle$ and $\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}T^{\mu_3\nu_3}\rangle$ correlation functions.

Part I finishes with a brief discussion of possible extensions in section 9. One such extension is to develop a helicity formalism. In this formalism one uses helicity projected operators, so we trade the transverse(-traceless) tensor for the two helicity components, so this approach may simplify the tensor decomposition of the 3-point functions. Section 9.1 contains a short introduction to this method. Another extension is to higher-point functions and we briefly discuss this in section 9.2.

Finally, this paper contains a number of appendices. Appendix A.1 contains a discussion of the tensorial decomposition of the 3-point function of a conserved but not traceless stress-energy tensor. In appendix A.2 we explain that the general tensorial decomposition is degenerate when $d = 3$. In this case the 3-point function of the stress-energy tensor is determined by only two

form factors. In appendix A.3 we discuss how to Fourier transform the scalar 3-point functions. In appendix A.4 we list useful properties of the triple-K integrals that are used in the derivations in the main text. Appendix A.5 contains a collection of facts about the Appell F_4 function: its definition, the differential equations it satisfies and useful integral representations. In appendix A.6 we give the explicit form of the master integral needed for the evaluation of the triple-K integrals and discuss a few related integrals. In appendix A.7 we prove the triviality of the $\langle T^{\mu_1\nu_1}J^{\mu_2}\mathcal{O}\rangle$ correlator and in appendix A.8 we present the general form of the contribution of conformal anomalies to 3-point functions. Appendix A.9 contains a set of useful identities involving projection operators, which are used in the derivations in the main text.

We finish this introduction with few comments about how to read this paper. Readers only interested in an overview and specific results may read this introduction and directly move to the relevant section of Part II. Section 2 serves also as a longer and more technical introduction to CFT in momentum space without the complication of tensorial decompositions. Section 6 may be read independently of all other sections.

Note Added: As this paper was finalised [34] appeared which has significant overlap with the discussion in section 2 of this paper. We also understand that our discussion of the tensor decomposition may have overlap with upcoming work by A. Dymarsky [35].

Theory

2 3-point function of scalar operators

2.1 From position to momentum space

Let us begin with a revision of some well-known facts regarding 3-point functions of conformal primary scalar operators in any CFT. Consider three scalar operators $\mathcal{O}_j$, of dimensions Δ_j, $j=1,2,3$. In position space the 3-point function is unique up to an overall constant c_{123} [36],

$$\langle\mathcal{O}_1(\boldsymbol{x}_1)\mathcal{O}_2(\boldsymbol{x}_2)\mathcal{O}_3(\boldsymbol{x}_3)\rangle=\frac{c_{123}}{|\boldsymbol{x}_1-\boldsymbol{x}_2|^{\Delta_1+\Delta_2-\Delta_3}|\boldsymbol{x}_2-\boldsymbol{x}_3|^{\Delta_2+\Delta_3-\Delta_1}|\boldsymbol{x}_3-\boldsymbol{x}_1|^{\Delta_3+\Delta_1-\Delta_2}},\tag{2.1}$$

where we work in Euclidean signature. This expression, in principle, can be Fourier transformed in order to obtain the result in momentum space. Extracting the overall Dirac delta function encoding momentum conservation, we define the reduced matrix element (denoted with double brackets) as in (1.1),

$$\langle\mathcal{O}_1(\boldsymbol{p}_1)\mathcal{O}_2(\boldsymbol{p}_2)\mathcal{O}_3(\boldsymbol{p}_3)\rangle=(2\pi)^d\delta(\boldsymbol{p}_1+\boldsymbol{p}_2+\boldsymbol{p}_3)\langle\!\langle\mathcal{O}_1(\boldsymbol{p}_1)\mathcal{O}_2(\boldsymbol{p}_2)\mathcal{O}_3(\boldsymbol{p}_3)\rangle\!\rangle.\tag{2.2}$$

Assuming $d\geq 3$, since $\boldsymbol{p}_1+\boldsymbol{p}_2+\boldsymbol{p}_3=0$ there are two independent momenta. We define

$$\Delta_t=\Delta_1+\Delta_2+\Delta_3,\qquad\qquad \delta_j=\frac{d-\Delta_t}{2}+\Delta_j,\quad j=1,2,3.\tag{2.3}$$

A useful representation of the Fourier transform of the position space expression (2.1) is

$$\begin{aligned}
&\langle\!\langle\mathcal{O}_1(\boldsymbol{p}_1)\mathcal{O}_2(\boldsymbol{p}_2)\mathcal{O}_3(\boldsymbol{p}_3)\rangle\!\rangle=\\
&\quad=c_{123}\pi^{\frac{3d}{2}}2^{3d-\Delta_t}\prod_{j=1}^{3}\frac{\Gamma(\delta_j)}{\Gamma\left(\frac{d}{2}-\delta_j\right)}\cdot\int\frac{\mathrm{d}^d\boldsymbol{k}}{(2\pi)^d}\frac{1}{|\boldsymbol{k}|^{2\delta_3}|\boldsymbol{p}_1-\boldsymbol{k}|^{2\delta_2}|\boldsymbol{p}_2+\boldsymbol{k}|^{2\delta_1}}\\
&\quad=\frac{c_{123}\pi^d2^{4+\frac{3d}{2}-\Delta_t}}{\Gamma\left(\frac{\Delta_t-d}{2}\right)\Gamma\left(\frac{\Delta_1+\Delta_2-\Delta_3}{2}\right)\Gamma\left(\frac{\Delta_2+\Delta_3-\Delta_1}{2}\right)\Gamma\left(\frac{\Delta_3+\Delta_1-\Delta_2}{2}\right)}\times\\
&\qquad\times p_1^{\Delta_1-\frac{d}{2}}p_2^{\Delta_2-\frac{d}{2}}p_3^{\Delta_3-\frac{d}{2}}\int_0^\infty\mathrm{d}x\,x^{\frac{d}{2}-1}K_{\Delta_1-\frac{d}{2}}(p_1x)K_{\Delta_2-\frac{d}{2}}(p_2x)K_{\Delta_3-\frac{d}{2}}(p_3x),
\end{aligned}\tag{2.4}$$

where $K_\nu(z)$ is a Bessel K function, *i.e.*, a modified Bessel function of the second kind. A derivation of this representation may be found in appendix A.3. As mentioned in the introduction, we will generally refer to integrals of the form above featuring three Bessel K functions and a power as *triple-K* integrals. This form of the 3-point function is familiar in the context of the AdS/CFT correspondence, where every bulk-to-boundary propagator for the field dual to the conformal operator $\mathcal{O}_j$ contains one Bessel K function [37].

The expression (2.4) may be severely divergent and requires a regularisation. This stems from the fact that the original position space expression (2.1) is valid at non-coincident points only and itself requires a regularisation. A simple solution is to analytically continue (2.4) to a function of d and Δ_j, with a regularisation in these parameters then yielding a finite result.

To illustrate this, consider the case of three operators of dimension one in $d = 3$, *i.e.*, $\Delta_j = 1$, $j = 1, 2, 3$. In this case, the Bessel functions can be expressed in terms of elementary functions (see (A.4.4)) and the integral in (2.4) has a logarithmic divergence. To regularise the result, we then substitute

$$d \mapsto d + 2\epsilon, \qquad \Delta_j \mapsto \Delta_j + \epsilon. \tag{2.5}$$

This regularisation scheme is extremely useful in context of triple-K integrals since it preserves the indices of Bessel functions in (2.4). In the present case, we find

$$\begin{aligned}\langle\!\langle \mathcal{O}(\boldsymbol{p}_1)\mathcal{O}(\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle\!\rangle &= \frac{16 c_{123}\pi^3}{\Gamma\left(\frac{\epsilon}{2}\right) p_1 p_2 p_3} \cdot \int_0^\infty \mathrm{d}x\, x^{-1+\epsilon} e^{-x(p_1+p_2+p_3)} \\ &= \frac{(2\pi)^3 c_{123}}{p_1 p_2 p_3} + O(\epsilon).\end{aligned} \tag{2.6}$$

This result can be confirmed by direct calculation using the fact that

$$\int \frac{\mathrm{d}^3\boldsymbol{k}}{(2\pi)^3} \frac{1}{|\boldsymbol{k}|^2 |\boldsymbol{p}_1 - \boldsymbol{k}|^2 |\boldsymbol{p}_2 + \boldsymbol{k}|^2} = \frac{1}{8 p_1 p_2 p_3}, \tag{2.7}$$

as follows from the substitution $\tilde{\boldsymbol{k}} = \boldsymbol{k}/k^2$.

In summary then, the Fourier transform of the position space expression (2.1) for the 3-point function of scalar operators in any CFT may be expressed, at least formally, and up to an overall multiplicative constant, in terms of the triple-K integral (2.4). In the next section we will show that this representation in terms of a triple-K integral is very natural in the context of the conformal Ward identities. In fact, we will be able to re-derive the expression (2.4) by solving the conformal Ward identities directly in momentum space, without any reference to position space.

2.2 Conformal Ward identities

The conformal Ward identities (CWIs) in position space may be found in any standard reference text, *e.g.*, [36]. In momentum space, the Ward identities for scalar operators have been partially analysed in [20, 21], and we will use these results here before generalising them in the following sections. First, observe that due to Lorentz invariance any 3-point function may be expressed in terms of the magnitudes of the momenta,

$$p_j = |\boldsymbol{p}_j| = \sqrt{\boldsymbol{p}_j^2}, \qquad j = 1, 2, 3. \tag{2.8}$$

The expression (2.4) is in accord with this fact. Regarding the 3-point function as a function of the momentum magnitudes, the dilatation Ward identity then reads

$$0 = \left[2d + \sum_{j=1}^{3} \left(p_j \frac{\partial}{\partial p_j} - \Delta_j\right)\right] \langle\!\langle \mathcal{O}_1(\boldsymbol{p}_1)\mathcal{O}_2(\boldsymbol{p}_2)\mathcal{O}_3(\boldsymbol{p}_3)\rangle\!\rangle. \tag{2.9}$$

Similarly, the Ward identity associated with special conformal transformations is

$$0 = \sum_{j=1}^{3} p_j^{\kappa} \left[\frac{\partial^2}{\partial p_j^2} + \frac{d+1-2\Delta_j}{p_j} \frac{\partial}{\partial p_j} \right] \langle\!\langle \mathcal{O}_1(\boldsymbol{p}_1)\mathcal{O}_2(\boldsymbol{p}_2)\mathcal{O}_3(\boldsymbol{p}_3) \rangle\!\rangle, \tag{2.10}$$

where κ is a free Lorentz index. Choosing $\boldsymbol{p}_1$ and $\boldsymbol{p}_2$ as independent momenta, we may split this vector equation into two independent scalar equations

$$0 = \left[\left(\frac{\partial^2}{\partial p_1^2} + \frac{d+1-2\Delta_1}{p_1} \frac{\partial}{\partial p_1} \right) - \left(\frac{\partial^2}{\partial p_3^2} + \frac{d+1-2\Delta_3}{p_3} \frac{\partial}{\partial p_3} \right) \right] \langle\!\langle \mathcal{O}_1(\boldsymbol{p}_1)\mathcal{O}_2(\boldsymbol{p}_2)\mathcal{O}_3(\boldsymbol{p}_3) \rangle\!\rangle, \tag{2.11}$$

$$0 = \left[\left(\frac{\partial^2}{\partial p_2^2} + \frac{d+1-2\Delta_2}{p_2} \frac{\partial}{\partial p_2} \right) - \left(\frac{\partial^2}{\partial p_3^2} + \frac{d+1-2\Delta_3}{p_3} \frac{\partial}{\partial p_3} \right) \right] \langle\!\langle \mathcal{O}_1(\boldsymbol{p}_1)\mathcal{O}_2(\boldsymbol{p}_2)\mathcal{O}_3(\boldsymbol{p}_3) \rangle\!\rangle. \tag{2.12}$$

As an immediate check, we may verify that the expression (2.4) satisfies (2.11, 2.12) using the well-known Bessel function relations [38]

$$\frac{\partial}{\partial a} \left[a^{\nu} K_{\nu}(ax) \right] = -x a^{\nu} K_{\nu-1}(ax), \tag{2.13}$$

$$K_{\nu-1}(x) + \frac{2\nu}{x} K_{\nu}(x) = K_{\nu+1}(x). \tag{2.14}$$

As we will see shortly, equations of the form (2.11, 2.12) also arise in the case of 3-point correlators of general tensor operators.

2.3 Uniqueness of the solution

To frame our analysis purely in momentum space, we need to show that there is a unique physically acceptable solution, up to an overall multiplicative constant, of the system (2.9, 2.11, 2.12) of dilatation and special CWIs. To accomplish this, it suffices to transform these equations into generalised hypergeometric form by writing

$$\langle\!\langle \mathcal{O}_1(\boldsymbol{p}_1)\mathcal{O}_2(\boldsymbol{p}_2)\mathcal{O}_3(\boldsymbol{p}_3) \rangle\!\rangle = p_3^{\Delta_t - 2d} \left(\frac{p_1^2}{p_3^2} \right)^{\mu} \left(\frac{p_2^2}{p_3^2} \right)^{\lambda} F\left(\frac{p_1^2}{p_3^2}, \frac{p_2^2}{p_3^2} \right), \tag{2.15}$$

where the overall power of momenta on the right-hand side is fixed by the dilatation Ward identity (2.9), and we have chosen to multiply the arbitrary function F by the prefactor $(p_1^2/p_3^2)^{\mu}(p_2^2/p_3^2)^{\lambda}$, where μ and λ are arbitrary constants. Substituting this parametrisation into (2.11, 2.12) then yields a pair of differential equations satisfied by F. Taking μ and λ to be any of the four combinations obtainable from the values

$$\mu = 0,\ \Delta_1 - \frac{d}{2}, \qquad\qquad \lambda = 0,\ \Delta_2 - \frac{d}{2}, \tag{2.16}$$

these equations for F read

$$\begin{aligned} 0 = \Big[& \xi(1-\xi)\frac{\partial^2}{\partial \xi^2} - \eta^2 \frac{\partial^2}{\partial \eta^2} - 2\xi\eta \frac{\partial^2}{\partial \xi \partial \eta} \\ & + \left(\gamma - (\alpha+\beta+1)\xi \right) \frac{\partial}{\partial \xi} - (\alpha+\beta+1)\eta \frac{\partial}{\partial \eta} - \alpha\beta \Big] F(\xi,\eta), \end{aligned} \tag{2.17}$$

$$\begin{aligned} 0 = \Big[& \eta(1-\eta)\frac{\partial^2}{\partial \eta^2} - \xi^2 \frac{\partial^2}{\partial \xi^2} - 2\xi\eta \frac{\partial^2}{\partial \xi \partial \eta} \\ & + \left(\gamma' - (\alpha+\beta+1)\eta \right) \frac{\partial}{\partial \eta} - (\alpha+\beta+1)\xi \frac{\partial}{\partial \xi} - \alpha\beta \Big] F(\xi,\eta), \end{aligned} \tag{2.18}$$

where

$$\xi = \frac{p_1^2}{p_3^2}, \qquad \eta = \frac{p_2^2}{p_3^2}, \tag{2.19}$$

and the values of the parameters $\alpha, \beta, \gamma, \gamma'$ depend on the choice of μ and λ. Specifically, parametrising the four choices for μ and λ by two variables $\epsilon_1, \epsilon_2 \in \{-1, +1\}$ according to

$$\mu = \frac{1}{2}\big(\Delta_1 - \frac{d}{2}\big)(\epsilon_1 + 1), \qquad \lambda = \frac{1}{2}\big(\Delta_2 - \frac{d}{2}\big)(\epsilon_2 + 1), \tag{2.20}$$

we have

$$\begin{aligned} \alpha &= \frac{1}{2}\left[\epsilon_1\left(\Delta_1 - \frac{d}{2}\right) + \epsilon_2\left(\Delta_2 - \frac{d}{2}\right) + \Delta_3\right], \qquad \beta = \alpha - \left(\Delta_3 - \frac{d}{2}\right), \\ \gamma &= 1 + \epsilon_1\left(\Delta_1 - \frac{d}{2}\right), \qquad \gamma' = 1 + \epsilon_2\left(\Delta_2 - \frac{d}{2}\right). \end{aligned} \tag{2.21}$$

The system of equations (2.17, 2.18) defines the generalised hypergeometric function of two variables Appell F_4. This function has been extensively studied by mathematicians (see, *e.g.*, [39, 40]), and its important properties are summarised in appendix A.5. In particular, the system of equations (2.17, 2.18) has at most four linearly independent solutions, each of which may be expressed in terms of the F_4 function [40, 41]. The four possible choices for μ and λ reproduce these four solutions exactly.

In a physical context only one linear combination of these four solutions is acceptable: all the others contain divergences for collinear momentum configurations, for example when $p_1 + p_2 = p_3$. To see this, consider the integral representation [42]

$$\begin{aligned} F_4\left(\alpha, \beta; \gamma, \gamma'; \frac{p_1^2}{p_3^2}, \frac{p_2^2}{p_3^2}\right) &= \frac{\Gamma(\gamma)\Gamma(\gamma')}{2^{\alpha+\beta-\gamma-\gamma'}\Gamma(\alpha)\Gamma(\beta)} \cdot \frac{p_3^{\alpha+\beta}}{p_1^{\gamma-1} p_2^{\gamma'-1}} \times \\ &\times \int_0^\infty \mathrm{d}x x^{\alpha+\beta-\gamma-\gamma'+1} I_{\gamma-1}(p_1 x) I_{\gamma'-1}(p_2 x) K_{\beta-\alpha}(p_3 x), \end{aligned} \tag{2.22}$$

where $I_\nu(x)$ is the Bessel I function. This expression is formal in the sense that the integral converges only for α, β, γ, γ' in certain ranges, see appendix A.5 for details. For the remaining parameter values the integral is defined by the analytic continuation (2.5). Using (2.21), one can then write the four solutions for the 3-point functions in the form

$$p_1^{\Delta_1 - \frac{d}{2}} p_2^{\Delta_2 - \frac{d}{2}} p_3^{\Delta_3 - \frac{d}{2}} \int_0^\infty \mathrm{d}x\, x^{\frac{d}{2}-1} I_{\pm(\Delta_1 - \frac{d}{2})}(p_1 x) I_{\pm(\Delta_2 - \frac{d}{2})}(p_2 x) K_{\Delta_3 - \frac{d}{2}}(p_3 x). \tag{2.23}$$

For large x we have the asymptotic expansions

$$I_\nu(x) = \frac{1}{\sqrt{2\pi}} \frac{e^x}{\sqrt{x}} + \ldots, \qquad K_\nu(x) = \sqrt{\frac{\pi}{2}} \frac{e^{-x}}{\sqrt{x}} + \ldots, \tag{2.24}$$

from which we see that the integral (2.23) converges at infinite x only for non-triangle (*i.e.*, unphysical) momentum configurations where $p_1 + p_2 < p_3$. Moreover, for the physical collinear momentum configuration $p_1 + p_2 = p_3$, the integral diverges for dimensions $d \geq 3$. However, the 3-point function itself is a linear combination of these four solutions and may be continued to the physical regime by choosing the linear combination for which the collinear divergences cancel. This may be accomplished by subtracting two integrals with the same asymptotics, *i.e.*, we sum the four terms of the form (2.23) with signs chosen so as to obtain Bessel K functions

$$K_\nu(x) = \frac{\pi}{2\sin(\nu\pi)} \left[I_\nu(x) - I_{-\nu}(x)\right]. \tag{2.25}$$

Therefore we arrive at the unique solution

$$\langle\!\langle \mathcal{O}_1(\boldsymbol{p}_1)\mathcal{O}_2(\boldsymbol{p}_2)\mathcal{O}_3(\boldsymbol{p}_3)\rangle\!\rangle = C_{123}\cdot p_1^{\Delta_1-\frac{d}{2}}p_2^{\Delta_2-\frac{d}{2}}p_3^{\Delta_3-\frac{d}{2}}\int_0^\infty \mathrm{d}x\, x^{\frac{d}{2}-1}K_{\Delta_1-\frac{d}{2}}(p_1x)K_{\Delta_2-\frac{d}{2}}(p_2x)K_{\Delta_3-\frac{d}{2}}(p_3x), \quad (2.26)$$

where C_{123} is an overall undetermined constant. From the asymptotic expansion (2.24), it is clear that this triple-K integral converges at infinite x for physical momentum configurations $p_1+p_2+p_3>0$. Depending on the values of the parameters Δ_j and d, however, the triple-K integral may still diverge at $x=0$. This divergence may be regularised using (2.5) as we will discuss in the next section.

In summary then, we have shown that the conformal Ward identities may be solved directly in momentum space leading to a unique result (2.26). As we will see shortly, a similar procedure also holds for tensorial correlation functions: solving the momentum space Ward identities will lead to a unique solution for 3-point correlators without any reference to the position space analysis.

2.4 Region of validity and anomalies

In this section, we now discuss the regularisation of the potential divergence of the triple-K integral at $x=0$. In general, assuming all parameters and variables are real, the triple-K integral (2.26) converges for [42]

$$\frac{d}{2} > \sum_{j=1}^{3}\left|\Delta_j-\frac{d}{2}\right|+2, \qquad p_1,p_2,p_3>0. \quad (2.27)$$

If the parameters in the integral do not satisfy this inequality, however, the integral may be defined via analytic continuation in d and Δ_j. If for some set of parameters the integral exhibits a singularity, then a regularisation is necessary and the scheme (2.5) can be used.

Let us consider a concrete example, also discussed in [20]. We set $d=3$ and consider three scalar operators of dimensions $\Delta_j=2$, $j=1,2,3$. The triple-K integral is then logarithmically divergent and a regularisation is necessary. We obtain

$$\begin{aligned}\langle\!\langle \mathcal{O}(\boldsymbol{p}_1)\mathcal{O}(\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle\!\rangle &= C_{123}\left(\frac{\pi}{2}\right)^{3/2}\cdot\int_0^\infty \mathrm{d}x\, x^{-1+\epsilon}e^{-x(p_1+p_2+p_3)}\\ &= C_{123}\left(\frac{\pi}{2}\right)^{3/2}\Gamma(\epsilon)(p_1+p_2+p_3)^{-\epsilon}\\ &= C_{123}\left(\frac{\pi}{2}\right)^{3/2}\left[\frac{1}{\epsilon}-(\gamma_E+\log(p_1+p_2+p_3))+O(\epsilon)\right]. \end{aligned} \quad (2.28)$$

The first two terms are proportional to the Fourier transform of $\delta(\boldsymbol{x}_1-\boldsymbol{x}_3)\delta(\boldsymbol{x}_2-\boldsymbol{x}_3)$ and may be removed by adding local counterterms. In the regularisation scheme (2.5), such a counterterm has the form

$$S_{ct} = c_\epsilon\int \mathrm{d}^{3+2\epsilon}\boldsymbol{x}\;\phi_0^3\mu^{-\epsilon}, \quad (2.29)$$

where μ is a scale that we introduce (as usual) so that the action is dimensionless. By taking three functional derivatives with respect to the source, we find the contribution of this term to the 3-point function is

$$3!c_\epsilon\mu^{-\epsilon}\delta(\boldsymbol{x}_1-\boldsymbol{x}_3)\delta(\boldsymbol{x}_2-\boldsymbol{x}_3). \quad (2.30)$$

Therefore, in order to cancel the divergence in (2.28), we need to choose

$$c_\epsilon = -\frac{C_{123}}{3!\epsilon}\left(\frac{\pi}{2}\right)^{3/2}+O(\epsilon^0). \quad (2.31)$$

Let us make a few comments. First, we can choose the subleading terms in c_ϵ in such a way that the constant in (2.28) can attain an arbitrary value. This is just scheme dependence. Secondly, notice that the same value of c_ϵ and the same factor of $\mu^{-\epsilon}$ in (2.29) are obtained by standard dimensional regularisation $d \mapsto d - \epsilon$. This observation will be a key point in a relation between regularisations of 2- and 3-point functions discussed in section 5.3.4.

With the renormalisation procedure carried out, we are then left with the finite result

$$\langle\!\langle \mathcal{O}(\boldsymbol{p}_1)\mathcal{O}(\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle\!\rangle = -C_{123}\left(\frac{\pi}{2}\right)^{3/2}\log\frac{p_1+p_2+p_3}{\tilde{\mu}}, \tag{2.32}$$

where

$$\tilde{\mu} = \mu\exp\left[\frac{c_\epsilon^{(0)}}{C_{123}}\left(\frac{2}{\pi}\right)^{3/2}\right] \tag{2.33}$$

and $c_\epsilon^{(0)}$ denotes the first subleading term in c_ϵ. Note that this 3-point function is anomalous, *i.e.*, it does not satisfy the dilatation Ward identity (2.9) and the scale dependence can be extracted from the μ dependence of the counterterm (2.30),

$$\mu\frac{\partial}{\partial\mu}\langle\!\langle \mathcal{O}(\boldsymbol{p}_1)\mathcal{O}(\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle\!\rangle = C_{123}\left(\frac{\pi}{2}\right)^{3/2} + O(\epsilon). \tag{2.34}$$

Of course, this can also be obtained directly from (2.32).

The violation of scale invariance means that the trace Ward identity is anomalous,

$$\langle T\rangle = -\phi_0\langle\mathcal{O}\rangle + \frac{1}{3!}C_{123}\left(\frac{\pi}{2}\right)^{3/2}\phi_0^3. \tag{2.35}$$

This in turn implies that the 4-point function of the stress-energy tensor with three $\mathcal{O}$s contains a specific ultra-local term that is not scheme-dependent because it is fixed by the anomalous Ward identity (2.35).

3 Decomposition of tensors

In this section, we present a natural decomposition of tensorial correlation functions. Correlation functions of conserved currents are transverse and/or traceless – up to local terms – and we would like to find a decomposition which reflects these properties. At this point, we will not yet impose conformal invariance.

The problem of decomposition has already been tackled in a number of papers, see for example [43, 30, 26, 27, 28, 13]. The usual approach consists of writing down the most general tensor structure before imposing the constraints following from symmetries and Ward identities. Here we refine this approach to take account of the permutation symmetries of operator insertions inside correlators, obtaining a convenient and natural decomposition applicable for any correlation function. In particular, our decomposition contains the minimal number of tensor structures, leading to the simplest form for the conformal Ward identities.

We remind the reader we will always be working in d-dimensional Euclidean field theory with a flat metric $\delta_{\mu\nu}$ for which indices are raised and lowered trivially.

3.1 Representations of tensor structures

The operator

$$\pi^\mu_\alpha(\boldsymbol{p}) = \delta^\mu_\alpha - \frac{p^\mu p_\alpha}{p^2} \tag{3.1}$$

is a projector onto tensors transverse to $\boldsymbol{p}$, *i.e.*, $p_\mu\pi^\mu_\alpha(\boldsymbol{p}) = 0$. Similarly, in d dimensions, the operator

$$\Pi^{\mu\nu}_{\alpha\beta}(\boldsymbol{p}) = \frac{1}{2}\left(\pi^\mu_\alpha(\boldsymbol{p})\pi^\nu_\beta(\boldsymbol{p}) + \pi^\mu_\beta(\boldsymbol{p})\pi^\nu_\alpha(\boldsymbol{p})\right) - \frac{1}{d-1}\pi^{\mu\nu}(\boldsymbol{p})\pi_{\alpha\beta}(\boldsymbol{p}) \tag{3.2}$$

is a projector onto transverse to $\boldsymbol{p}$, traceless, symmetric tensors of rank two. In particular

$$\delta_{\mu\nu}\Pi^{\mu\nu}_{\alpha\beta}(\boldsymbol{p}) = 0, \qquad p_\mu \Pi^{\mu\nu}_{\alpha\beta}(\boldsymbol{p}) = 0. \tag{3.3}$$

Therefore, any transverse to $\boldsymbol{p}$, traceless, symmetric tensor $t^{\mu\nu}$ of rank two may be written as $t^{\mu\nu} = \Pi^{\mu\nu}_{\alpha\beta}(\boldsymbol{p}) X^{\alpha\beta}$, where $X^{\alpha\beta}$ is an arbitrary tensor.

As we are interested in correlation functions, we must consider tensor functions that depend on a number of momenta. Let $\mathcal{T}_j^{\mu_{j1}\mu_{j2}\dots\mu_{jr_j}}$, $j = 1, 2, \dots, n$ be a given set of QFT operators. Due to the momentum conservation, the n-point function contains a delta function which may be written explicitly by introducing the reduced matrix element which we denote with double brackets $\langle\!\langle \dots \rangle\!\rangle$,

$$\begin{aligned}
&\langle \mathcal{T}_1^{\mu_{11}\mu_{12}\dots\mu_{1r_1}}(\boldsymbol{p}_1)\mathcal{T}_2^{\mu_{21}\mu_{22}\dots\mu_{2r_2}}(\boldsymbol{p}_2)\dots\mathcal{T}_n^{\mu_{n1}\mu_{n2}\dots\mu_{nr_n}}(\boldsymbol{p}_n)\rangle \\
&= (2\pi)^d \delta\left(\sum_{k=1}^n \boldsymbol{p}_k\right) \langle\!\langle \mathcal{T}_1^{\mu_{11}\mu_{12}\dots\mu_{1r_1}}(\boldsymbol{p}_1)\mathcal{T}_2^{\mu_{21}\mu_{22}\dots\mu_{2r_2}}(\boldsymbol{p}_2)\dots\mathcal{T}_n^{\mu_{n1}\mu_{n2}\dots\mu_{nr_n}}(\boldsymbol{p}_n)\rangle\!\rangle.
\end{aligned} \tag{3.4}$$

The n-point function thus depends on at most $n-1$ vectors, say $\boldsymbol{p}_1, \dots, \boldsymbol{p}_{n-1}$. If $n-1 < d$, then all $n-1$ momenta are independent. If $n-1 \geq d$, however, then only d generic momenta are independent. In this case we can write [44]

$$\delta^{\mu\nu} = \sum_{j,k=1}^d p_j^\mu p_k^\nu (Z^{-1})_{kj}, \tag{3.5}$$

where Z is the Gram matrix, $Z = [\boldsymbol{p}_k \cdot \boldsymbol{p}_l]_{k,l=1}^d$, hence the metric $\delta^{\mu\nu}$ is no longer an independent tensor.

From now on we assume $d \geq 3$. Since we are primarily interested in 3-point functions, the degeneracy does not occur. Nevertheless, the case $d = 3$ is still special since the existence of the cross-product allows the metric tensor to be re-expressed purely in terms of the momenta. This degeneracy serves to reduce the number of independent form factors for certain correlators, as we discuss in appendix A.2. In the following discussion we will ignore this degeneracy however and concentrate on the general case. We will therefore choose two out of the three $\boldsymbol{p}_1, \boldsymbol{p}_2, \boldsymbol{p}_3$ as independent momenta, and treat the metric $\delta^{\mu\nu}$ as an independent tensor.

As an example consider a 3-point function of two transverse, traceless, symmetric rank two operators $t^{\mu\nu}$ and a scalar operator $\mathcal{O}$. Using the projectors (3.2) one can write the most general form

$$\langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1) t^{\mu_2\nu_2}(\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle\!\rangle = \Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1)\Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2) X^{\alpha_1\beta_1\alpha_2\beta_2}, \tag{3.6}$$

where $X^{\alpha_1\beta_1\alpha_2\beta_2}$ is a general tensor of rank four built from the metric and momenta. Usually one chooses two independent momenta once and for all. On the other hand, there is no obstacle to choosing different independent momenta for different Lorentz indices. In this paper we always choose

$$\boldsymbol{p}_1, \boldsymbol{p}_2 \text{ for } \mu_1, \nu_1; \ \boldsymbol{p}_2, \boldsymbol{p}_3 \text{ for } \mu_2, \nu_2 \text{ and } \boldsymbol{p}_3, \boldsymbol{p}_1 \text{ for } \mu_3, \nu_3. \tag{3.7}$$

Such a choice enhances the symmetry properties of the decomposition, as we will discuss at length in the next section.

Let us now enumerate all possible tensors that can appear in $X^{\alpha_1\beta_1\alpha_2\beta_2}$. Observe that whenever a simple tensor contains at least one of the following tensors

$$\delta^{\alpha_1\beta_1}, \delta^{\alpha_2\beta_2}, p_1^{\alpha_1}, p_1^{\beta_1}, p_2^{\alpha_2}, p_2^{\beta_2}, \tag{3.8}$$

then the contraction with the projectors in (3.6) vanishes. Therefore, in accordance with the choice (3.7), the only tensors giving a non-zero result after contraction with the projectors are

$$\delta^{\alpha_1\alpha_2}, \delta^{\alpha_1\beta_2}, \delta^{\beta_1\alpha_2}, \delta^{\beta_1\beta_2}, p_2^{\alpha_1}, p_2^{\beta_1}, p_3^{\alpha_2}, p_3^{\beta_2}. \tag{3.9}$$

Since the projector (3.2) is symmetric in $\mu \leftrightarrow \nu$ and $\alpha \leftrightarrow \beta$, the most general form of our 3-point function is then

$$\langle\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1)t^{\mu_2\nu_2}(\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle\rangle = \Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1)\Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)\Big[A_1 p_2^{\alpha_1}p_2^{\beta_1}p_3^{\alpha_2}p_3^{\beta_2} + A_2 p_2^{\alpha_1}p_3^{\alpha_2}\delta^{\beta_1\beta_2} + A_3\delta^{\alpha_1\alpha_2}\delta^{\beta_1\beta_2}\Big], \tag{3.10}$$

where the coefficients A_1, A_2 and A_3 are scalar functions of momenta. We will refer to the coefficients A_j, and their analogous counterparts in more general correlation functions, as *form factors*. By Lorentz invariance, these form factors are functions of the momentum magnitudes

$$p_j = |\boldsymbol{p}_j| = \sqrt{\boldsymbol{p}_j^2}, \qquad j = 1, 2, 3, \tag{3.11}$$

i.e., $A_j = A_j(p_1, p_2, p_3)$. In particular, any scalar product of two momenta can be written as a combination of momentum magnitudes, for example

$$\boldsymbol{p}_1 \cdot \boldsymbol{p}_2 = \frac{1}{2}(p_3^2 - p_1^2 - p_2^2). \tag{3.12}$$

For brevity, we will henceforth suppress the dependence of form factors on the momentum magnitudes, writing $A_j(p_1, p_2, p_3)$ as simply A_j.

Note that the correlation function on the left-hand side of (3.10) is symmetric under a transposition $(\boldsymbol{p}_1, \mu_1, \nu_1) \leftrightarrow (\boldsymbol{p}_2, \mu_2, \nu_2)$. One can apply this symmetry to the right-hand side to find that all form factors A_1, A_2 and A_3 are symmetric under $p_1 \leftrightarrow p_2$. To prove this, observe that one has, for example, $\pi^{\mu_1}_{\alpha_1}(\boldsymbol{p}_1)p_3^{\alpha_1} = -\pi^{\mu_1}_{\alpha_1}(\boldsymbol{p}_1)p_2^{\alpha_1}$. Therefore $\boldsymbol{p}_2$ and $-\boldsymbol{p}_3$ can be exchanged under both $\pi^{\mu_1}_{\alpha_1}(\boldsymbol{p}_1)$ and $\Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1)$, and similarly for other momenta.

For any form factor A_j we define an associated non-negative integer N_j, the *tensorial dimension* of A_j, similar to that defined in [30]. Specifically, the tensorial dimension N_j is the number of momenta that appear in the tensorial expression multiplying A_j (excluding those in the transverse-traceless projectors) in the decomposition of the correlation function. As we will see later, this quantity will appear explicitly in the conformal Ward identities. In the example (3.10), we have the following tensorial dimensions: $N_1 = 4$, $N_2 = 2$ and $N_3 = 0$.

Decompositions for other correlation functions may be found in the second part of the paper. Observe that in each case the form factor A_1 stands in front of the unique tensor containing momenta only. The tensorial dimension N_1 is therefore always equal to the number of Lorentz indices in the 3-point function, and tensorial dimensions of all remaining form factors are smaller than N_1.

3.2 Decomposition of $\langle t^{\mu_1\nu_1}t^{\mu_2\nu_2}t^{\mu_3\nu_3}\rangle$

In the previous section we introduced a natural decomposition of tensor structures. Rather than fixing two independent momenta (as is done for example in [43, 30, 26, 27, 28, 13]) we chose a different set of independent momenta for different Lorentz indices according to (3.7). Such a choice respects all symmetries of the correlation function, as we now discuss.

In [30], it was shown that the transverse-traceless correlation function $\langle\langle t^{\mu_1\nu_1}t^{\mu_2\nu_2}t^{\mu_3\nu_3}\rangle\rangle$ can be decomposed into eight tensor structures plus their $\boldsymbol{p}_1 \leftrightarrow \boldsymbol{p}_2$ symmetric versions. In our method, however, we arrive at only five tensor structures (for the general case $d \geq 3$, see

appendix A.2 for the case $d = 3$) according to the following decomposition

$$\begin{aligned}
&\langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1)t^{\mu_2\nu_2}(\boldsymbol{p}_2)t^{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\!\rangle \\
&\quad = \Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1)\Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)\Pi^{\mu_3\nu_3}_{\alpha_3\beta_3}(\boldsymbol{p}_3)\Big[A_1 p_2^{\alpha_1}p_2^{\beta_1}p_3^{\alpha_2}p_3^{\beta_2}p_1^{\alpha_3}p_1^{\beta_3} \\
&\qquad + A_2\delta^{\beta_1\beta_2}p_2^{\alpha_1}p_3^{\alpha_2}p_1^{\alpha_3}p_1^{\beta_3} + A_2(p_1\leftrightarrow p_3)\delta^{\beta_2\beta_3}p_2^{\alpha_1}p_2^{\beta_1}p_3^{\alpha_2}p_1^{\alpha_3} \\
&\qquad\quad + A_2(p_2\leftrightarrow p_3)\delta^{\beta_1\beta_3}p_2^{\alpha_1}p_3^{\alpha_2}p_3^{\beta_2}p_1^{\alpha_3} \\
&\qquad + A_3\delta^{\alpha_1\alpha_2}\delta^{\beta_1\beta_2}p_1^{\alpha_3}p_1^{\beta_3} + A_3(p_1\leftrightarrow p_3)\delta^{\alpha_2\alpha_3}\delta^{\beta_2\beta_3}p_2^{\alpha_1}p_2^{\beta_1} \\
&\qquad\quad + A_3(p_2\leftrightarrow p_3)\delta^{\alpha_1\alpha_3}\delta^{\beta_1\beta_3}p_3^{\alpha_2}p_3^{\beta_2} \\
&\qquad + A_4\delta^{\alpha_1\alpha_3}\delta^{\alpha_2\beta_3}p_2^{\beta_1}p_3^{\beta_2} + A_4(p_1\leftrightarrow p_3)\delta^{\alpha_1\alpha_3}\delta^{\alpha_2\beta_1}p_3^{\beta_2}p_1^{\beta_3} \\
&\qquad\quad + A_4(p_2\leftrightarrow p_3)\delta^{\alpha_1\alpha_2}\delta^{\alpha_3\beta_2}p_2^{\beta_1}p_1^{\beta_3} \\
&\qquad + A_5\delta^{\alpha_1\beta_2}\delta^{\alpha_2\beta_3}\delta^{\alpha_3\beta_1}\Big]\,.
\end{aligned} \tag{3.13}$$

By $p_1 \leftrightarrow p_3$ we denote the exchange of the arguments p_1 and p_3, $A_2(p_1 \leftrightarrow p_3) = A_2(p_3, p_2, p_1)$. If no arguments are specified, then the standard ordering is assumed, *i.e.*, $A_2 = A_2(p_1, p_2, p_3)$.

First observe that this decomposition is manifestly invariant under the permutation group S_3 of the set $\{1, 2, 3\}$, *i.e.*, for any $\sigma \in S_3$,

$$\langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1)t^{\mu_2\nu_2}(\boldsymbol{p}_2)t^{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\!\rangle = \langle\!\langle t^{\mu_{\sigma(1)}\nu_{\sigma(1)}}(\boldsymbol{p}_{\sigma(1)})t^{\mu_{\sigma(2)}\nu_{\sigma(2)}}(\boldsymbol{p}_{\sigma(2)})t^{\mu_{\sigma(3)}\nu_{\sigma(3)}}(\boldsymbol{p}_{\sigma(3)})\rangle\!\rangle. \tag{3.14}$$

In particular, the form factors A_1 and A_5 are S_3-invariant,

$$A_j(p_1, p_2, p_3) = A_j(p_{\sigma(1)}, p_{\sigma(2)}, p_{\sigma(3)}), \quad j \in \{1, 5\}, \tag{3.15}$$

since the tensors they multiply are S_3-invariant. The action of the symmetry group on the remaining terms is then clearly visible. As an example, let us concentrate on the third line of (3.13) with the A_2 form factor. The $(\boldsymbol{p}_1, \mu_1, \nu_1) \leftrightarrow (\boldsymbol{p}_2, \mu_2, \nu_2)$ permutation leaves the tensor in the first term invariant, therefore the A_2 factor exhibits the $p_1 \leftrightarrow p_2$ symmetry. On the other hand, the $(\boldsymbol{p}_1, \mu_1, \nu_1) \leftrightarrow (\boldsymbol{p}_3, \mu_3, \nu_3)$ permutation swaps tensor structures of the first and the second term in the third line. This requires that the form factor of the second term is related to the form factor of the first term by the $p_1 \leftrightarrow p_3$ permutation, as indicated. Working out the remaining lines of (3.13) one finds that both remaining factors A_3 and A_4 are $p_1 \leftrightarrow p_2$ symmetric.

Let us comment then on the apparent disagreement between the number of tensor structures between (3.13) and the results of [30]. As already mentioned, the mismatch follows from the choice of two independent momenta in [30] to be $\boldsymbol{p}_1$ and $\boldsymbol{p}_2$, in our notation. Such a choice breaks the S_3 symmetry down to the $(\boldsymbol{p}_1, \mu_1, \nu_1) \leftrightarrow (\boldsymbol{p}_2, \mu_2, \nu_2)$ symmetry. One can easily recover eight tensor structures from (3.13) by substituting $\boldsymbol{p}_3 = -\boldsymbol{p}_1 - \boldsymbol{p}_2$ and writing the decomposition in terms of $\boldsymbol{p}_1$ and $\boldsymbol{p}_2$ only. One finds

$$\begin{aligned}
&\langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1)t^{\mu_2\nu_2}(\boldsymbol{p}_2)t^{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\!\rangle \\
&\quad = \Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1)\Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)\Pi^{\mu_3\nu_3}_{\alpha_3\beta_3}(\boldsymbol{p}_3)\left[\frac{1}{2}A_1 p_2^{\alpha_1}p_2^{\beta_1}p_1^{\alpha_2}p_1^{\beta_2}p_1^{\alpha_3}p_1^{\beta_3}\right. \\
&\qquad - \frac{1}{2}A_2\delta^{\beta_1\beta_2}p_2^{\alpha_1}p_1^{\alpha_2}p_1^{\alpha_3}p_1^{\beta_3} - A_2(p_1\leftrightarrow p_3)\delta^{\beta_2\beta_3}p_2^{\alpha_1}p_2^{\beta_1}p_1^{\alpha_2}p_1^{\alpha_3} \\
&\qquad + \frac{1}{2}A_3\delta^{\alpha_1\alpha_2}\delta^{\beta_1\beta_2}p_1^{\alpha_3}p_1^{\beta_3} + A_3(p_1\leftrightarrow p_3)\delta^{\alpha_2\alpha_3}\delta^{\beta_2\beta_3}p_2^{\alpha_1}p_2^{\beta_1} \\
&\qquad - \frac{1}{2}A_4\delta^{\alpha_1\alpha_3}\delta^{\alpha_2\beta_3}p_2^{\beta_1}p_1^{\beta_2} - A_4(p_1\leftrightarrow p_3)\delta^{\alpha_1\alpha_3}\delta^{\alpha_2\beta_1}p_1^{\beta_2}p_1^{\beta_3} \\
&\qquad + \frac{1}{2}A_5\delta^{\alpha_1\beta_2}\delta^{\alpha_2\beta_3}\delta^{\alpha_3\beta_1} \\
&\qquad \left. + \text{everything with } (\boldsymbol{p}_1, \alpha_1, \beta_1) \leftrightarrow (\boldsymbol{p}_2, \alpha_2, \beta_2)\right]\,.
\end{aligned} \tag{3.16}$$

As we can see, the number of tensor structures increases to exactly eight, as the symmetry group is diminished.

Summarising, our decomposition method based on (3.7) gives the minimal number of tensor structures obeying the symmetries of the correlation function. It is an improvement over the standard method with two independent momenta fixed, since such a choice breaks symmetries and leads therefore to the larger number of tensor structures.

Finally, we should comment on the fact that we decompose the transverse-traceless part of the correlation function only. This is because the difference between the full 3-point function and its transverse-traceless part is semi-local, and hence may be entirely reconstructed from the Ward identities. We will discuss this method for recovering the full correlation function from its transverse-traceless part in the next section.

Let us note in passing that the decomposition method described here may also be used for correlation functions in non-conformal theories. For example, in cases where the stress-energy tensor is transverse but no longer traceless one should use the π^μ_α projectors (3.1) in place of $\Pi^{\mu\nu}_{\alpha\beta}$ in (3.13). In this way one obtains ten tensor structures, five of which have nonzero trace. This decomposition is given in appendix A.1.

3.3 Finding the form factors

We would like to apply the results of the previous section to spin-1 and spin-2 conserved currents J^μ and a stress-energy tensor $T^{\mu\nu}$. These quantum operators, however, are only transverse and traceless on-shell, and in the quantum case, we need to analyse Ward identities. To proceed, we define transverse, transverse-traceless and local parts of J^μ and $T^{\mu\nu}$ by

$$j^\mu \equiv \pi^\mu_\alpha J^\alpha, \qquad j^\mu_{\text{loc}} \equiv J^\mu - j^\mu, \tag{3.17}$$

$$t^{\mu\nu} \equiv \Pi^{\mu\nu}_{\alpha\beta} T^{\alpha\beta}, \qquad t^{\mu\nu}_{\text{loc}} \equiv T^{\mu\nu} - t^{\mu\nu}, \tag{3.18}$$

as well as longitudinal and trace parts

$$r = p_\mu J^\mu, \qquad R^\nu = p_\mu T^{\mu\nu}, \qquad R = p_\nu R^\nu, \qquad T = T^\mu_\mu. \tag{3.19}$$

From these definitions, we then have

$$j^\mu_{\text{loc}} = \frac{p^\mu}{p^2} r, \tag{3.20}$$

$$\begin{aligned} t^{\mu\nu}_{\text{loc}} &= \frac{p^\mu}{p^2} R^\nu + \frac{p^\nu}{p^2} R^\mu - \frac{p^\mu p^\nu}{p^4} R + \frac{1}{d-1} \pi^{\mu\nu} \left(T - \frac{R}{p^2} \right) \\ &= \mathscr{T}^{\mu\nu}_\alpha R^\alpha + \frac{\pi^{\mu\nu}}{d-1} T, \end{aligned} \tag{3.21}$$

where the operator

$$\mathscr{T}^{\mu\nu}_\alpha(\boldsymbol{p}) = \frac{1}{p^2} \left[2p^{(\mu} \delta^{\nu)}_\alpha - \frac{p_\alpha}{d-1} \left(\delta^{\mu\nu} + (d-2) \frac{p^\mu p^\nu}{p^2} \right) \right]. \tag{3.22}$$

In the following, we will also use $\mathscr{T}^{\mu\nu\alpha} = \delta^{\alpha\beta} \mathscr{T}^{\mu\nu}_\beta$.

We now observe that in a CFT, all terms in (3.20) and (3.21) are computable by means of the transverse and trace Ward identities. We can therefore divide a 3-point function into two parts: the *transverse-traceless part* represented as in section 3.1, and the *semi-local part* (indicated by the subscript *loc*) expressible through the transverse Ward identities. For simplicity we will use the term 'transverse-traceless part' in all cases, even if the correlation function does not contain the stress-energy tensor.

As an example, consider

$$\langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1) t^{\mu_2\nu_2}(\boldsymbol{p}_2) \mathcal{O}(\boldsymbol{p}_3) \rangle\!\rangle = \Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1) \Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2) \langle\!\langle T^{\alpha_1\beta_1}(\boldsymbol{p}_1) T^{\alpha_2\beta_2}(\boldsymbol{p}_2) \mathcal{O}(\boldsymbol{p}_3) \rangle\!\rangle. \tag{3.23}$$

One can recover the full 3-point function by writing

$$\begin{aligned}\langle\!\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}\mathcal{O}\rangle\!\rangle &= \langle\!\langle t^{\mu_1\nu_1}t^{\mu_2\nu_2}\mathcal{O}\rangle\!\rangle + \langle\!\langle t^{\mu_1\nu_1}t_{\text{loc}}^{\mu_2\nu_2}\mathcal{O}\rangle\!\rangle + \langle\!\langle t_{\text{loc}}^{\mu_1\nu_1}t^{\mu_2\nu_2}\mathcal{O}\rangle\!\rangle + \langle\!\langle t_{\text{loc}}^{\mu_1\nu_1}t_{\text{loc}}^{\mu_2\nu_2}\mathcal{O}\rangle\!\rangle \\ &= \langle\!\langle t^{\mu_1\nu_1}t^{\mu_2\nu_2}\mathcal{O}\rangle\!\rangle - \langle\!\langle T^{\mu_1\nu_1}t_{\text{loc}}^{\mu_2\nu_2}\mathcal{O}\rangle\!\rangle - \langle\!\langle t_{\text{loc}}^{\mu_1\nu_1}T^{\mu_2\nu_2}\mathcal{O}\rangle\!\rangle + \langle\!\langle t_{\text{loc}}^{\mu_1\nu_1}t_{\text{loc}}^{\mu_2\nu_2}\mathcal{O}\rangle\!\rangle \end{aligned} \tag{3.24}$$

All terms on the right-hand side apart from the first may be computed by means of Ward identities. The exact form of the Ward identities depends on the exact definition of the operators involved, but more importantly, all these terms depend on 2-point functions only.

Due to the complicated nature of contractions of the projectors (3.1) and (3.2) one might fear that it is very difficult to calculate the form factors by means of Feynman rules, given some particular QFT. Reassuringly, this is not the case, as we can see in the following example. First, we decompose the full 3-point function $\langle\!\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}\mathcal{O}\rangle\!\rangle$ into simple tensors using the choice of momenta (3.7) and denote

$$\langle\!\langle T^{\alpha_1\beta_1}T^{\alpha_2\beta_2}\mathcal{O}\rangle\!\rangle = \tilde{A}_1 p_2^{\alpha_1}p_2^{\beta_1}p_3^{\alpha_2}p_3^{\beta_2} + \tilde{A}_2 p_2^{\alpha_1}p_3^{\alpha_2}\delta^{\beta_1\beta_2} + \tilde{A}_3\delta^{\alpha_1\alpha_2}\delta^{\beta_1\beta_2} + \dots, \tag{3.25}$$

where the omitted terms do not contain the tensors we have listed explicitly. Next, we apply the projectors (3.2). Obverse, for example, that the projector $\Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1)$ is constructed from the metric and the momentum $\boldsymbol{p}_1$ only, and therefore when applied to the 3-point function it cannot change the coefficient of any tensor containing $p_2^{\alpha_1}p_2^{\beta_1}$, *i.e.*,

$$\Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1)p_2^{\alpha_1}p_2^{\beta_1} = p_2^{\mu_1}p_2^{\nu_1} + \dots, \tag{3.26}$$

where the omitted terms do not contain $p_2^{\mu_1}p_2^{\nu_1}$. Using the same argument for $\Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)$, we see that the coefficients of $p_2^{\alpha_1}p_2^{\beta_1}p_3^{\alpha_2}p_3^{\beta_2}$ in (3.25) and $p_2^{\mu_1}p_2^{\nu_1}p_3^{\mu_2}p_3^{\nu_2}$ in $\langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1)t^{\mu_2\nu_2}(\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle\!\rangle$ in (3.23) are equal, *i.e.*, $A_1 = \tilde{A}_1$. Similarly, we find that

$$\begin{aligned}&\Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1)\Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)\langle\!\langle T^{\alpha_1\beta_1}(\boldsymbol{p}_1)T^{\alpha_2\beta_2}(\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle\!\rangle = \\ &\quad = \tilde{A}_1 p_2^{\mu_1}p_2^{\nu_1}p_3^{\mu_2}p_3^{\nu_2} + \frac{1}{4}\tilde{A}_2 p_2^{\mu_1}p_3^{\mu_2}\delta^{\nu_1\nu_2} + \frac{1}{2}\tilde{A}_3\delta^{\mu_1\mu_2}\delta^{\nu_1\nu_2} + \dots,\end{aligned} \tag{3.27}$$

where the omitted terms do not contain the tensors we have listed explicitly. We therefore have

$$A_1 = \text{coefficient of } p_2^{\mu_1}p_2^{\nu_1}p_3^{\mu_2}p_3^{\nu_2} \text{ in } \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle\!\rangle, \tag{3.28}$$

$$A_2 = 4\cdot\text{coefficient of } p_2^{\mu_1}p_3^{\mu_2}\delta^{\nu_1\nu_2} \text{ in } \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle\!\rangle, \tag{3.29}$$

$$A_3 = 2\cdot\text{coefficient of } \delta^{\mu_1\mu_2}\delta^{\nu_1\nu_2} \text{ in } \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle\!\rangle. \tag{3.30}$$

We list the analogous formulae for all other 3-point functions in the second part of the paper.

3.4 Example

Let us consider a conformally coupled free scalar free massless field ϕ in d Euclidean dimensions. In the presence of a non-trivial source $g^{\mu\nu}$ for the stress-energy tensor, the action reads

$$S = \int \mathrm{d}^d x\,\sqrt{g}\left[\frac{1}{2}g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi + \frac{d-2}{8(d-1)}R\phi^2\right], \tag{3.31}$$

where R is the Ricci scalar of $g_{\mu\nu}$. The stress-energy tensor in the presence of the sources is then

$$T_{\mu\nu} = \frac{2}{\sqrt{g}}\frac{\delta S}{\delta g^{\mu\nu}} = \partial_\mu\phi\partial_\nu\phi - \frac{1}{2}g_{\mu\nu}\partial_\alpha\phi\partial^\alpha\phi + \frac{d-2}{4(d-1)}\left(g_{\mu\nu}\nabla^2 - \nabla_\mu\nabla_\nu + R_{\mu\nu}\right)\phi^2. \tag{3.32}$$

In this CFT, $\mathcal{O}(x) = \phi^2(x)$ is a conformal primary operator of dimension $\Delta_3 = d-2$.

For later use we quote the result for the form factors of $\langle\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}\mathcal{O}\rangle\rangle$ in this theory. Writing down the expression for $\langle\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}\mathcal{O}\rangle\rangle$ using the regular Feynman rules, from (3.28, 3.29, 3.30) we may then read off expressions for the form factors. Explicitly evaluating these integrals for the case $d = 3$, we find

$$A_1 = \frac{3(p_1^2+p_2^2)+p_3^2+12p_1p_2+4p_3(p_1+p_2)}{48\,p_3(p_1+p_2+p_3)^4}, \tag{3.33}$$

$$A_2 = \frac{2(p_1^2+p_2^2)+p_3^2+6p_1p_2+3p_3(p_1+p_2)}{12\,(p_1+p_2+p_3)^3}, \tag{3.34}$$

$$A_3 = -\frac{2(p_1^2p_2+p_1p_2^2+p_1p_3^2+p_3p_1^2+p_2p_3^2+p_3p_2^2)+2p_1p_2p_3+p_1^3+p_2^3+p_3^3}{24\,(p_1+p_2+p_3)^2}, \tag{3.35}$$

in agreeement with the direct evaluation of this correlator given in [17].

4 Conformal Ward identities in momentum space

In section 2.2 we wrote down the Ward identities associated with dilatations and special conformal transformation for the case of correlators involving three scalars. In this section, we discuss the corresponding Ward identities for 3-point correlators involving insertions of the stress-energy tensor and conserved currents. First, in section 4.1, we obtain the dilatation and special conformal Ward identities in momentum space by Fourier transforming the well-known position space expressions; in sections 4.2 and 4.3 we then reduce these identities to a set of simple scalar equations using the tensor decomposition introduced in section 3.1. Finally, in sections 4.4 and 4.5, we write down the transverse and trace Ward identities.

4.1 From position space to momentum space

Let $\mathcal{T}_1, \mathcal{T}_2, \ldots, \mathcal{T}_n$ represent quantum operators of dimensions $\Delta_1, \Delta_2, \ldots, \Delta_n$ and of arbitrary Lorentz structure in some CFT. The dilatation Ward identity in position space is especially simple and reads [36]

$$0 = \left[\sum_{j=1}^{n}\Delta_j + \sum_{j=1}^{n} x_j^\alpha \frac{\partial}{\partial x_j^\alpha}\right]\langle \mathcal{T}_1(\boldsymbol{x}_1)\ldots\mathcal{T}_n(\boldsymbol{x}_n)\rangle. \tag{4.1}$$

The Ward identity associated with special conformal transformations for the n-point function of scalar operators $\mathcal{O}_1, \mathcal{O}_2, \ldots, \mathcal{O}_n$ is

$$0 = \left[\sum_{j=1}^{n}\left(2\Delta_j x_j^\kappa + 2x_j^\kappa x_j^\alpha\frac{\partial}{\partial x_j^\alpha} - x_j^2\frac{\partial}{\partial x_{j\kappa}}\right)\right]\langle \mathcal{O}_1(\boldsymbol{x}_1)\ldots\mathcal{O}_n(\boldsymbol{x}_n)\rangle, \tag{4.2}$$

where κ is a free Lorentz index. For general tensors $\mathcal{T}_j$ one needs to add an additional term to the equation. This term depends on the Lorentz structure, and to write it down, we assume that the tensor $\mathcal{T}_j$ has r_j Lorentz indices, *i.e.*, $\mathcal{T}_j = \mathcal{T}_j^{\mu_{j1}\ldots\mu_{jr_j}}$, for $j = 1, 2, \ldots, n$. In this case, the contribution

$$2\sum_{j=1}^{n}\sum_{k=1}^{r_j}\left[(x_j)_{\alpha_{jk}}\delta^{\kappa\mu_{jk}} - x_j^{\mu_{jk}}\delta^\kappa_{\alpha_{jk}}\right]\langle \mathcal{T}_1^{\mu_{11}\ldots\mu_{1n_1}}(\boldsymbol{x}_1)\ldots\mathcal{T}_j^{\mu_{j1}\ldots\alpha_{jk}\ldots\mu_{jr_j}}(\boldsymbol{x}_j)\ldots\mathcal{T}_n^{\mu_{n1}\ldots\mu_{nr_n}}(\boldsymbol{x}_n)\rangle \tag{4.3}$$

must then be added to the right-hand side of (4.2).

Expressions (4.1, 4.2, 4.3) may be Fourier transformed in a similar manner to that discussed in [45]. Due to the translation invariance the position space correlators depend only on the differences $\boldsymbol{x}_j - \boldsymbol{x}_n$. Therefore, we can set $\boldsymbol{x}_n = 0$ and take

$$\boldsymbol{p}_n = -\sum_{j=1}^{n-1} \boldsymbol{p}_j. \tag{4.4}$$

The Ward identities (4.1) and (4.2) then transform to

$$0 = \left[\sum_{j=1}^{n} \Delta_j - (n-1)d - \sum_{j=1}^{n-1} p_j^\alpha \frac{\partial}{\partial p_j^\alpha}\right] \langle\!\langle \mathcal{T}_1(\boldsymbol{p}_1) \dots \mathcal{T}_n(\boldsymbol{p}_n) \rangle\!\rangle, \tag{4.5}$$

$$0 = \left[\sum_{j=1}^{n-1} \left(2(\Delta_j - d)\frac{\partial}{\partial p_j^\kappa} - 2p_j^\alpha \frac{\partial}{\partial p_j^\alpha}\frac{\partial}{\partial p_j^\kappa} + (p_j)_\kappa \frac{\partial}{\partial p_j^\alpha}\frac{\partial}{\partial p_{j\alpha}}\right)\right] \langle\!\langle \mathcal{O}_1(\boldsymbol{p}_1) \dots \mathcal{O}_n(\boldsymbol{p}_n) \rangle\!\rangle, \tag{4.6}$$

while the additional contribution (4.3) transforms to

$$2\sum_{j=1}^{n}\sum_{k=1}^{n_j} \left(\delta^{\mu_{jk}\kappa}\frac{\partial}{\partial p_j^{\alpha_{jk}}} - \delta^\kappa_{\alpha_{jk}}\frac{\partial}{\partial p_{j\mu_{jk}}}\right) \langle\!\langle \mathcal{T}_1^{\mu_{11}\dots\mu_{1r_1}}(\boldsymbol{p}_1) \dots \mathcal{T}_j^{\mu_{j1}\dots\alpha_{jk}\dots\mu_{jr_j}}(\boldsymbol{p}_j) \dots \mathcal{T}_n^{\mu_{n1}\dots\mu_{nr_n}}(\boldsymbol{p}_n), \rangle\!\rangle \tag{4.7}$$

and once again must be added to the right-hand side of (4.6). It will be useful to denote the differential operator obtained by summing the right-hand side of (4.6) and (4.7) as $\mathcal{K}^\kappa$, so that the CWIs may be compactly expressed as

$$\mathcal{K}^\kappa \langle\!\langle \mathcal{T}_1(\boldsymbol{p}_1) \dots \mathcal{T}_n(\boldsymbol{p}_n) \rangle\!\rangle = 0. \tag{4.8}$$

In view of (4.7), note that $\mathcal{K}^\kappa$ acts non-trivially on Lorentz indices and so in fact is really of the form

$$\mathcal{K}^\kappa = \mathcal{K}^{\mu_{11}\dots\mu_{nr_n},\kappa}_{\alpha_{11}\dots\alpha_{nr_n}}, \tag{4.9}$$

however for simplicity we will omit the tensor indices on $\mathcal{K}^\kappa$.

In the following analysis we will focus specifically on 3-point functions. The idea will be to take the tensor decomposition the 3-point function described in section 3.1, then apply the operators (4.6) and (4.7) yielding differential equations for the form factors. Since by Lorentz invariance the form factors are purely functions of the momentum magnitudes, the action of momentum derivatives on form factors may be obtained using the chain rule,

$$\begin{aligned}\frac{\partial}{\partial p_{1\mu}} &= \frac{\partial p_1}{\partial p_{1\mu}}\frac{\partial}{\partial p_1} + \frac{\partial p_2}{\partial p_{1\mu}}\frac{\partial}{\partial p_2} + \frac{\partial p_3}{\partial p_{1\mu}}\frac{\partial}{\partial p_3} \\ &= \frac{p_1^\mu}{p_1}\frac{\partial}{\partial p_1} + \frac{p_1^\mu + p_2^\mu}{p_3}\frac{\partial}{\partial p_3},\end{aligned} \tag{4.10}$$

noting that $\boldsymbol{p}_3$ is fixed via (4.4). We may express derivatives with respect to $\boldsymbol{p}_2$ similarly, and the final results may then be re-expressed purely in terms of the momentum magnitudes.

4.2 Dilatation Ward identity

Using (4.10), it is simple to rewrite the dilatation Ward identity (4.5) for any 3-point function $\langle\!\langle \mathcal{T}_1\mathcal{T}_2\mathcal{T}_3 \rangle\!\rangle$ in terms of its form factors as

$$0 = \left[2d + N_n + \sum_{j=1}^{3}\left(p_j\frac{\partial}{\partial p_j} - \Delta_j\right)\right] A_n(p_1, p_2, p_3), \tag{4.11}$$

where N_n is the tensorial dimension of A_n, *i.e.*, the number of momenta in the tensor multiplying the form factor A_n and the transverse-traceless projectors. As previously, Δ_j, $j = 1, 2, 3$ denote the conformal dimensions of the operators $\mathcal{T}_j$ in the 3-point function: for a conserved current we thus have $\Delta = d - 1$ while for a stress-energy tensor $\Delta = d$.

The dilatation Ward identity determines the total degree of the 3-point function and hence of its form factors. In general, if a function A satisfies

$$0 = \left[-D + \sum_{j=1}^{3} p_j \frac{\partial}{\partial p_j} \right] A(p_1, p_2, p_3) \tag{4.12}$$

for some constant D then we will refer to D as the *degree* of A, denoted $\deg(A) = D$. (A homogeneous polynomial in momenta of degree D has dilatation degree D.) Therefore (4.11) implies that the form factor A_n has degree

$$\deg(A_n) = \Delta_t - 2d - N_n, \tag{4.13}$$

where $\Delta_t = \Delta_1 + \Delta_2 + \Delta_3$.

4.3 Special conformal Ward identities

In this section, we now extract scalar equations for the form factors by inserting our tensor decomposition for the transverse-traceless part of the 3-point functions into the special conformal Ward identities. While the details are somewhat involved, the procedure is nonetheless conceptually straightforward. We will outline the method using as an example the 3-point function $\langle\!\langle T^{\mu_1\nu_1} T^{\mu_2\nu_2} \mathcal{O} \rangle\!\rangle$, which captures all the important features.

Consider then the action of the CWI operator $\mathcal{K}^\kappa$ defined in (4.8) on the decomposition (3.24),

$$\begin{aligned} 0 &= \mathcal{K}^\kappa \langle\!\langle T^{\mu_1\nu_1} T^{\mu_2\nu_2} \mathcal{O} \rangle\!\rangle \\ &= \mathcal{K}^\kappa \langle\!\langle t^{\mu_1\nu_1} t^{\mu_2\nu_2} \mathcal{O} \rangle\!\rangle + \mathcal{K}^\kappa \langle\!\langle t^{\mu_1\nu_1} t^{\mu_2\nu_2}_{\text{loc}} \mathcal{O} \rangle\!\rangle + \mathcal{K}^\kappa \langle\!\langle t^{\mu_1\nu_1}_{\text{loc}} t^{\mu_2\nu_2} \mathcal{O} \rangle\!\rangle + \mathcal{K}^\kappa \langle\!\langle t^{\mu_1\nu_1}_{\text{loc}} t^{\mu_2\nu_2}_{\text{loc}} \mathcal{O} \rangle\!\rangle, \end{aligned} \tag{4.14}$$

recalling that our notation for $\mathcal{K}^\kappa$ suppresses Lorentz indices so that in reality, *e.g.*,

$$\mathcal{K}^\kappa \langle\!\langle t^{\mu_1\nu_1} t^{\mu_2\nu_2} \mathcal{O} \rangle\!\rangle = \mathcal{K}^{\mu_1\nu_1\mu_2\nu_2,\kappa}_{\alpha_1\beta_1\alpha_2\beta_2} \langle\!\langle t^{\alpha_1\beta_1} t^{\alpha_2\beta_2} \mathcal{O} \rangle\!\rangle. \tag{4.15}$$

Through a direct but lengthy calculation we find that the first term on the right-hand side of (4.14), $\mathcal{K}^\kappa \langle\!\langle t^{\mu_1\nu_1} t^{\mu_2\nu_2} \mathcal{O} \rangle\!\rangle$, is transverse-traceless in μ_1, ν_1 and μ_2, ν_2 with respect to the corresponding momenta,

$$\delta_{\mu_1\nu_1} \left[\mathcal{K}^\kappa \langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1) t^{\mu_2\nu_2}(\boldsymbol{p}_2) \mathcal{O}(\boldsymbol{p}_3) \rangle\!\rangle \right] = \delta_{\mu_2\nu_2} \left[\mathcal{K}^\kappa \langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1) t^{\mu_2\nu_2}(\boldsymbol{p}_2) \mathcal{O}(\boldsymbol{p}_3) \rangle\!\rangle \right] = 0, \tag{4.16}$$

$$p_{1\mu_1} \left[\mathcal{K}^\kappa \langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1) t^{\mu_2\nu_2}(\boldsymbol{p}_2) \mathcal{O}(\boldsymbol{p}_3) \rangle\!\rangle \right] = p_{2\mu_2} \left[\mathcal{K}^\kappa \langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1) t^{\mu_2\nu_2}(\boldsymbol{p}_2) \mathcal{O}(\boldsymbol{p}_3) \rangle\!\rangle \right] = 0, \tag{4.17}$$

where we used the definitions (4.6) and (4.7) for $\mathcal{K}^\kappa$ and the identities given in appendix A.9. For correlators involving conserved currents, we find that the analogue of (4.17) similarly applies.

We are now free to apply transverse-traceless projectors (3.1) and (3.2) to (4.14), in order to isolate equations for the form factors appearing in the decomposition of $\langle\!\langle t^{\mu_1\nu_1} t^{\mu_2\nu_2} \mathcal{O} \rangle\!\rangle$. Evaluating the action of $\mathcal{K}^\kappa$ on the semi-local terms in (4.14) via the formulae in appendix A.9, we find

$$\pi^\mu_\alpha \mathcal{K}^\kappa j^\alpha_{\text{loc}} = \frac{2(d-2)}{p^2} \pi^{\kappa\mu} r, \tag{4.18}$$

$$\Pi^{\mu\nu}_{\alpha\beta} \mathcal{K}^\kappa t^{\alpha\beta}_{\text{loc}} = \frac{4d}{p^2} \Pi^{\mu\nu\kappa}{}_\alpha R^\alpha, \tag{4.19}$$

$$\pi^\mu_\alpha j^\alpha_{\text{loc}} = \Pi^{\mu\nu}_{\alpha\beta} t^{\alpha\beta}_{\text{loc}} = 0. \tag{4.20}$$

The last equation implies that any correlation function with more than one insertion of $t^{\mu\nu}_{\rm loc}$ or $j^{\mu}_{\rm loc}$ vanishes when the CWI operator $\mathcal{K}^\kappa$ and the projectors (3.1) and (3.2) are applied. This is because the CWI operator $\mathcal{K}^\kappa$ can be written as a sum of two terms

$$\mathcal{K}^\kappa = \mathcal{K}^\kappa_1\left(\frac{\partial}{\partial p_1^\mu}\right) + \mathcal{K}^\kappa_2\left(\frac{\partial}{\partial p_2^\mu}\right), \tag{4.21}$$

each depending only on derivatives with respect to the appropriate momenta, hence

$$\Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1)\Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)\mathcal{K}^\kappa\langle\!\langle t^{\alpha_1\beta_1}_{\rm loc}(\boldsymbol{p}_1)t^{\alpha_2\beta_2}_{\rm loc}(\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle\!\rangle = 0. \tag{4.22}$$

Substituting all results into (4.14), we find

$$\begin{aligned}0 = {}& \Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1)\Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)\mathcal{K}^\kappa\langle\!\langle t^{\alpha_1\beta_1}(\boldsymbol{p}_1)t^{\alpha_2\beta_2}(\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle\!\rangle \\ &+ \frac{4d}{p_1^2}\Pi^{\mu_1\nu_1\kappa}{}_{\alpha_1}(\boldsymbol{p}_1)\left[p_{1\beta_1}\langle\!\langle T^{\alpha_1\beta_1}(\boldsymbol{p}_1)t^{\mu_2\nu_2}(\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle\!\rangle\right] \\ &+ \frac{4d}{p_2^2}\Pi^{\mu_2\nu_2\kappa}{}_{\alpha_2}(\boldsymbol{p}_2)\left[p_{2\beta_2}\langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1)T^{\alpha_2\beta_2}(\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle\!\rangle\right].\end{aligned} \tag{4.23}$$

Two last terms are semi-local and may be re-expressed in terms of 2-point functions via the transverse Ward identities. The remaining task is then to rewrite the first term of (4.23) in terms of form factors and extract the CWIs. Via the method of section 3.1, we can write the most general form of the result as

$$\begin{aligned}&\Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1)\Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)\mathcal{K}^\kappa\langle\!\langle t^{\alpha_1\beta_1}(\boldsymbol{p}_1)t^{\alpha_2\beta_2}(\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle\!\rangle = \Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1)\Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)\times \\ &\quad\times\Big[p_1^\kappa\left(C_{11}p_2^{\alpha_1}p_2^{\beta_1}p_3^{\alpha_2}p_3^{\beta_2} + C_{12}p_2^{\alpha_1}p_3^{\alpha_2}\delta^{\beta_1\beta_2} + C_{13}\delta^{\alpha_1\alpha_2}\delta^{\beta_1\beta_2}\right) \\ &\qquad + p_2^\kappa\left(C_{21}p_2^{\alpha_1}p_2^{\beta_1}p_3^{\alpha_2}p_3^{\beta_2} + C_{22}p_2^{\alpha_1}p_3^{\alpha_2}\delta^{\beta_1\beta_2} + C_{23}\delta^{\alpha_1\alpha_2}\delta^{\beta_1\beta_2}\right) \\ &\qquad + \Big(C_{31}\delta^{\kappa\alpha_1}p_2^{\beta_1}p_3^{\alpha_2}p_3^{\beta_2} + C_{32}\delta^{\kappa\alpha_1}\delta^{\alpha_2\beta_1}p_3^{\beta_2} \\ &\qquad\quad + C_{41}\delta^{\kappa\alpha_2}p_2^{\alpha_1}p_2^{\beta_1}p_3^{\beta_2} + C_{42}\delta^{\kappa\alpha_2}\delta^{\alpha_1\beta_2}p_2^{\beta_1}\Big)\Big].\end{aligned} \tag{4.24}$$

In this expression, the coefficients C_{jk} are differential equations involving the form factors A_1, A_2 and A_3 of (3.10). Each CWI can then be presented in terms of the momentum magnitudes $p_j = |\boldsymbol{p}_j|$.

As we can see, there are ten coefficients C_{jk} in (4.24), so there are at most ten equations to consider. Usually not all of the CWIs, however, are independent. In this example, the $\boldsymbol{p}_1 \leftrightarrow \boldsymbol{p}_2$ symmetry implies that the equations following from C_{1j} and C_{2j}, as well as C_{3j} and C_{4j}, are pairwise equivalent.

For any 3-point function, the resulting equations can be divided into two groups: the *primary* and the *secondary* CWIs. The primary CWIs are second-order differential equations and appear as the coefficients of transverse or transverse-traceless tensors containing p_1^κ or p_2^κ, where κ is the 'special' index in the CWI operator $\mathcal{K}^\kappa$. In the expression (4.24) above, the primary CWIs are equivalent to the vanishing of the coefficients C_{1j} and C_{2j}. The remaining equations, following from all other transverse or transverse-traceless terms, are then the secondary CWIs and are first-order differential equations. In the expression (4.24), the secondary CWIs are equivalent to the vanishing of the coefficients C_{3j} and C_{4j}.

In the next two subsections we will examine the general form of the primary and secondary CWIs and discuss some of their properties. In section 5, we will return to analyse their solution for the form factors. In outline our strategy will be, first, to solve each of the primary CWIs up to an overall multiplicative constant, then second, to insert these solutions into the secondary CWIs typically allowing the number of undetermined constants to be further reduced. In the case of the correlator $\langle\!\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}\mathcal{O}\rangle\!\rangle$, for example, we will find that the final result is then uniquely determined up to one numerical constant, in agreement with the position space analysis of [3].

4.3.1 Primary conformal Ward identities

It turns out that in all cases the primary CWIs look very similar to the CWIs (2.11) for scalar operators. In order to write the primary CWIs in a simple way, we define the following fundamental differential operators

$$\mathrm{K}_j = \frac{\partial^2}{\partial p_j^2} + \frac{d+1-2\Delta_j}{p_j}\frac{\partial}{\partial p_j}, \quad j = 1,2,3, \tag{4.25}$$

$$\mathrm{K}_{ij} = \mathrm{K}_i - \mathrm{K}_j, \quad j = 1,2,3, \tag{4.26}$$

where Δ_j is the conformal dimension of the j-th operator in the 3-point function under consideration. (Observe that this same operator appeared earlier in (2.11, 2.12).)

In the case of our example $\langle\langle T^{\mu_1\nu_1} T^{\mu_2\nu_2} \mathcal{O} \rangle\rangle$, the primary CWIs for the form factors defined in (3.10) read

$$\begin{array}{lll} \mathrm{K}_{13}\, A_1 = 0, & \mathrm{K}_{13}\, A_2 = 8A_1, & \mathrm{K}_{13}\, A_3 = 2A_2, \\ \mathrm{K}_{23}\, A_1 = 0, & \mathrm{K}_{23}\, A_2 = 8A_1, & \mathrm{K}_{23}\, A_3 = 2A_2, \end{array} \tag{4.27}$$

Note that, from the definition (4.26), we have

$$\mathrm{K}_{ii} = 0, \qquad \mathrm{K}_{ji} = -\mathrm{K}_{ij}, \qquad \mathrm{K}_{ij} + \mathrm{K}_{jk} = \mathrm{K}_{ik}, \tag{4.28}$$

for any $i, j, k \in \{1,2,3\}$. One can therefore subtract corresponding pairs of equations and obtain the following system of independent partial differential equations

$$\begin{array}{lll} \mathrm{K}_{12}\, A_1 = 0, & \mathrm{K}_{12}\, A_2 = 0, & \mathrm{K}_{12}\, A_3 = 0, \\ \mathrm{K}_{13}\, A_1 = 0, & \mathrm{K}_{13}\, A_2 = 8A_1, & \mathrm{K}_{13}\, A_3 = 2A_2. \end{array} \tag{4.29}$$

As we will prove, each equation has a unique solution up to one numerical constant. This means that at this point the 3-point function $\langle\langle T^{\mu_1\nu_1} T^{\mu_2\nu_2} \mathcal{O} \rangle\rangle$ is determined by three numerical constants. After the application of the secondary CWIs this number will decrease further.

The primary CWIs for all 3-point functions are listed explicitly in the second part of the paper. They share the following properties:

1. All primary CWIs are second-order linear differential equations.

2. The equations for the coefficient A_1 are always homogeneous and given by (2.11, 2.12) exactly, *i.e.*,
$$\mathrm{K}_{12}\, A_1 = 0, \qquad \mathrm{K}_{13}\, A_1 = 0. \tag{4.30}$$

3. The equations for the remaining form factors are similar to (2.11, 2.12), but they may contain a linear inhomogeneous part. For a form factor A_n multiplying a tensor of tensorial dimension N_n, the only form factors A_j which can appear in the inhomogeneous part are those with $N_j = N_n + 2$. It is therefore always possible to solve the primary CWIs recursively, starting with A_1.

 In the case of our example, the recursive structure of the equations (4.29) is clearly visible.

4. There is no semi-local contribution to any primary CWI. In our example, last two terms in (4.23) do not contribute to the primary CWIs. This conclusion is valid in general and can be checked explicitly in all cases.

5. The solution to each pair of primary CWIs is unique up to one numerical constant, as we will prove in section 5.

4.3.2 Secondary conformal Ward identities

The secondary CWIs are first-order partial differential equations and in principle involve the semi-local information contained in j^μ_{loc} and $t^{\mu\nu}_{\text{loc}}$. In order to write them compactly, we define the two differential operators

$$\begin{aligned}\mathrm{L}_{s,N} &= p_1(p_1^2+p_2^2-p_3^2)\frac{\partial}{\partial p_1} + 2p_1^2 p_2 \frac{\partial}{\partial p_2} \\ &\quad + \left[(2d-\Delta_1-2\Delta_2+N+s)p_1^2 + (\Delta_1-2+s)(p_3^2-p_2^2)\right], \qquad (4.31)\\ \mathrm{R}_s &= p_1\frac{\partial}{\partial p_1} - (\Delta_1-2+s), \qquad (4.32)\end{aligned}$$

as well as their symmetric versions

$$\begin{aligned}\mathrm{L}'_{s,N} &= \mathrm{L}_{s,N} \text{ with } p_1 \leftrightarrow p_2 \text{ and } \Delta_1 \leftrightarrow \Delta_2, \qquad (4.33)\\ \mathrm{R}'_s &= \mathrm{R}_s \text{ with } p_1 \mapsto p_2 \text{ and } \Delta_1 \mapsto \Delta_2. \qquad (4.34)\end{aligned}$$

These operators depend on two parameters N and s determined by the Ward identity in question. Physically, while N has no clear interpretation, the parameter s denotes the spin of the first operator insertion in the 3-point function (or the second in the case of $\mathrm{L}'_{s,N}$ and R'_s). For conserved currents, we therefore have $s=1$ while for the stress-energy tensor $s=2$.

In our example (3.10) one finds two independent secondary CWIs following from the coefficients C_{31} and C_{32} in (4.24), namely

$$\begin{aligned}&\mathrm{L}_{2,0}\,A_1 + \mathrm{R}_2\,A_2 = 2d\cdot\text{coeff. of } p_2^{\mu_1}p_3^{\mu_2}p_3^{\nu_2} \text{ in } p_{1\nu_1}\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle\!\rangle, \qquad (4.35)\\ &\mathrm{L}_{2,0}\,A_2 + 4\,\mathrm{R}_2\,A_3 = 8d\cdot\text{coeff. of } \delta^{\mu_1\mu_2}p_3^{\mu_2} \text{ in } p_{1\nu_1}\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle\!\rangle, \qquad (4.36)\end{aligned}$$

with $\Delta_1=\Delta_2=d$. Note that in order to correctly extract the coefficient of a tensor, the rule (3.7) regarding the momenta associated with a given Lorentz index must be observed. The semi-local terms on the right-hand sides may be computed by means of transverse Ward identities, to which we now turn our attention.

4.4 Transverse Ward identities

In this section we review briefly the transverse (or diffeomorphism) Ward identities in momentum space. These Ward identities arise from the conservation law for currents. In particular we will need the precise form of all semi-local terms that appear in these Ward identities since these terms are required for the explicit evaluation of the right-hand sides of the secondary CWIs such as (4.35, 4.36).

We assume the CFT contains the following data:

- A symmetry group G. The conserved current $J^{\mu a}$, $a=1,\ldots,\dim G$, is then the Noether current associated with the symmetry and is sourced by a potential A^a_μ.
- Scalar primary operators $\mathcal{O}^I$ all of the same dimension Δ, sourced by ϕ^I_0.
- A stress-energy tensor $T_{\mu\nu}$ sourced by the metric $g^{\mu\nu}$.

Under a symmetry transformation with parameter α^a the sources transform as

$$\begin{aligned}\delta g^{\mu\nu} &= 0, \qquad (4.37)\\ \delta A^a_\mu &= -D^{ac}_\mu \alpha^c = -\partial_\mu\alpha^a - f^{abc}A^b_\mu\alpha^c, \qquad (4.38)\\ \delta\phi^I_0 &= -\mathrm{i}\alpha^a (T^a_R)^{IJ}\phi^J_0, \qquad (4.39)\end{aligned}$$

where T^a_R are matrices of a representation R and f^{abc} are structure constants of the group G. The gauge field transforms in the adjoint representation while the ϕ^I may transform in any representation R. The convariant derivative is $D^{IJ}_\mu = \delta^{IJ}\partial_\mu - \mathrm{i}A^a_\mu (T^a_R)^{IJ}$.

Similarly, under a diffeomorphism ξ^μ the sources transform as

$$\delta g^{\mu\nu} = -(\nabla^\mu \xi^\nu + \nabla^\nu \xi^\mu), \tag{4.40}$$

$$\delta A^a_\mu = \xi^\nu \nabla_\nu A^a_\mu + \nabla_\mu \xi^\nu \cdot A^a_\nu, \tag{4.41}$$

$$\delta \phi^I_0 = \xi^\nu \partial_\nu \phi^I_0, \tag{4.42}$$

where ∇ is a Levi-Civita connection.

From the generating functional

$$Z[\phi^I_0, A^a_\mu, g^{\mu\nu}] = \int \mathcal{D}\Phi \, \exp\left(-S[A^a_\mu, g^{\mu\nu}] - \int \mathrm{d}^d \boldsymbol{x} \, \sqrt{g} \phi^I_0 \mathcal{O}^I \right), \tag{4.43}$$

we have the one-point functions in the presence of sources

$$\langle T_{\mu\nu}(\boldsymbol{x})\rangle = -\frac{2}{\sqrt{g(\boldsymbol{x})}} \frac{\delta}{\delta g^{\mu\nu}(\boldsymbol{x})} Z, \tag{4.44}$$

$$\langle J^{\mu a}(\boldsymbol{x})\rangle = -\frac{1}{\sqrt{g(\boldsymbol{x})}} \frac{\delta}{\delta A^a_\mu(\boldsymbol{x})} Z, \tag{4.45}$$

$$\langle \mathcal{O}^I(\boldsymbol{x})\rangle = -\frac{1}{\sqrt{g(\boldsymbol{x})}} \frac{\delta}{\delta \phi^I_0(\boldsymbol{x})} Z. \tag{4.46}$$

By taking more functional derivatives we can obtain higher-point correlation functions, *e.g.*,

$$\begin{aligned}\langle T_{\mu_1\nu_1}(\boldsymbol{x}) T_{\mu_2\nu_2}(\boldsymbol{y}) T_{\mu_3\nu_3}(\boldsymbol{z})\rangle = {} & \frac{-2}{\sqrt{g(\boldsymbol{z})}} \frac{\delta}{\delta g^{\mu_3\nu_3}(\boldsymbol{z})} \frac{-2}{\sqrt{g(\boldsymbol{y})}} \frac{\delta}{\delta g^{\mu_2\nu_2}(\boldsymbol{y})} \frac{-2}{\sqrt{g(\boldsymbol{x})}} \frac{\delta}{\delta g^{\mu_1\nu_1}(\boldsymbol{x})} Z[g^{\mu\nu}] \\ & + 2\langle \frac{\delta T_{\mu_1\nu_1}(\boldsymbol{x})}{\delta g^{\mu_2\nu_2}(\boldsymbol{y})} T_{\mu_3\nu_3}(\boldsymbol{z})\rangle + 2\langle \frac{\delta T_{\mu_2\nu_2}(\boldsymbol{y})}{\delta g^{\mu_3\nu_3}(\boldsymbol{z})} T_{\mu_1\nu_1}(\boldsymbol{x})\rangle + 2\langle \frac{\delta T_{\mu_3\nu_3}(\boldsymbol{z})}{\delta g^{\mu_1\nu_1}(\boldsymbol{x})} T_{\mu_2\nu_2}(\boldsymbol{y})\rangle. \end{aligned} \tag{4.47}$$

Note that here we define the 3-point function of the stress-energy tensor to be the correlator of three separate stress-energy tensor insertions (and similarly for other correlators involving conserved currents), rather than the correlator obtained by functionally differentiating the generating functional with respect to the metric three times. While the latter definition is used in [3, 4, 13, 30], our definition here is simpler for direct QFT computations. To convert between the two definitions simply requires the addition or subtraction of the semi-local terms in the formula above.

Requiring the partition function to be invariant under variation of the sources then leads to the transverse Ward identities

$$\begin{aligned} 0 &= D^{ac}_\mu \langle J^{\mu a}\rangle - \mathrm{i}(T^a_R)^{IJ} \phi^J_0 \langle \mathcal{O}^I\rangle \\ &= \nabla_\mu \langle J^{\mu a}\rangle + f^{abc} A^b_\mu \langle J^{\mu c}\rangle - \mathrm{i}(T^a_R)^{IJ} \phi^J_0 \langle \mathcal{O}^I\rangle, \end{aligned} \tag{4.48}$$

$$\begin{aligned} 0 &= \nabla^\mu \langle T_{\mu\nu}\rangle + \nabla_\nu A^a_\mu \cdot \langle J^{\mu a}\rangle - \nabla_\mu A^a_\nu \cdot \langle J^{\mu a}\rangle + \partial_\nu \phi^I_0 \cdot \langle \mathcal{O}^I\rangle - A^a_\nu \nabla_\mu \langle J^{\mu a}\rangle \\ &= \nabla^\mu \langle T_{\mu\nu}\rangle - F^a_{\mu\nu} \langle J^{\mu a}\rangle + D^{IJ}_\nu \phi^J_0 \cdot \langle \mathcal{O}^I\rangle. \end{aligned} \tag{4.49}$$

These equations may then be differentiated with respect to the sources to obtain the corresponding Ward identities for higher point functions.

Explicit expressions for all the higher-point transverse Ward identities we need are listed in the second part of the paper. In obtaining these expressions we have used the assumptions:

1. $\mathcal{O}^I$ is independent of the sources, *i.e.*,

$$\frac{\delta \mathcal{O}^I}{\delta \phi^J_0} = 0, \qquad \frac{\delta \mathcal{O}^I}{\delta A^a_\mu} = 0, \qquad \frac{\delta \mathcal{O}^I}{\delta g^{\mu\nu}} = 0. \tag{4.50}$$

2. The source ϕ_0^I appears only as in (4.43), so that

$$\frac{\delta T_{\mu\nu}(\boldsymbol{x})}{\delta\phi_0^I(\boldsymbol{y})} = -g_{\mu\nu}(\boldsymbol{x})\mathcal{O}(\boldsymbol{x})\delta(\boldsymbol{x}-\boldsymbol{y}), \qquad \frac{\delta J^{\mu a}}{\delta\phi_0^I} = 0. \tag{4.51}$$

3. The gauge field A_μ^a couples either through covariant derivatives or acts as an external source for the current in the form of $A_\mu^a J^{\mu a}$. This means there are no kinetic terms for A_μ^a, *i.e.*, no derivatives acting on A_μ^a in the action, hence

$$\frac{\delta T_{\mu\nu}(\boldsymbol{x})}{\delta A_\rho^a(\boldsymbol{y})} = F_{\mu\nu}^{\rho a}(\boldsymbol{x})\delta(\boldsymbol{x}-\boldsymbol{y}), \qquad \frac{\delta J^{\mu a}(\boldsymbol{x})}{\delta A_\nu^b(\boldsymbol{y})} = G^{\mu\nu ab}(\boldsymbol{x})\delta(\boldsymbol{x}-\boldsymbol{y}) \tag{4.52}$$

where F and G are functions of the CFT fields.

Of course, it may happen that renormalisation requires us to add counterterms violating one or more of the assumptions above, in which case the relevant Ward identities would need to be modified accordingly.

As a specific illustration of the general discussion above, let us consider the transverse Ward identity for $\langle\!\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}\mathcal{O}\rangle\!\rangle$ for a matter content consisting of conformal scalars, as defined in section 3.4. We will take the operator $\mathcal{O}=\phi^2$. The relevant Ward identity is

$$p_1^{\nu_1}\langle\!\langle T_{\mu_1\nu_1}(\boldsymbol{p}_1)T_{\mu_2\nu_2}(\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle = 2p_1^{\nu_1}\langle\!\langle \frac{\delta T_{\mu_1\nu_1}}{\delta g^{\mu_2\nu_2}}(\boldsymbol{p}_1,\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle, \tag{4.53}$$

where $\delta T_{\mu_1\nu_1}/\delta g^{\mu_2\nu_2}$ denotes taking the functional derivative of the stress-energy tensor with respect to the metric, after which we restore the metric to its background value $g_{\mu\nu}=\delta_{\mu\nu}$. Evaluating this functional derivative explicitly using (3.32), we find [17]

$$\begin{aligned}\frac{\delta T_{\mu\nu}(\boldsymbol{x})}{\delta g^{\rho\sigma}(\boldsymbol{y})} = &-\frac{1}{2}\left[\delta_{\mu\nu}\delta_\rho^\alpha\delta_\sigma^\beta + 2\delta_{\mu(\rho}\delta_{\sigma)\nu} - \delta_{\mu\nu}\delta_{\rho\sigma}\delta^{\alpha\beta}\right]\delta(\boldsymbol{x}-\boldsymbol{y})T_{\alpha\beta}(\boldsymbol{x})\\ &+\frac{1}{16}\Big[C^{(1)\alpha\beta}_{\mu\nu\rho\sigma}\delta(\boldsymbol{x}-\boldsymbol{y})\partial_\alpha\partial_\beta + C^{(2)\alpha\beta}_{\mu\nu\rho\sigma}\partial_\alpha\delta(\boldsymbol{x}-\boldsymbol{y})\partial_\beta\\ &\quad + C^{(3)\alpha\beta}_{\mu\nu\rho\sigma}\partial_\alpha\partial_\beta\delta(\boldsymbol{x}-\boldsymbol{y})\Big]\mathcal{O}(\boldsymbol{x}),\end{aligned} \tag{4.54}$$

where partial derivatives are taken with respect to $\boldsymbol{x}$ and the prefactors are

$$C^{(1)\alpha\beta}_{\mu\nu\rho\sigma} = \delta_{\mu\nu}\delta_\rho^{(\alpha}\delta_\sigma^{\beta)} + 2\delta_{\mu(\rho}\delta_{\sigma)\nu}\delta^{\alpha\beta} - \delta_{\mu\nu}\delta_{\rho\sigma}\delta^{\alpha\beta}, \tag{4.55}$$

$$C^{(2)\alpha\beta}_{\mu\nu\rho\sigma} = 2\delta_{\mu\nu}\delta_\rho^{(\alpha}\delta_\sigma^{\beta)} + \delta_{\mu(\rho}\delta_{\sigma)\nu}\delta^{\alpha\beta} - \delta_{\mu\nu}\delta_{\rho\sigma}\delta^{\alpha\beta} - 2\delta^{\alpha}_{(\mu}\delta_{\nu)(\rho}\delta^{\beta}_{\sigma)}, \tag{4.56}$$

$$C^{(3)\alpha\beta}_{\mu\nu\rho\sigma} = \delta_{\mu\nu}\delta_\rho^{(\alpha}\delta_\sigma^{\beta)} + \delta_{\mu(\rho}\delta_{\sigma)\nu}\delta^{\alpha\beta} - \delta_{\mu\nu}\delta_{\rho\sigma}\delta^{\alpha\beta} - 2\delta^{\alpha}_{(\mu}\delta_{\nu)(\rho}\delta^{\beta}_{\sigma)} + \delta_{\rho\sigma}\delta_\mu^{(\alpha}\delta_\nu^{\beta)}. \tag{4.57}$$

After Fourier transforming and using the result for the 2-point function

$$\langle\!\langle\mathcal{O}(\boldsymbol{p})\mathcal{O}(-\boldsymbol{p})\rangle\!\rangle = \frac{1}{4p} \tag{4.58}$$

we obtain

$$p_{1\nu_1}\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle\!\rangle = -\frac{1}{32\,p_3}p_2^{\mu_1}p_3^{\mu_2}p_3^{\nu_2} + \frac{p_3}{32}\delta^{\mu_1\mu_2}p_3^{\nu_2} + \ldots, \tag{4.59}$$

where we have retained only the terms appearing in the right-hand sides of the secondary CWIs (4.35) and (4.36). The omitted terms do not contain the tensors listed explicitly and will play no further role in our analysis. As usual, we use the convention (3.7) for the Lorentz indices.

4.5 Trace Ward identities

Invariance of the generating functional (4.43) under the Weyl transformations

$$\delta g^{\mu\nu} = 2g^{\mu\nu}\delta\sigma, \qquad \delta\phi_0^I = (d-\Delta)\phi_0^I\delta\sigma, \qquad \delta A_\mu^a = 0. \tag{4.60}$$

leads to the trace (or Weyl) Ward identity in the presence of the sources

$$\langle T(\boldsymbol{x})\rangle = (\Delta - d)\phi_0^I(\boldsymbol{x})\langle\mathcal{O}^I(\boldsymbol{x})\rangle, \tag{4.61}$$

where $T = T_\mu^\mu$. Functionally differentiating with respect to the sources then yields trace Ward identities for 3-point functions, *e.g.*,

$$\langle\!\langle T(\boldsymbol{p}_1)\mathcal{O}^I(\boldsymbol{p}_2)\mathcal{O}^J(\boldsymbol{p}_3)\rangle\!\rangle = -\Delta\left[\langle\!\langle\mathcal{O}^I(\boldsymbol{p}_2)\mathcal{O}^J(-\boldsymbol{p}_2)\rangle\!\rangle + \langle\!\langle\mathcal{O}^I(\boldsymbol{p}_3)\mathcal{O}^J(-\boldsymbol{p}_3)\rangle\!\rangle\right], \tag{4.62}$$

$$\langle\!\langle T(\boldsymbol{p}_1)T_{\mu\nu}(\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle = 2\langle\!\langle\frac{\delta T}{\delta g^{\mu\nu}}(\boldsymbol{p}_1,\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle. \tag{4.63}$$

A complete list of all trace Ward identities is given in the second part of the paper.

As is well known, due to renormalisation the trace Ward identity may acquire an anomalous contribution. The exact contribution depends strongly on the specifics of the theory, but its form is universal. In this paper we assume no anomalies in the transverse Ward identities (4.48) and (4.49) can appear. The anomalous contributions are therefore still transverse.

In section 8 we will consider in detail the divergences and anomalies in the correlation functions $\langle\!\langle T^{\mu_1\nu_1}J^{\mu_2}J^{\mu_3}\rangle\!\rangle$ and $\langle\!\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}T^{\mu_3\nu_3}\rangle\!\rangle$ for the case $d = 4$. Here, as an example, we consider the most general form of the trace anomalies in the correlation functions considered above, which is

$$\langle\!\langle T(\boldsymbol{p}_1)\mathcal{O}^I(\boldsymbol{p}_2)\mathcal{O}^J(\boldsymbol{p}_3)\rangle\!\rangle_{\text{anomaly}} = B_1^{IJ}, \tag{4.64}$$

$$\langle\!\langle T(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle_{\text{anomaly}} = \Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)p_3^{\alpha_2}p_3^{\beta_2}B_1^I + \pi^{\mu_2\nu_2}(\boldsymbol{p}_2)B_2^I, \tag{4.65}$$

where the form factors B_1^{IJ}, B_1^I and B_2^I are functions of the momentum magnitudes specific to the theory in question. For example, the contribution to $\langle\!\langle T^{\mu_1\nu_1}\mathcal{O}^I\mathcal{O}^J\rangle\!\rangle_{\text{anomaly}}$ follows from the counterterm, [3, 32]

$$\frac{1}{2}\int \mathrm{d}^d\boldsymbol{x}\sqrt{g}P^{IJ}\phi_0^I\Box^{\Delta-\frac{d}{2}}\phi_0^J, \tag{4.66}$$

where P^{IJ} are numerical coefficients and we assume that $\Delta - \frac{d}{2}$ is a non-negative integer. In this case we then find

$$B_1^{IJ} = (-p_1^2)^{\Delta-\frac{d}{2}}P^{IJ}. \tag{4.67}$$

In section 8 we provide a short worked example of the renormalisation procedure in the case of the $\langle\!\langle T^{\mu_1\nu_1}J^{\mu_2}J^{\mu_3}\rangle\!\rangle$ and $\langle\!\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}T^{\mu_3\nu_3}\rangle\!\rangle$ correlators and show how the anomalies arise.

5 Solutions to conformal Ward identities

It is a rather pleasant fact that all the primary CWIs can be solved in terms of the triple-K integrals similar to (2.26). We will start by analysing some properties of the triple-K integrals before proceeding to show how this class of integrals solves the primary CWIs. In particular, we will find that the solution to each primary CWI is unique up to one numerical constant. Finally, we will analyse the structure and implications of the secondary CWIs.

5.1 Triple-K integrals

All primary CWIs can be solved in terms of the general triple-K integral

$$I_{\alpha\{\beta_1\beta_2\beta_3\}}(p_1,p_2,p_3)=\int_0^\infty \mathrm{d}x\, x^\alpha \prod_{j=1}^3 p_j^{\beta_j} K_{\beta_j}(p_j x), \tag{5.1}$$

where K_ν is a Bessel K function. This integral depends on four parameters, namely the power α of the integration variable x, and the three Bessel function indices β_j. Its arguments, p_1, p_2, p_3, are magnitudes of momenta $p_j = |\boldsymbol{p}_j|$, $j = 1, 2, 3$. In the following we will generically refer to these as α and β parameters respectively.

It will be useful to define a reduced version $J_{N\{k_1k_2k_3\}}$ of the triple-K integral by substituting

$$\alpha=\frac{d}{2}-1+N, \qquad \beta_j=\Delta_j-\frac{d}{2}+k_j, \quad j=1,2,3. \tag{5.2}$$

Here we assume that we concentrate on some particular 3-point function and the conformal dimensions Δ_j, $j = 1, 2, 3$ are therefore fixed. In other words we define

$$J_{N\{k_j\}}=I_{\frac{d}{2}-1+N\{\Delta_j-\frac{d}{2}+k_j\}}, \tag{5.3}$$

where we use a shortened notation $\{k_j\} = \{k_1k_2k_3\}$, etc. Finally we define

$$\Delta_t=\Delta_1+\Delta_2+\Delta_3, \qquad \beta_t=\beta_1+\beta_2+\beta_3, \qquad k_t=k_1+k_2+k_3. \tag{5.4}$$

The main point of this section is to present relations showing that all primary CWIs for a given 3-point function can be solved by the triple-K integrals (5.1). The representation (5.3) turns out to be extremely useful, as the parameters N and k_j are fixed by the primary CWIs and have no dependence on either Δ_j or d. If desired, these triple-K integrals may also be re-expressed in terms of other familiar integrals such as Feynman or Schwinger parametrised integrals, as discussed in appendix A.3.

5.1.1 Region of validity, regularisation and renormalisation

We assume all parameters and variables in the triple-K integral (5.1) are real. From the asymptotic expansion (A.4.8) the integral converges at large x, however in general there may still be a divergence at $x = 0$. From the series expansion (A.4.1) and the definition (A.4.2), we see the triple-K integral only converges if [40]

$$\alpha>\sum_{j=1}^3|\beta_j|+1, \qquad p_1,p_2,p_3>0. \tag{5.5}$$

If α does not satisfy this inequality, we can regard the triple-K integral (5.1) as a function of α

$$\alpha\mapsto I_{\alpha\{\beta_1\beta_2\beta_3\}}(p_1,p_2,p_3). \tag{5.6}$$

with the other parameters and momenta fixed and use analytic continuation. The scheme (2.5) is very convenient here as it does not change the indices of the Bessel functions. In terms of the parameters α and β_j this corresponds to the substitutions

$$\alpha\mapsto\alpha+\epsilon, \qquad \beta_j \text{ does not change, } j=1,2,3 \tag{5.7}$$

in (5.1). Generically one finds that the limit $\epsilon \to 0$ then exists, except for cases where

$$\alpha+1\pm\beta_1\pm\beta_2\pm\beta_3=-2n, \tag{5.8}$$

for some non-negative n. In these cases we find poles in ϵ. This can be seen as follows. The only source of the singularity is the divergence of the integral at $x = 0$. Therefore we can expand the integrand about $x = 0$ and look for possible divergences.

First, assume that the condition (5.8) is met but all $\beta_j \notin \mathbb{Z}$. We will show that in such a case we should expect single poles in ϵ. Indeed, equations (A.4.1) and (A.4.2) show that the expansion contains power terms x^a for various $a \in \mathbb{R}$ only. Note that the function (5.6) is singular only if there exists a term x^a in the expansion of the integrand with $a = -1$. Indeed, the integral of such a term near $x = 0$ is

$$\int_0 x^a \mathrm{d}x = \frac{\text{const}}{a+1}, \tag{5.9}$$

which has a single pole at $a = -1$ only. Note that while the triple-K integral as it stands will diverge if the expansion contains any x^a terms with $a < -1$, the analytic continuation will exist due to (5.9). Therefore, the function (5.6) is well-defined as long as the expansion of the integrand near $x = 0$ does not contain a $1/x$ term, which is exactly the condition (5.8).

Now observe that if some $\beta_j \in \mathbb{Z}$, then the series expansion of the integrand in the triple-K integral about $x = 0$ contains logarithms due to (A.4.7). Since, for example

$$\int_0 x^a \log x \, \mathrm{d}x = -\frac{\text{const}_1}{(a+1)^2} + \frac{\text{const}_2}{a+1}, \tag{5.10}$$

we may expect a double pole in α in (5.6) when some of the β_j are integer. While the order of the pole may increase, its position remains at $a = -1$. Using the series expansions (A.4.1) and (A.4.7), we can see that the positions of the poles in any triple-K integral are given by (5.8).

5.1.2 Basic properties

Having defined triple-K integrals, we want to show that they solve the primary CWIs. We will therefore now analyse the basic properties of the triple-K integrals. The most obvious of these is the permutation symmetry

$$I_{\alpha\{\beta_{\sigma(1)}\beta_{\sigma(2)}\beta_{\sigma(3)}\}}(p_1,p_2,p_3) = I_{\alpha\{\beta_1\beta_2\beta_3\}}(p_{\sigma^{-1}(1)},p_{\sigma^{-1}(2)},p_{\sigma^{-1}(3)}), \tag{5.11}$$

where σ is any permutation of the set $\{1,2,3\}$. We also have the relations

$$\frac{\partial}{\partial p_n} I_{\alpha\{\beta_j\}} = -p_n I_{\alpha+1\{\beta_j-\delta_{jn}\}}, \tag{5.12}$$

$$I_{\alpha\{\beta_j+\delta_{jn}\}} = p_n^2 I_{\alpha\{\beta_j-\delta_{jn}\}} + 2\beta_n I_{\alpha-1\{\beta_j\}}, \tag{5.13}$$

$$I_{\alpha\{\beta_1\beta_2,-\beta_3\}} = p_3^{-2\beta_3} I_{\alpha\{\beta_1\beta_2\beta_3\}}, \tag{5.14}$$

for any $n = 1,2,3$, as follows from the basic Bessel function relations

$$\frac{\partial}{\partial a}\left[a^\nu K_\nu(ax)\right] = -x a^\nu K_{\nu-1}(ax), \tag{5.15}$$

$$K_{\nu-1}(x) + \frac{2\nu}{x} K_\nu(x) = K_{\nu+1}(x), \tag{5.16}$$

$$K_{-\nu}(x) = K_\nu(x). \tag{5.17}$$

Some additional properties of Bessel functions and triple-K integrals are listed in appendix A.4.

5.1.3 Dilatation degree of the triple-K integral

As the triple-K integral solves CWIs, it should also solve the dilatation Ward identity (4.12). In other words it should have a definite dilatation dimension. Using (5.15) and (5.12) we can write

$$\begin{aligned}\int_0^\infty \mathrm{d}x \frac{\partial}{\partial x}\left(x^{\alpha+1}\prod_{j=1}^3 p_j^{\beta_j} K_{\beta_j}(p_j x)\right) &= \\ &= (\alpha+1-\beta_t) I_{\alpha\{\beta_k\}} - \sum_{j=1}^3 p_j^2 I_{\alpha+1\{\beta_k-\delta_{jk}\}} \\ &= (\alpha+1-\beta_t) I_{\alpha\{\beta_k\}} + \sum_{j=1}^3 p_j \frac{\partial}{\partial p_j} I_{\alpha\{\beta_k\}}. \end{aligned} \tag{5.18}$$

The expression on the left-hand side leads to a boundary term at $x = 0$. In the region of convergence (5.5) all integrals in this expression are well-defined and the boundary term vanishes. Now we can use the analytic continuation (5.7) in order to argue that the analytically continued left-hand side vanishes, except in the case where (5.8) is satisfied. Indeed, if we regard both sides of (5.18) as analytic functions of α with other parameters and momenta fixed, then the validity of (5.18) in the region (5.5) implies its validity in the entire domain of analyticity. Therefore, we have shown that

$$\deg I_{\alpha\{\beta_j\}} = \beta_t - \alpha - 1, \qquad \deg J_{N\{k_j\}} = \Delta_t + k_t - 2d - N \tag{5.19}$$

provided $\alpha + 1 \pm \beta_1 \pm \beta_2 \pm \beta_3 \neq -2n$ for some non-negative n and independent choice of signs.

Finally we can argue that, if (5.8) holds, we should expect scaling anomalies in $I_{\alpha\{\beta_j\}}$. Using the power series expansion (A.4.1) of the Bessel I functions one can see that the series expansion of the boundary term $x^{\alpha+1}\prod_{j=1}^3 p_j^{\beta_j} K_{\beta_j}(p_j x)$ in (5.18) about $x = 0$ contains a constant piece exactly when (5.8) holds. This indicates that the dilatation Ward identity for the $I_{\alpha\{\beta_j\}}$ is not satisfied in such a case. Note that this is not a strict argument since the regulator cannot be removed from the integrals appearing in (5.18). One should instead expand both sides in the regulator ϵ and match terms order by order.

5.2 Solutions to the primary conformal Ward identities

In the previous section we defined the triple-K integral and analysed its basic properties. We now want to use this knowledge in order to write a solution to the CWIs. For this we need the following fundamental identity. For any $m, n = 1, 2, 3$,

$$\mathrm{K}_{mn}\, J_{N\{k_j\}} = -2k_m J_{N+1\{k_j-\delta_{jm}\}} + 2k_n J_{N+1\{k_j-\delta_{jn}\}}, \tag{5.20}$$

for $k_1, k_2, k_3, N \in \mathbb{R}$. The operator K_{mn} is the CWI operator defined in (4.26). This relation is a direct consequence of the identities (5.12) and (5.13).

Let us first consider the pair of primary CWIs for the form factor A_1. As discussed in section 4.3.1, such CWIs are always homogeneous and take the form (4.30). Observe that if we set all $k_j = 0$ in (5.20) then $A_1 = \alpha_1 J_{N\{000\}}$ is a solution for arbitrary $N \in \mathbb{R}$ and an integration constant $\alpha_1 \in \mathbb{R}$. Furthermore, observe that, if we impose only one homogeneous equation, say $\mathrm{K}_{12}\, A = 0$, then the most general solution in terms of the triple-K integrals is $\alpha J_{N\{00k_3\}}$ for any $\alpha, N, k_3 \in \mathbb{R}$. In general the equation (5.20) is sufficient to write down solutions to all primary CWIs.

The remaining piece of information is the value of N. In general, if $A_n = \alpha_n J_{N\{k_j\}}$ is a form factor of tensorial dimension N_n, then (4.13) and (5.19) determine the value of $N = N(A_n)$ to be

$$N(A_n) = N_n + k_t. \tag{5.21}$$

Let us see how the procedure works for our example $\langle\!\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}\mathcal{O}\rangle\!\rangle$. The primary CWIs are given by (4.29) and, in particular,

$$\begin{aligned}
&\mathrm{K}_{12}\,A_1 = 0, \qquad \mathrm{K}_{13}\,A_1 = 0,\\
&\mathrm{K}_{12}\,A_2 = 0, \qquad \mathrm{K}_{12}\,\mathrm{K}_{13}\,A_2 = 0, \qquad \mathrm{K}_{13}^2\,A_2 = 0,\\
&\mathrm{K}_{12}\,A_3 = 0, \qquad \mathrm{K}_{12}\,\mathrm{K}_{13}\,A_3 = 0, \qquad \mathrm{K}_{12}\,\mathrm{K}_{13}^2\,A_3 = 0, \qquad \mathrm{K}_{13}^3\,A_3 = 0,
\end{aligned} \tag{5.22}$$

Therefore, using (5.20) and (5.21), we can write the most general solution given in terms of the triple-K integrals,

$$A_1 = \alpha_1 J_{4\{000\}}, \tag{5.23}$$

$$A_2 = \alpha_{21} J_{3\{001\}} + \alpha_2 J_{2\{000\}}, \tag{5.24}$$

$$A_3 = \alpha_{31} J_{2\{002\}} + \alpha_{32} J_{1\{001\}} + \alpha_3 J_{0\{000\}}, \tag{5.25}$$

where all the α are numerical constants. Finally, the inhomogeneous parts of (4.29) fix some of these constants. When the solution above is substituted into the primary CWIs, (5.20) requires that

$$\alpha_{21} = 4\alpha_1, \qquad \alpha_{31} = 2\alpha_1, \qquad \alpha_{32} = \alpha_2. \tag{5.26}$$

The three remaining undetermined constants $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}$ multiply integrals of the form $J_{N\{000\}}$. Such integrals solve the homogeneous parts of the CWIs. Therefore the remaining constants, undetermined by the primary CWIs, will be called *primary constants*.

Let us summarise our results. We have analysed the primary CWIs for the $\langle\!\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}\mathcal{O}\rangle\!\rangle$ correlation function and we found a solution

$$A_1 = \alpha_1 J_{4\{000\}}, \tag{5.27}$$

$$A_2 = 4\alpha_1 J_{3\{001\}} + \alpha_2 J_{2\{000\}}, \tag{5.28}$$

$$A_3 = 2\alpha_1 J_{2\{002\}} + \alpha_2 J_{1\{001\}} + \alpha_3 J_{0\{000\}}, \tag{5.29}$$

with three undetermined constants $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}$. We will show shortly that this solution to the primary CWIs is indeed unique, specifying $\langle\!\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}\mathcal{O}\rangle\!\rangle$ in momentum space up to three constants. Following application of the secondary CWIs, we will find that the number of undetermined constants is reduced to just one. The method we have described is based purely in momentum space and is applicable to all 3-point functions. Explicit solutions for all primary CWIs are listed in the second part of the paper.

The triple-K integrals we discuss here also arise in AdS/CFT calculations of momentum space 3-point functions using a dual gravitational theory (recent papers include, *e.g.*, [46, 47, 29]). As such, these calculations apply only to the specific CFTs dual to particular gravitational theories. In contrast, our approach here is completely general, showing that all 3-point functions of conserved currents, stress-energy tensors and scalar operators in *any* CFT can be expressed in terms of triple-K integrals.

Finally, let us return to the issue of regularisation. In some cases, despite the finiteness of the final result, the regularisation procedure described in section 5.1.1 may still be necessary. Furthermore, it can happen that single triple-K integrals may diverge on their own while the form factor they build remains finite. It is therefore essential to keep track of the expansion in the regulator ϵ carefully. In particular one must consider the primary constants as functions of the regulator and take the $\epsilon \to 0$ limit only after the substitution into the final expression for the form factors. We will show an example of this behaviour in the next section.

5.2.1 More on $\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}\mathcal{O}\rangle$

In this section we wish to illustrate that the solution to the primary CWIs in terms of the triple-K integrals can be evaluated explicitly with ease. A systematic discussion of the evaluation of

the triple-K integrals will be given in section 6. For concreteness, consider $\langle\!\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}\mathcal{O}\rangle\!\rangle$ with a scalar operator $\mathcal{O}$ of dimension $\Delta_3 = 1$ in $d = 3$ dimensional CFT. The solution to the primary CWIs is given by (5.27) - (5.29) with constants fixed according to (5.26). In order to write the solution explicitly, we can use expressions (A.4.4) and (A.4.5), after which all integrals turn out to be elementary. The following integrals converge and can be easily computed

$$J_{4\{000\}} = I_{\frac{9}{2}\{\frac{3}{2}\frac{3}{2},-\frac{1}{2}\}} = \left(\frac{\pi}{2}\right)^{3/2} \cdot \frac{3(p_1^2+p_2^2)+p_3^2+12p_1p_2+4p_3(p_1+p_2)}{p_3(p_1+p_2+p_3)^4}, \tag{5.30}$$

$$J_{3\{001\}} = I_{\frac{7}{2}\{\frac{3}{2}\frac{3}{2}\frac{1}{2}\}} = \left(\frac{\pi}{2}\right)^{3/2} \cdot \frac{2(p_1^2+p_2^2)+p_3^2+6p_1p_2+3p_3(p_1+p_2)}{(p_1+p_2+p_3)^3}, \tag{5.31}$$

assuming $\Delta_1 = \Delta_2 = 3$ and $\Delta_3 = 1$. The remaining integrals diverge and require a regularisation. As discussed in section 5.1.1, we consider the integrals $J_{N+\epsilon\{k_j\}}$ and we expand the result in ϵ. In this manner, we find

$$\begin{aligned} J_{2+\epsilon\{002\}} = I_{\frac{5}{2}+\epsilon\{\frac{3}{2}\frac{3}{2}\frac{3}{2}\}} = &-\left(\frac{\pi}{2}\right)^{3/2} \frac{1}{(p_1+p_2+p_3)^2}\left[2p_1p_2p_3+p_1^3+p_2^3+p_3^3\right. \\ &\left.+2(p_1^2p_2+p_1p_2^2+p_1^2p_3+p_1p_3^2+p_2^2p_3+p_2p_3^2)\right]+O(\epsilon), \end{aligned} \tag{5.32}$$

$$J_{2+\epsilon\{000\}} = I_{\frac{5}{2}+\epsilon\{\frac{3}{2}\frac{3}{2},-\frac{1}{2}\}} = \left(\frac{\pi}{2}\right)^{3/2} \cdot \frac{1}{p_3\epsilon} + O(\epsilon^0), \tag{5.33}$$

$$J_{1+\epsilon\{001\}} = I_{\frac{3}{2}+\epsilon\{\frac{3}{2}\frac{3}{2}\frac{1}{2}\}} = -\left(\frac{\pi}{2}\right)^{3/2} \cdot \frac{p_3}{\epsilon} + O(\epsilon^0), \tag{5.34}$$

$$J_{0+\epsilon\{000\}} = I_{\frac{1}{2}+\epsilon\{\frac{3}{2}\frac{3}{2},-\frac{1}{2}\}} = -\left(\frac{\pi}{2}\right)^{3/2} \frac{p_1^2+p_2^2-p_3^2}{2p_3\epsilon} + O(\epsilon^0). \tag{5.35}$$

As we will see the omitted terms make no contribution in our subsequent analysis. In order to further constrain the primary constants $\alpha_1, \alpha_2, \alpha_3$ we must consider the secondary CWIs. We will return to this task in section 5.3.1.

At this point we can compare the result given by (5.27) - (5.29) with the direct calculations of the 3-point function for the free scalar field carried out in section 3.4. We see that the form of the integrals $J_{4\{000\}}$, $J_{3\{001\}}$ and $J_{2\{002\}}$ match the form factors A_1, A_2 and A_3 in the equations (3.33) - (3.35). Therefore one finds the primary constants for this particular model to be

$$\alpha_1 = \frac{1}{48}\left(\frac{2}{\pi}\right)^{\frac{3}{2}}, \qquad \alpha_2 = 0, \qquad \alpha_3 = 0. \tag{5.36}$$

Note that the relations (5.26) provide a cross-check on our solution. Later, we will see that the secondary Ward identities impose two additional constraints on the primary constants that are not yet visible.

5.2.2 Uniqueness of the solution

In the previous sections we argued that all CWIs may be solved in terms of triple-K integrals (5.1). A case-by-case analysis confirms this and the list of complete solutions is given in the second part of the paper. Here we want to establish that these solutions are unique. To be more precise, we want to argue that each pair of the primary CWIs determines a form factor A_n uniquely up to one numerical constant. This may be achieved by essentially the same reasoning as in section 2.3.

First, assume that A_n satisfies a pair of homogeneous primary CWIs

$$\mathrm{K}_{12}\, A_n = 0, \qquad\qquad \mathrm{K}_{13}\, A_n = 0, \tag{5.37}$$

together with the dilatation Ward identity (4.11) with tensorial dimension N_n. We can then use the substitution

$$A_n(p_1,p_2,p_3) = p_3^{\Delta_t-2d-N_n}\left(\frac{p_1^2}{p_3^2}\right)^{\mu}\left(\frac{p_2^2}{p_3^2}\right)^{\lambda} F\left(\frac{p_1^2}{p_3^2},\frac{p_2^2}{p_3^2}\right) \tag{5.38}$$

and proceed with the analysis analogous to that following equation (2.15). The substitution leads to the system of equations (2.17, 2.18) with four possible choices of parameters

$$\alpha = \frac{1}{2}\left[N_n + \epsilon_1\left(\Delta_1 - \frac{d}{2}\right) + \epsilon_2\left(\Delta_2 - \frac{d}{2}\right) + \Delta_3\right], \qquad \beta = \alpha - \left(\Delta_3 - \frac{d}{2}\right),$$
$$\gamma = 1 + \epsilon_1\left(\Delta_1 - \frac{d}{2}\right), \qquad \gamma' = 1 + \epsilon_2\left(\Delta_2 - \frac{d}{2}\right), \tag{5.39}$$

parametrised by $\epsilon_1, \epsilon_2 = \pm 1$. We can now use equation (2.22) and the analysis that follows. This leads to the conclusion that the only physically acceptable solution to the homogenous part of the CWIs is given by the triple-K integral $\alpha_n J_{N_n\{000\}}(p_1, p_2, p_3)$, where α_n is a single undetermined constant.

In general, the primary CWIs for a form factor A_n contain inhomogeneous parts. The recursive nature of the primary CWIs discussed in section 4.3.1 then allows us to solve these equations one-by-one. Since the inhomogeneous part is linear in the other form factors, every two solutions to a given pair of equations differ by a solution to the homogeneous part of the equation. The full solution to the pair of primary CWIs and the dilatation Ward identity is therefore unique up to one numerical constant.

It is important to emphasise that while the solution to each pair of primary CWIs is unique up to one primary constant, the representation in terms of triple-K integrals may not be. For example, for generic parameter values the equation (5.18) shows that

$$(\alpha + \beta_t) I_{\alpha-1\{\beta_1\beta_2\beta_3\}} = I_{\alpha\{\beta_1+1,\beta_2,\beta_3\}} + I_{\alpha\{\beta_1,\beta_2+1,\beta_3\}} + I_{\alpha\{\beta_1,\beta_2,\beta_3+1\}}. \tag{5.40}$$

One can therefore rewrite one triple-K integral as a combination of others hence the representation is not unique.

5.3 Solutions to the secondary conformal Ward identities

In this section we will finalise our theoretical considerations by solving the secondary CWIs. In general, the secondary CWIs lead to linear algebraic equations between the various primary constants appearing in solutions to the primary CWIs. The precise form of the secondary CWIs depends on the semi-local information provided by transverse Ward identities, which may be written in terms of the 2-point functions.

We will first return to our example from section 5.2.1 and show how the two secondary CWIs (4.35, 4.36) constrain the values of the three primary constants appearing in the solution (5.27) - (5.29) to the primary CWIs. As expected, we will find two algebraic linear equations between the three primary constants. From this we may conclude that the 3-point function $\langle\langle T^{\mu_1\nu_1} T^{\mu_2\nu_2} \mathcal{O}\rangle\rangle$ depends on a single undetermined primary constant.

Next, we will discuss how the secondary CWIs in the general case lead into a set of algebraic equations for the primary constants. The discussion is sensitive on whether or not the regulator can be thoroughly removed from all triple-K integrals building a given 3-point function. In either cases the procedure is based on taking a zero-momentum limit. In this limit the triple-K integrals simplify and the secondary CWIs can be shown to lead to a set of equations between primary constants.

The procedure is considerably simpler in the case where the regulator can be removed from all triple-K integrals, while in the case one needs to keep the regulator special care must be taken when regulating the 2-point functions that appear in the right-hand side of the secondary CWIs.

5.3.1 $\langle T^{\mu_1\nu_1} T^{\mu_2\nu_2} \mathcal{O}\rangle$ for free scalars

Let us begin by discussing our example correlation function $\langle\langle T^{\mu_1\nu_1} T^{\mu_2\nu_2} \mathcal{O}\rangle\rangle$. We derived the secondary CWIs earlier in (4.35) and (4.36), where the terms on the right-hand side of these

equations are given by the transverse Ward identity (4.59). We now want to show that these data fix two out of three primary constants in the solution (5.27) - (5.29) of the primary CWIs. To fix the final remaining constant then requires additional physical input in the form of the specific field content.

Since the regulator ϵ in the triple-K integrals (5.33) - (5.35) cannot be removed, we must assume that the primary constants α_2 and α_3 depend on the regulator ϵ as well. As remarked earlier in section 5.1.1, while each individual component may depend on the regulator, the full expression for the form factors A_j cannot. Let us therefore define the power series expansions

$$\alpha_j = \sum_{n=-\infty}^{\infty} \alpha_j^{(n)} \epsilon^n, \quad j = 2, 3. \tag{5.41}$$

Since the integral $J_{4\{000\}}$ is finite, we can assume that the constant α_1 does not depend on the regulator, *i.e.*, $\alpha_1 = \alpha_1^{(0)}$.

We start by substituting the solutions (5.27, 5.28) together with the series expansions (5.41) into the secondary CWI (4.35), with right-hand side given by (4.59). Organising the equations according to powers of ϵ, upon sending $\epsilon \to 0$ all equations associated with negative powers of ϵ must vanish. In the present case, this yields $\alpha_2^{(n)} = 0$ for all $n \leq 0$. The equation coming from the ϵ^0 terms then reads

$$-\frac{3}{p_3}\left(\frac{\pi}{2}\right)^{\frac{3}{2}}(\alpha_2^{(1)} + 3\alpha_1^{(0)}) = -\frac{3}{16 p_3}. \tag{5.42}$$

The same procedure may now be applied to the remaining secondary CWI (4.36), yielding $\alpha_3^{(n)} = 0$ for all $n \leq 0$ and

$$\left(\frac{\pi}{2}\right)^{\frac{3}{2}}\left[2\alpha_3^{(1)} + 3\alpha_1^{(0)} - \frac{1}{16}\left(\frac{2}{\pi}\right)^{\frac{3}{2}}\right]\frac{p_1^2 + 3p_2^2 - 3p_3^2}{p_3} + \frac{3}{4}p_3 = \frac{3}{4}p_3. \tag{5.43}$$

Putting everything together, we have

$$\alpha_2 = \left[-3\alpha_1 + \frac{1}{16}\left(\frac{2}{\pi}\right)^{\frac{3}{2}}\right]\epsilon + O(\epsilon^2), \tag{5.44}$$

$$\alpha_3 = \frac{1}{2}\left[-3\alpha_1 + \frac{1}{16}\left(\frac{2}{\pi}\right)^{\frac{3}{2}}\right]\epsilon + O(\epsilon^2), \tag{5.45}$$

where the constant α_1 remains undetermined by Ward identities. When we take the limit $\epsilon \to 0$, the leading terms of order ϵ in these expressions then multiply $1/\epsilon$ poles in the $J_{2+\epsilon\{000\}}$, $J_{1+\epsilon\{001\}}$ and $J_{0+\epsilon\{000\}}$ integrals yielding the correct finite result. The omitted higher order terms in (5.44) and(5.45) make no contribution.

Finally, we can check the results of this section against the result (5.36) for the specific theory discussed in section 5.2.1. Inserting the value of α_1 from (5.36) into (5.44) and (5.45) we indeed recover the correct result $\alpha_2 = \alpha_3 = 0$ up to insignificant $O(\epsilon^2)$ terms.

5.3.2 Simplifications in the generic case

In the previous section we substituted the full solutions to the primary CWIs into the secondary CWIs in order to extract more information about the primary constants. At first sight this procedure might appear hard to carry out in general since the triple-K integrals usually cannot be expressed in terms of elementary functions. It turns out, however, that examining the zero-momentum limit leads to simple algebraic equations for the primary constants.

In this section, for reasons of simplicity, we will assume that each triple-K integral in a solution to the primary CWIs can be defined by an analytic continuation and the regulator can

be completely removed. We will refer to this as the 'generic case' in the present and following sections. We will then analyse the remaining cases later.

In the zero-momentum limit

$$\boldsymbol{p}_3 \to 0, \qquad \boldsymbol{p}_1 = \boldsymbol{p}_2 = \boldsymbol{p}, \tag{5.46}$$

we have

$$p_3^{\beta_3} K_{\beta_3}(p_3 x) = \left[\frac{2^{\beta_3-1}\Gamma(\beta_3)}{x^{\beta_3}} + O(p_3^2)\right] + p_3^{2\beta_3}\left[2^{-\beta_3-1}\Gamma(-\beta_3)x^{\beta_3} + O(p_3^2)\right], \quad \beta_3 \notin \mathbb{Z}, \tag{5.47}$$

$$K_0(p_3 x) = -\log p_3 - \log x + \log 2 - \gamma_E + O(p_3^2), \tag{5.48}$$

$$\begin{aligned} p_3^n K_n(p_3 x) = {} & \left[\frac{2^{n-1}\Gamma(n)}{x^n} + O(p_3^2)\right] \\ & + p_3^{2n}\left[\frac{(-1)^{n+1}}{2^n\Gamma(n+1)} x^n \log p_3 + \text{ultralocal} + O(p_3^2)\right], \quad n = 1, 2, 3, \ldots \end{aligned} \tag{5.49}$$

From these expressions one can see that the zero momentum limit of $p^{\beta_3} K_{\beta_3}(p_3 x)$ exists if $\beta_3 > 0$. Since for any correlation function and any form factor $\beta_3 = \Delta_3 - \frac{d}{2} + k_3$ with non-negative k_3, this condition is satisfied if $\Delta_3 > \frac{d}{2}$. (For conserved currents and for the stress-energy tensor we thus have $\beta_3 > 0$ automatically.) We will return to discuss the case where $\beta_3 \leq 0$ later in the text.

Assuming $\beta_3 > 0$ then, we can calculate the limit of the triple-K integrals in the generic case

$$\lim_{p_3 \to 0} I_{\alpha\{\beta_j\}}(p, p, p_3) = l_{\alpha\{\beta_j\}} \cdot p^{\beta_t - \alpha - 1}, \tag{5.50}$$

where, using the result (A.5.19), we find

$$l_{\alpha\{\beta_k\}} = \frac{2^{\alpha-3}\Gamma(\beta_3)}{\Gamma(\alpha - \beta_3 + 1)} \prod_{\epsilon_1, \epsilon_2 \in \{-1,1\}} \Gamma\left(\frac{\alpha - \beta_3 + 1 + \epsilon_1\beta_1 + \epsilon_2\beta_2}{2}\right), \tag{5.51}$$

which is valid away from poles of the gamma function. Since the derivatives in the L and R operators defined in (4.31) and (4.32) acting on triple-K integrals can also be expressed via (5.12) in terms of triple-K integrals, this procedure leads to algebraic constraints on the primary constants.

5.3.3 Derivation of the equations in the generic case

Let us illustrate the considerations above in the case of the correlator $\langle\!\langle T^{\mu_1\nu_1} T^{\mu_2\nu_2} \mathcal{O} \rangle\!\rangle$. The secondary CWIs are given by (4.35) and (4.36),

$$\mathrm{L}_{2,0}\, A_1 + \mathrm{R}_2\, A_2 = 2d \cdot \text{coeff. of } p_2^{\mu_1} p_3^{\mu_2} p_3^{\nu_2} \text{ in } p_{1\nu_1} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) T^{\mu_2\nu_2}(\boldsymbol{p}_2) \mathcal{O}(\boldsymbol{p}_3) \rangle\!\rangle, \tag{5.52}$$

$$\mathrm{L}_{2,0}\, A_2 + 4\,\mathrm{R}_2\, A_3 = 8d \cdot \text{coeff. of } \delta^{\mu_1\mu_2} p_3^{\mu_2} \text{ in } p_{1\nu_1} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) T^{\mu_2\nu_2}(\boldsymbol{p}_2) \mathcal{O}(\boldsymbol{p}_3) \rangle\!\rangle, \tag{5.53}$$

with $\Delta_1 = \Delta_2 = d$ and L and R defined by (4.31) and (4.32). The right-hand sides are semi-local and can be expressed in terms of 2-point functions by means of the transverse Ward identities. In section 4.4 we found the Ward identity (4.53), which reads

$$p_1^{\nu_1} \langle\!\langle T_{\mu_1\nu_1}(\boldsymbol{p}_1) T_{\mu_2\nu_2}(\boldsymbol{p}_2) \mathcal{O}(\boldsymbol{p}_3) \rangle\!\rangle = 2 p_1^{\nu_1} \langle\!\langle \frac{\delta T_{\mu_1\nu_1}}{\delta g^{\mu_2\nu_2}}(\boldsymbol{p}_1, \boldsymbol{p}_2) \mathcal{O}(\boldsymbol{p}_3) \rangle\!\rangle. \tag{5.54}$$

We omit the group index on $\mathcal{O}$ as we consider only one scalar operator. First, we want to argue that the right-hand side of (5.54) vanishes if $\beta_3 > 0$, unless some specific conditions on conformal dimensions are met. Therefore, in this section we will assume that the right-hand sides of (5.52)

and (5.53) vanish, leaving a discussion of the various special cases to the following sections. Indeed, the only possibility for a non-vanishing right-hand side of (5.54) is if the functional derivative $\delta T_{\mu_1\nu_1}/\delta g^{\mu_2\nu_2}$ contains the operator $\mathcal{O}$ or its descendants. Since the dilatation degree of $\delta T_{\mu_1\nu_1}/\delta g^{\mu_2\nu_2}$ is equal to d, this requires $d = \Delta_3 + 2n$ where n is a non-negative integer. Consider first the case $\Delta_3 = d$. In this case, we can write the most general form of $\delta T_{\mu_1\nu_1}/\delta g^{\mu_2\nu_2}$ which contains $\mathcal{O}$ as

$$\frac{\delta T_{\mu_1\nu_1}(\boldsymbol{x})}{\delta g^{\mu_2\nu_2}(\boldsymbol{y})} = \left[c_1\delta_{\mu_1\nu_1}\delta_{\mu_2\nu_2} + c_2\delta_{(\mu_1(\mu_2}\delta_{\nu_2)\nu_1)}\right]\delta(\boldsymbol{x}-\boldsymbol{y})\mathcal{O}(\boldsymbol{x}) + \dots \tag{5.55}$$

where c_1 and c_2 are numerical constants. If, on the other hand, $d = \Delta_3 + 2n$ with $n > 0$ then derivatives acting on both $\mathcal{O}$ and $\delta(\boldsymbol{x}-\boldsymbol{y})$ may also appear. In all cases, the Fourier transform reads

$$\langle\!\langle \frac{\delta T_{\mu_1\nu_1}}{\delta g^{\mu_2\nu_2}}(\boldsymbol{p}_1,\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle\!\rangle = P_{\mu_1\nu_1\mu_2\nu_2}(p_1^2,p_2^2,p_3^2)\langle\!\langle\mathcal{O}(\boldsymbol{p}_3)\mathcal{O}(-\boldsymbol{p}_3)\rangle\!\rangle, \tag{5.56}$$

where P is some polynomial built from momenta and the metric $\delta_{\mu\nu}$, with kinematic dependence on squares of momenta only. This form arises from the Fourier transform of the position space expression containing derivatives acting on delta functions and on the 2-point function. Since $\langle\!\langle\mathcal{O}(\boldsymbol{p}_3)\mathcal{O}(-\boldsymbol{p}_3)\rangle\!\rangle \sim p_3^{2\beta_3}$, the expression vanishes in the $p_3 \to 0$ limit as long as $\beta_3 > 0$.

We now substitute the solutions of the primary CWIs (5.27, 5.28) into the left-hand side of (5.52) and take the zero-momentum limit. Assuming the regulator can be removed (see section 5.3.6 if not), the result is

$$-\frac{l_{\frac{d}{2}+1\{\frac{d}{2},\frac{d}{2},\Delta_3-\frac{d}{2}\}}}{2}p^{\Delta_3-2}(2+2d-\Delta_3)\left[\alpha_2+\alpha_1(\Delta_3+2)(\Delta_3-d+2)\right] = 0, \tag{5.57}$$

which leads to

$$\alpha_2 = -(\Delta_3+2)(\Delta_3+2-d)\alpha_1. \tag{5.58}$$

The same reasoning as above applied to (5.53) leads to the equation

$$\alpha_3 = \frac{1}{4}\Delta_3(\Delta_3+2)(\Delta_3-d)(\Delta_3+2-d)\alpha_1. \tag{5.59}$$

Summarising, in this and the previous section we presented a method for extracting algebraic dependencies between the primary constants following from the secondary CWIs. The analysis was performed in the generic case, where the regulator can be removed from all triple-K integrals involved. Note that the results (5.58) and (5.59) agree with our example (5.44) and (5.45) in the leading term in ϵ only, *i.e.*, they correctly predict $\alpha_2 = \alpha_3 = 0 + O(\epsilon)$. This is due to the fact that in our example the regulator cannot be removed from each triple-K integral separately. Therefore, it does not satisfy the assumption of this section. Note, however, that the analysis of the generic case is sufficient if one is merely interested in finding the solution up to semi-local terms. This is because the possible non-generic cases arise due to the regularisation procedure, correcting the generic solution by at most semi-local terms.

5.3.4 Triple-K integrals and 2-point functions

Before we discuss the general procedure applicable to all cases, we first need to analyse the possible singularities associated with the 2-point functions. This is because the secondary CWIs connect triple-K integrals with semi-local terms expressible in terms of 2-point functions. Therefore, if the regulator is kept explicitly, the singular terms in triple-K integrals must match the singularities appearing in the 2-point functions.

An initial obstacle is that our convenient regularisation scheme (2.5) does not work for 2-point functions. The Fourier transform of the position space expression for a generic 2-point function is

$$\langle\mathcal{O}(\boldsymbol{x})\mathcal{O}(\boldsymbol{y})\rangle = \frac{c}{|\boldsymbol{x}-\boldsymbol{y}|^{2\Delta}} \;\longmapsto\; \langle\!\langle\mathcal{O}(\boldsymbol{p})\mathcal{O}(-\boldsymbol{p})\rangle\!\rangle = \frac{c\pi^{d/2}2^{d-\alpha}\Gamma\left(\frac{d-2\Delta}{2}\right)}{\Gamma(\Delta)}\cdot p^{2\Delta-d}, \tag{5.60}$$

where Δ is a conformal dimension of a scalar operator $\mathcal{O}$. The singularity occurs if $2\Delta = d+2n$ for a non-negative integer n and is not regularised by the scheme (2.5).

Let us now try to find a different scheme which regularises 2-point functions and it yields results equivalent to (2.5, 5.7) when applied to the expressions entering the left-hand side of the secondary CWIs. It turns out that standard dimensional regularization has this property. We explicitly checked this in all cases discussed in this paper. To motivate this choice consider the following integral

$$\int \frac{\mathrm{d}^d \boldsymbol{k}}{(2\pi)^d} \frac{k^{\mu_1} \dots k^{\mu_r}}{\boldsymbol{k}^{2\delta_3} |\boldsymbol{k}-\boldsymbol{p}_1|^{2\delta_2} |\boldsymbol{k}+\boldsymbol{p}_2|^{2\delta_1}} \tag{5.61}$$

regulated by shifting the space-time dimension while keeping the δ_j parameters fixed. In terms of the α and β parameters in (5.1), this corresponds to $\alpha \mapsto \alpha - \frac{\epsilon}{2}$ and $\beta_j \mapsto \beta_j - \frac{\epsilon}{2}$ as can be seen from (2.4) and (5.2). We can evaluate the integral (5.61) both by the triple-K integrals or the usual Feynman parametrised integrals. This leads to equation (A.3.23), namely

$$\begin{aligned} I_{\alpha\{\beta_1\beta_2\beta_3\}} = 2^{\alpha-3}\Gamma\left(\frac{\alpha-\beta_t+1}{2}\right)\Gamma\left(\frac{\alpha+\beta_t+1}{2}\right)\times \\ \times \int_{[0,1]^3} \mathrm{d}X\, D^{\frac{1}{2}(\beta_t-\alpha-1)} \prod_{j=1}^{3} x_j^{\frac{1}{2}(\alpha-1-\beta_t)+\beta_j}, \end{aligned} \tag{5.62}$$

where

$$\mathrm{d}X = \mathrm{d}x_1\mathrm{d}x_2\mathrm{d}x_3\delta(x_1+x_2+x_3-1), \qquad D = p_1^2 x_2 x_3 + p_2^2 x_1 x_3 + p_3^2 x_1 x_2. \tag{5.63}$$

The leading divergence comes from a possible zero under the first gamma function and both schemes regularise $\alpha-\beta_t+1$ in the same way, shifting it by $\epsilon/2$.

A detailed analysis shows that all singularities and the finite part (modulo terms analytic in all momenta) in the expressions appearing in the left-hand side of the secondary CWIs match between the two schemes (*i.e.*, dimensional regularisation, $d \mapsto d-\epsilon$, and the scheme (2.5, 5.7)). Thus we can consistently take into account the contributions of the 2-point functions by computing them using dimensional regularization.

In the renormalised theory the divergent terms can be removed. Therefore we define constants $c_{\mathcal{O}}$, c_J and c_T by

$$\langle\!\langle \mathcal{O}^I(\boldsymbol{p})\mathcal{O}^J(-\boldsymbol{p})\rangle\!\rangle = c_{\mathcal{O}}\delta^{IJ} \times \begin{cases} p^{2\Delta-d} & \text{if } 2\Delta \neq d+2n,\ n\in\mathbb{N}, \\ p^{2\Delta-d}\left(-\log p^2 + \text{local}\right) & \text{if } 2\Delta = d+2n,\ n\in\mathbb{N} \end{cases} \tag{5.64}$$

$$\langle\!\langle J^{\mu a}(\boldsymbol{p})J^{\nu b}(-\boldsymbol{p})\rangle\!\rangle = c_J\delta^{ab} \times \begin{cases} \pi^{\mu\nu}(\boldsymbol{p})p^{d-2} & \text{if } d=3,5,7,\dots \\ \pi^{\mu\nu}(\boldsymbol{p})p^{d-2}\left(-\log p^2 + \text{local}\right) & \text{if } d=4,6,8,\dots \end{cases} \tag{5.65}$$

$$\langle\!\langle T^{\mu\nu}(\boldsymbol{p})T^{\rho\sigma}(-\boldsymbol{p})\rangle\!\rangle = c_T \times \begin{cases} \Pi^{\mu\nu\rho\sigma}(\boldsymbol{p})p^{d} & \text{if } d=3,5,7,\dots \\ \Pi^{\mu\nu\rho\sigma}(\boldsymbol{p})p^{d}\left(-\log p^2 + \text{local}\right) & \text{if } d=4,6,8,\dots \end{cases} \tag{5.66}$$

where all $\mathcal{O}^I$ operators have the same conformal dimension Δ and the dimensions of $J^{\mu a}$ and $T^{\mu\nu}$ are $d-1$ and d respectively. In general the normalisation constants $c_{\mathcal{O}}$ and c_J carry group indices. For simplicity we assume that the Killing form is trivial, *i.e.*,

$$c^{IJ}_{\mathcal{O}} = c_{\mathcal{O}}\delta^{IJ}, \qquad c^{ab}_J = c_J\delta^{ab}. \tag{5.67}$$

5.3.5 Secondary conformal Ward identities in all cases

Let us now return to the discussion of the secondary CWIs in the case where the regulator cannot be removed in certain triple-K integrals. In principle, the procedure is simple. One must keep the explicit dependence on ϵ, both in the triple-K integrals as well as in the primary constants, and carry out the analysis of sections 5.3.2 and 5.3.3 order by order in the regulator.

Note that if the index of a Bessel function is integral, then the expansions (5.48) and (5.49) should be used instead of (5.47).

The only difference with section 5.3.3 is that looking at the zero-momentum limit may not be enough. We should look at both terms following from the first and second brackets in (5.47), *i.e.*, the coefficients of p_3^0 and $p_3^{2\beta_3}$ in the expansion in powers of p_3 with $p_1 = p_2 = p$. If the Bessel index is integral, then we should use (5.48) and (5.49) and look for the coefficients of p_3^0 and $p_3^{2\beta_3}\log p_3$. This procedure will provide a set of algebraic equations relating the primary constants.

Let us now explain why this procedure is valid for $\beta_3 < 0$ and why the remaining terms in the expansions (5.47) - (5.49) are irrelevant. First, the unitarity bound requires $-1 \leq \beta_3$. The unitarity bound can only be saturated by a non-composite scalar operator in a free field theory, [8] and [48]. We can therefore assume $-1 < \beta_3 < 0$. It turns out that the considerations encountered in the case $\beta_3 > 0$ remain valid here. Since the zero-momentum limit does not exist in this case, we are going to look for the coefficient of p_3^0 in the expansion in p_3. The key observation is that on the left-hand sides of the secondary CWIs such as (5.52, 5.53), the differential operators L and R defined by (4.31) and (4.32) do not contain derivatives with respect to p_3, and can only increase powers of p_3 by two. Therefore, the coefficient of p_3^0 in the series expansion in p_3 remains unaltered provided $-1 < \beta_3$. A similar analysis applies to the right-hand sides of the secondary CWIs.

Let us now examine why it is sufficient to look at the leading coefficients in (5.47) - (5.49) only. From (A.4.1), we know that in each successive term the power of the integration variable x increases by two. After taking the zero-momentum limit, the integral (A.5.19) therefore leads to essentially the same expression as (5.51) with $\beta_3 \mapsto \beta_3 + 2n$, $n \in \mathbb{N}$, plus some finite pre-factor following from the series expansion of the Bessel K function. Since the singularities manifest themselves as poles of the gamma functions, we see that the result cannot be more singular than the original $l_{\alpha\{\beta_j\}}$.

5.3.6 Back to the example

Finally, let us see how the general consideration of the previous section work in the case of the $\langle\!\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}\mathcal{O}\rangle\!\rangle$ correlation function. First, we carry out the same analysis as in sections 5.3.2 and 5.3.3 but keep the regulator explicitly. Instead of (5.57), we then find

$$-\frac{1}{2}l_{\frac{d}{2}+1+\epsilon\{\frac{d}{2},\frac{d}{2},\Delta_3-\frac{d}{2}\}}p^{\Delta_3-2+\epsilon}(2+2d-\Delta_3+\epsilon)\times \\ \times\left[a_2+a_1((\Delta_3-\epsilon)^2-d(\Delta+2-\epsilon)+4(1+\Delta+\epsilon))\right]=0. \tag{5.68}$$

If the $\epsilon \to 0$ limit exists, we recover (5.57). The limit does not exist, however, if Δ_3 satisfies some non-generic relations. From the definition of $l_{\frac{d}{2}+1+\epsilon\{\frac{d}{2},\frac{d}{2},\Delta_3-\frac{d}{2}\}}$ in (5.51), we see that this happens if at least one of the following conditions are satisfied:

- $\Delta_3 = 2 + 2n_1$, $n_1 \in \mathbb{N}$,
- $\Delta_3 = d + 2 + 2n_2$, $n_2 \in \mathbb{N}$,
- $\Delta_3 = d + 4 + 2n_3$, $n_3 \in \mathbb{N}$,
- $\Delta_3 = 2d + 4 + 2n_4$, $n_4 \in \mathbb{N}$.

The order of the singularity increases if more than one of these conditions is satisfied. Since there exists a choice of Δ_3, d and the n_j constants such that all conditions are satisfied, we might expect a pole of order four in ϵ. Note however that there is another gamma function in the denominator of (5.51) which becomes singular if the numerator has a pole of order four. The maximal order of the pole is therefore only three.

The discussion above leads to the conclusion that we should expand the primary constant α_2 up to third order in ϵ as in (5.41), hence

$$\begin{aligned}\alpha_2 J_{2\{000\}} = &\frac{1}{\epsilon^3}\alpha_2^{(0)} J^{(-3)}_{2\{000\}} + \frac{1}{\epsilon^2}\left[\alpha_2^{(0)} J^{(-2)}_{2\{000\}} + \alpha_2^{(1)} J^{(-3)}_{2\{000\}}\right] \\ &+ \frac{1}{\epsilon}\left[\alpha_2^{(0)} J^{(-1)}_{2\{000\}} + \alpha_2^{(1)} J^{(-2)}_{2\{000\}} + \alpha_3^{(2)} J^{(-3)}_{2\{000\}}\right] \\ &+ \left[\alpha_2^{(0)} J^{(0)}_{2\{000\}} + \alpha_2^{(1)} J^{(-1)}_{2\{000\}} + \alpha_2^{(2)} J^{(-2)}_{2\{000\}} + \alpha_2^{(3)} J^{(-3)}_{2\{000\}}\right] + O(\epsilon). \end{aligned} \tag{5.69}$$

With the appropriate choice of primary constants we will now see that the singular terms in the various triple-K integrals building a given form factor cancel out, leaving ultralocal singular terms at most.

Let us now extract the primary constants from the secondary CWI (4.35). We substitute (5.69) into (5.68) and expand the result in ϵ in any possible combination of the cases itemised above. One can subsequently solve the equations starting from the most singular one. In this way, one finds

$$\begin{aligned} \alpha_2^{(0)} &= -(\Delta_3+2)(\Delta_3+2-d)\alpha_1, & \alpha_2^{(2)} &= -\alpha_1, \\ \alpha_2^{(1)} &= (2\Delta_3 - d + 4)\alpha_1, & \alpha_2^{(3)} &= 0. \end{aligned} \tag{5.70}$$

We assume here that α_1 is a true constant, *i.e.*, $\alpha_1 = \alpha_1^{(0)}$. These solutions are valid for the special cases listed at the beginning of this section. Notice that they still do not cover all special cases, for example they do not cover the case of the example we studied in section 5.3.1. This is because so far we have considered the equations following from the first parts of the expansions (5.47) - (5.49) only: we must now turn our attention to the equations following from the second parts. In many cases the equations to follow will agree with (5.70), but in some special cases new contributions will arise.

The equation (5.47) and the integral (A.5.19) lead to the equation

$$\begin{aligned} &-\frac{l_{\frac{d}{2}+1+\epsilon\{\frac{d}{2},\frac{d}{2},\frac{d}{2}-\Delta_3\}}}{2}(2+d+\Delta_3+\epsilon)\left[a_2 + a_1(\Delta_3+2+\epsilon)(\Delta_3-d+2+\epsilon)\right] = \\ &\quad = 4d\cdot\text{coefficient of } p_2^{\mu_1}p_3^{\mu_2}p_3^{\nu_2}\cdot p_3^{2\Delta_3-d}p^{d-\Delta_3-2} \text{ in } p_2^{\nu_1}\langle\!\langle \frac{\delta T_{\mu_1\nu_1}}{\delta g^{\mu_2\nu_2}}(\boldsymbol{p}_1,\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle \end{aligned} \tag{5.71}$$

following from the coefficient of $p_3^{2\beta_3}$ in the series expansion (5.47). First note that if the $\epsilon\to 0$ limit exists, then the left-hand side vanishes when the solution (5.58) is substituted. On the other hand, we know from section 5.3.4 that the right-hand side can be non-zero only if $\Delta_3 = d-2-2n$ for some non-negative integer n. Therefore, in such a case we expect the left-hand side to be more singular than the right-hand side, so that the solution (5.58) cancels the leading order singularity while the sub-leading terms match the right-hand side. Indeed, the left-hand side is singular if $l_{\frac{d}{2}+1\{\frac{d}{2},\frac{d}{2},\frac{d}{2}-\Delta_3\}}$ is singular. Analysing the expression (5.51), we see that this can happen only if $\Delta_3 = d-2-2n$, $n\in\mathbb{N}$. Note that our example analysed in section 5.3.1 is of this kind as there $d=3$, $\Delta_3=1$.

Finally, we would like to extract the sub-leading equations in the case where $\Delta_3 = d-2-2n$. In order to do this, we write $\alpha_2 = \alpha_2^{(0)} + \epsilon\alpha_2^{(1)}$ in (5.71) and expand the result in ϵ. At zeroth order we recover (5.58), after which we find

$$C\cdot\Gamma\left(\frac{d}{2}-\Delta_3\right)\left[(2\Delta_3-d+4)\alpha_1 + \alpha_2^{(1)}\right] = 4d\cdot c_1^I c_{\mathcal{O}}, \tag{5.72}$$

where

$$C = \frac{(-1)^{\frac{d-\Delta_3}{2}}2^{\frac{d}{2}-\Delta_3-2}\sqrt{\pi}\,\Gamma\left(\frac{\Delta_3+2}{2}\right)\Gamma\left(\frac{\Delta_3+d+4}{2}\right)}{\Gamma\left(\frac{d-\Delta_3}{2}\right)\Gamma\left(\frac{\Delta_3+3}{2}\right)} \tag{5.73}$$

and the constant c_1^K is defined as

$$p_1^{\nu_1}\langle\!\langle\frac{\delta T_{\mu_1\nu_1}}{\delta g^{\mu_2\nu_2}}(\boldsymbol{p}_1,\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p})_3\rangle\!\rangle\Big|_{p_1=p_2=p} = c_1^K p_2^{\mu_1}p_3^{\mu_2}p_3^{\nu_2}p^{d-\Delta_3-2}\langle\!\langle\mathcal{O}^I(\boldsymbol{p}_3)\mathcal{O}^K(-\boldsymbol{p}_3)\rangle\!\rangle + c_2^K\delta^{\mu_1\mu_2}p_3^{\nu_2}p^{d-\Delta_3}\langle\!\langle\mathcal{O}^I(\boldsymbol{p}_3)\mathcal{O}^K(-\boldsymbol{p}_3)\rangle\!\rangle + \dots \quad (5.74)$$

By $p_1 = p_2 = p$, we mean here the following procedure: First, the correlation function on the left-hand side is expanded in terms of simple tensors according to the convention (3.7), then secondly, one applies $p_1 = p_2 = p$ to each coefficient separately.

In order to derive (5.72) we used (5.47). If $\Delta_3 = d - 2 - 2n$ and $2\Delta_3 = d + 2m$ for some $m, n \in \mathbb{N}$, *i.e.*, if β_3 is an integer, then one must use instead (5.49). The procedure remains identical and the final result is

$$-\frac{C}{\Gamma\left(\Delta_3 - \frac{d}{2} + 1\right)}\left[(2\Delta_3 - d + 4)\alpha_1 + \alpha_2^{(1)}\right] = 4d \cdot c_1^I c_{\mathcal{O}}. \quad (5.75)$$

In total, we have that

$$\alpha_2^{(0)} = -(\Delta_3 + 2)(\Delta_3 + 2 - d)\alpha_1 \quad (5.76)$$

in all cases while

$$\alpha_2^{(1)} = (2\Delta_3 - d + 4)\alpha_1, \qquad \alpha_2^{(2)} = -\alpha_1 \quad (5.77)$$

if $\Delta_3 \neq d - 2 - 2n$, $n \in \mathbb{N}$ and

$$4d \cdot c_1^I c_{\mathcal{O}} = \left[(2\Delta_3 - d + 4)\alpha_1 + \alpha_2^{(1)}\right] \cdot C \cdot \begin{cases} \Gamma\left(\frac{d}{2} - \Delta_3\right) & \text{if } 2\Delta_3 \neq d + 2m,\ m \in \mathbb{N}, \\ -\dfrac{1}{\Gamma\left(\Delta_3 - \frac{d}{2} + 1\right)} & \text{if } 2\Delta_3 = d + 2m,\ m \in \mathbb{N}, \end{cases} \quad (5.78)$$

if $\Delta_3 = d - 2 - 2n$, $n \in \mathbb{N}$.

A similar analysis may be carried out for the second CWI, (5.53). Putting all the ingredients together, we can now write the most general form of the correlator $\langle\!\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}\mathcal{O}^I\rangle\!\rangle$ for $d = 3$ and $\Delta_2 = \Delta_3 = 1$. Using the results for the triple-K integrals from section 5.2.1 we have

$$A_1^I = \frac{\alpha_1^I}{p_3 a_{123}^4}\left[p_3^2 + 4p_3 a_{12} + 3(a_{12}^2 + 2b_{12})\right], \quad (5.79)$$

$$A_2^I = \frac{\alpha_1^I}{p_3 a_{123}^3}\left[p_3^3 + 3p_2^2 a_{12} + p_3(-a_{12}^2 + 8b_{12}) - 3a_{12}^3\right] - \frac{4c_1^I c_{\mathcal{O}}}{p_3}, \quad (5.80)$$

$$A_3^I = \frac{\alpha_1^I(a_{12} - p_3)}{4p_3 a_{123}^2}\left[-p_3^3 - 3p_3^2 a_{12} + p_3(a_{12}^2 - 10b_{12}) + 3a_{12}(a_{12}^2 - 2b_{12})\right] + \frac{c_{\mathcal{O}}}{p_3}\left[(c_1^I - 3c_2^I)(p_1^2 + p_2^2) + 3(c_1^I + c_2^I)p_3^2\right], \quad (5.81)$$

where we have defined the symmetric polynomials in momentum magnitudes

$$a_{123} = p_1 + p_2 + p_3, \qquad b_{123} = p_1p_2 + p_1p_3 + p_2p_3, \qquad c_{123} = p_1p_2p_3, \quad (5.82)$$

$$a_{ij} = p_i + p_j, \qquad b_{ij} = p_ip_j, \quad (5.83)$$

where $i, j = 1, 2, 3$. The solution for this correlator is thus uniquely determined up to one numerical constant α_1^I. The remaining constants in the solution, namely $c_{\mathcal{O}}$, c_1^I and c_2^I, are determined by the 2-point function normalisations: $c_{\mathcal{O}}$ is given in (5.64) while c_1^I and c_2^I are given in (5.74).

One can check this result against our example in section 5.3.1. From (4.58) and (4.59), the solution for the parameters is

$$c_{\mathcal{O}} = \frac{1}{4}, \qquad c_1^I = -\frac{1}{16}, \qquad c_2^I = 0. \quad (5.84)$$

6 Evaluation of triple-K integrals

In the preceding sections we have developed a method for calculating all 3-point functions of the stress-energy tensor, conserved currents and scalar operators in any CFT. The method operates purely in momentum space, and is based on a direct solution of the conformal Ward identities. The results we obtain are expressed in the form of triple-K integrals.

In the present section we now turn to discuss the evaluation of these triple-K integrals. The ease with which this may be accomplished depends on the dimensionality of the space d. In an odd number of dimensions, it turns out that all 3-point functions of conserved currents and the stress-energy tensor are expressible in terms of triple-K integrals in which the Bessel function indices are half-integer. In this case, the Bessel functions reduce to elementary functions (see appendix A.4) and the triple-K integral may be straightforwardly evaluated. When the spatial dimension d is even, we obtain instead triple-K integrals in which the Bessel function indices are integer. In this case, the triple-K integrals we encounter may be evaluated in terms of a single master integral through the use of a reduction scheme. In the following we will focus primarily on the case $d = 4$, but method we present extends straightforwardly to higher even dimensions.

Our reduction scheme is based on the observation that, given a triple-K integral $I_{\alpha\{\beta_1\beta_2\beta_3\}}$, through differentiation we may easily obtain the integrals $I_{\alpha+1\{\beta_1+1,\beta_2,\beta_3\}}$ and $I_{\alpha+1\{\beta_1-1,\beta_2,\beta_3\}}$, and similarly for β_2 and β_3 by permutation. We may then start with a known integral with a sufficiently small value of α and obtain integrals with larger α through repeated differentiation. In some cases a relation for lowering α also exists. For 3-point functions of conserved currents and stress-energy tensors in $d = 4$ the necessary triple-K integrals are:

$$I_{4\{111\}} \mapsto I_{5\{211\}}, I_{6\{221\}}, I_{7\{222\}}, \tag{6.1}$$

$$I_{2\{111\}} \mapsto I_{3\{211\}}, I_{4\{221\}}, I_{4\{311\}}, I_{5\{222\}}, I_{5\{321\}}, I_{6\{322\}}, \tag{6.2}$$

$$I_{0\{111\}} \mapsto I_{1\{211\}}, I_{2\{221\}}, I_{3\{222\}}, I_{3\{321\}}, I_{4\{322\}}, I_{5\{422\}}, I_{5\{332\}}, \tag{6.3}$$

$$I_{1\{222\}} \mapsto I_{3\{332\}}, I_{4\{333\}}. \tag{6.4}$$

The integrals are organised into four families, each with constant $\alpha - \beta_t$, where each integral within a given family may be obtained from the corresponding integral on the leftmost end. The four leftmost integrals can then be derived from the single master integral, $I_{0\{111\}}$, as we will show in section 6.4.

6.1 Reduction scheme

The elementary properties of Bessel functions imply

$$I_{\alpha\{\beta_{\sigma(1)}\beta_{\sigma(2)}\beta_{\sigma(3)}\}}(p_1,p_2,p_3) = I_{\alpha\{\beta_1\beta_2\beta_3\}}(p_{\sigma^{-1}(1)},p_{\sigma^{-1}(2)},p_{\sigma^{-1}(3)}), \tag{6.5}$$

$$I_{\alpha\{\beta_1\beta_2\beta_3\}} = -\frac{1}{p_1}\frac{\partial}{\partial p_1} I_{\alpha-1\{\beta_1+1,\beta_2\beta_3\}}, \tag{6.6}$$

$$I_{\alpha\{\beta_1\beta_2,-\beta_3\}} = p_3^{-2\beta_3} I_{\alpha\{\beta_1\beta_2\beta_3\}}, \tag{6.7}$$

$$(\alpha-\beta_t) I_{\alpha-1\{\beta_1\beta_2\beta_3\}} = p_1^2 I_{\alpha\{\beta_1-1,\beta_2,\beta_3\}} + p_2^2 I_{\alpha\{\beta_1,\beta_2-1,\beta_3\}} + p_3^2 I_{\alpha\{\beta_1,\beta_2,\beta_3-1\}}, \quad \alpha-\beta_t \neq -2n,\ n\in\mathbb{N}, \tag{6.8}$$

where the triple-K integral $I_{\alpha\{\beta_1\beta_2\beta_3\}}$ was defined in (5.1). The first of these equations appeared previously as (5.11) and simply expresses the fact that the triple symmetry under permutation, with σ representing a permutation of the set $\{1,2,3\}$. The second and third of these equations appeared earlier as (5.13) and (5.14), while the last follows from the second line of (5.18). Starting from these relations, we may now set up our reduction scheme as follows.

First, assume we are given an integral $I_{\alpha\{\beta_1\beta_2\beta_3\}}$. Applying (6.6) in the form

$$I_{\alpha+1\{\beta_1-1,\beta_2,\beta_3\}} = -\frac{1}{p_1}\frac{\partial}{\partial p_1} I_{\alpha\{\beta_1\beta_2\beta_3\}}, \tag{6.9}$$

we increase α by one while decreasing β_1 by one. Equivalently, this operation increases the difference $\alpha - \beta_t$ by two but keeps the sum $\alpha + \beta_t$ fixed.

If instead we first use equation (6.7), followed by (6.6) then (6.7) again, we find

$$\begin{aligned} I_{\alpha+1\{\beta_1+1,\beta_2,\beta_3\}} &= -p_1^{2\beta_1+1}\frac{\partial}{\partial p_1}\left[p_1^{-2\beta_1} I_{\alpha\{\beta_1\beta_2\beta_3\}}\right] \\ &= \left(2\beta_1 - p_1\frac{\partial}{\partial p_1}\right) I_{\alpha\{\beta_1\beta_2\beta_3\}}, \end{aligned} \tag{6.10}$$

where the sum $\alpha+\beta_t$ increases by two but the difference $\alpha-\beta_t$ remains fixed. Similarly, applying this operation and its permutations repeatedly we obtain

$$I_{\alpha+n\{\beta_j+k_j\}} = (-1)^{k_t}\left[\prod_{j=1}^{3} p_j^{2(\beta_j+k_j)}\left(\frac{1}{p_j}\frac{\partial}{\partial p_j}\right)^{k_j}\right]\left[p_1^{-2\beta_1}p_2^{-2\beta_2}p_3^{-2\beta_3} I_{\alpha\{\beta_j\}}\right]. \tag{6.11}$$

where $k_t = \sum_j k_j$ and the k_j are non-negative integers.

Now, both reduction relations (6.6) and (6.7) obtained by differentiation happen to increase α by one. To reduce α we may instead use the (6.8) in the form

$$I_{\alpha-1\{\beta_1\beta_2\beta_3\}} = \frac{1}{\alpha-\beta_t}\left[p_1^2 I_{\alpha\{\beta_1-1,\beta_2,\beta_3\}} + p_2^2 I_{\alpha\{\beta_1,\beta_2-1,\beta_3\}} + p_3^2 I_{\alpha\{\beta_1,\beta_2,\beta_3-1\}}\right], \tag{6.12}$$

assuming $\alpha - \beta_t \neq -2n$, where n is a non-negative integer. This equation is closely related to Davydychev's recursion relation (3.4) introduced in [33]. Indeed, using (A.3.17) one can rewrite Davydychev's J integral defined in (2.1) of [33] as

$$J(\delta_1,\delta_2,\delta_3) = \frac{4\pi^2}{\Gamma(\delta_1)\Gamma(\delta_2)\Gamma(\delta_3)\Gamma(4-\delta_t)} I_{1\{2-\delta_2-\delta_3,2-\delta_1-\delta_3,2-\delta_1-\delta_2\}}. \tag{6.13}$$

Note that the rather complicated form of equation (3.4) in [33] is a consequence of the specific index structure in the triple-K integral above.

6.2 The case $d = 4$

Let us now apply the reduction relations above to the integrals listed in (6.1) - (6.4). Integrals within a given family have constant $\alpha-\beta_t$ and hence are connected by (6.10). Similarly, integrals belonging to different families but with the same sum $\alpha + \beta_t$ may be connected using (6.9). We summarise in table 1 the dependencies between all integrals necessary for the evaluation of 3-point functions of conserved currents and the stress-energy tensor in $d = 4$.

Integrals within a single row (except for the first) are connected via equation (6.9), while integrals within a single column are related by (6.10). The difference $\alpha - \beta_t$ is thus constant within each column while the sum $\alpha + \beta_t$ is constant along each row. Similarly, the index α is constant along diagonals from top-right to bottom-left, while β_t is constant along diagonals from the top-left to the bottom-right. In cases where there are two integrals in a given table entry, the arrows entering and leaving this entry have two labels, with the upper label referring to the upper integral and the lower label referring to the lower integral, thus, *e.g.*,

$$I_{3\{222\}} = -\frac{1}{p_1}\frac{\partial}{\partial p_1} I_{2\{322\}}, \qquad I_{3\{321\}} = -\frac{1}{p_3}\frac{\partial}{\partial p_3} I_{2\{322\}}. \tag{6.14}$$

We assume all integrals are regularised in our standard scheme (5.7). In this case, the operations assigned to the arrows should be applied order by order in the regulator. Note that




Table 1: *Reduction scheme for the integrals required in the calculations of 3-point functions of conserved currents and stress-energy tensor in $d = 4$. All integrals can be obtained as indicated from a single master integral $I_{0\{111\}}$ given by* (A.6.1). *The dotted line indicates the use of* (6.12).

if one uses a different regularisation scheme which changes the values of the β parameters, one cannot apply the operators on the vertical lines order by order in ϵ. This is because (6.10) contains a β parameter which would become a function of the regulator.

In section 6.4 we will evaluate the master integral $I_{0\{111\}}$. However, as the entries on the left side of the table are generally more singular (and more complicated) than the entries on the right, in practice it is more convenient to move in columns rather than in rows. The required expressions for the three integrals $I_{0\{111\}}$, $I_{2\{111\}}$ and $I_{1\{000\}}$, which generate all the integrals in the three rightmost columns are given in appendix A.6.

Starting from $I_{0+\epsilon\{111\}}$ we can follow the arrows in table 1 to obtain $I_{2+\epsilon\{221\}}$. Using (6.5)

and (6.12) we find

$$\begin{aligned}
I_{1+\epsilon\{222\}} &= \frac{1}{\epsilon-4}\left[p_1^2 I_{2+\epsilon\{122\}} + p_2^2 I_{2+\epsilon,\{212\}} + p_3^2 I_{2+\epsilon\{221\}}\right] \\
&= \frac{p_1^4+p_2^4+p_3^4}{2\epsilon^2} \\
&\quad + \frac{1}{8}\left(\frac{4}{\epsilon}+1\right)\times\left[\left(p_1^2p_2^2 + (\tfrac{3}{4}-\gamma_E+\log 2)p_3^4 - p_3^4\log p_3\right)\right. \\
&\qquad\qquad \left. + (p_1\leftrightarrow p_3) + (p_2\leftrightarrow p_3)\right] \\
&\quad - \frac{1}{4}\left[p_3^2\left(2-p_1\frac{\partial}{\partial p_1}\right)\left(2-p_2\frac{\partial}{\partial p_2}\right) + (p_1\leftrightarrow p_3) + (p_2\leftrightarrow p_3)\right] I^{(0)}_{0\{111\}} \\
&\quad + O(\epsilon),
\end{aligned} \tag{6.15}$$

where $I^{(0)}_{0\{111\}}$ denotes the ϵ^0 term in the series expansion in ϵ of $I_{0+\epsilon\{111\}}$. Thus all integrals in the table can be obtained from $I_{0\{111\}}$.

6.3 Integrals in even dimensions $d \geq 4$

In the previous section we presented a comprehensive procedure for the evaluation of all triple-K integrals appearing in 3-point functions of conserved currents and the stress-energy tensor in $d = 4$. In this section we want to extend our analysis to all even dimensions $d \geq 4$. We will present a recursive procedure which yields all required integrals from the master integral $I_{0\{111\}}$.

Let us start with the case $d = 6$. Looking at the solutions to the primary CWIs, one sees that new integrals arise only in the form factor A_5 of the $\langle\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}T^{\mu_3\nu_3}\rangle\rangle$ correlator. One of these new integrals, $J_{0\{000\}}$, is equal to $I_{2\{333\}}$ when $d = \Delta_j = 6$, $j = 1, 2, 3$ is used in (5.3). All remaining integrals in the form factor A_5 then satisfy the relation $\alpha - \beta_t = -7$ meaning they can be obtained from $I_{2\{333\}}$ using (6.10). Finally, notice that by (6.9),

$$-\frac{1}{p_3}\frac{\partial}{\partial p_3} I_{2\{333\}} = I_{3\{332\}}. \tag{6.16}$$

Thus to obtain all required integrals in $d = 6$ we need to add another column to the left side of table 1.

This discussion generalises for general even d. Assume first that we have all integrals required for the evaluation of the 3-point functions of conserved currents and the stress-energy tensor in a given even dimension. The leftmost integrals in table 1 have the lowest value of $\alpha - \beta_t$ and it is these integrals which appear in the A_5 form factor in the correlator of three stress-energy tensors. For these integrals $\alpha - \beta_t = -(d+1)$, and the lowest value of $\alpha + \beta_t$ is attained for the $J_{0\{000\}}$ integral. For the correlator of three stress-energy tensors in d dimensions

$$J_{0\{000\}} = I_{\frac{d}{2}-1\{\frac{d}{2}\frac{d}{2}\frac{d}{2}\}} \tag{6.17}$$

and all remaining integrals in the form factor A_5 can be obtained from this one by means of (6.10). Therefore, if one knows the integrals in a dimension $d-2$, in order to obtain the missing integrals in dimension d, one needs to reduce the value of $\alpha-\beta_t$. This can be achieved by (6.12),

$$I_{\frac{d}{2}-1+\epsilon\{\frac{d}{2}\frac{d}{2}\frac{d}{2}\}} = -\frac{1}{d-\epsilon}\left[p_1^2 I_{\frac{d}{2}+\epsilon\{\frac{d}{2}-1,\frac{d}{2},\frac{d}{2}\}} + p_2^2 I_{\frac{d}{2}+\epsilon\{\frac{d}{2},\frac{d}{2}-1,\frac{d}{2}\}} + p_3^2 I_{\frac{d}{2}+\epsilon\{\frac{d}{2},\frac{d}{2},\frac{d}{2}-1\}}\right]. \tag{6.18}$$

If one denotes $d = d' + 2$, then the integrals featuring in this expression can be obtained from the integral (6.17) in dimension d' as follows

$$\begin{aligned}
I_{\frac{d}{2}+\epsilon\{\frac{d}{2}-1,\frac{d}{2},\frac{d}{2}\}} &= I_{\frac{d'}{2}+1+\epsilon\{\frac{d'}{2},\frac{d'}{2}+1,\frac{d'}{2}+1\}} \\
&= \left(d' - p_2\frac{\partial}{\partial p_2}\right)\left(d' - p_3\frac{\partial}{\partial p_3}\right) I_{\frac{d'}{2}-1+\epsilon\{\frac{d'}{2}\frac{d'}{2}\frac{d'}{2}\}}.
\end{aligned} \tag{6.19}$$

This shows that the recursive use of (6.19) and (6.10) allows analytic expressions to be found for all triple-K integrals required for the evaluation of 3-point functions of conserved currents and the stress-energy tensor in arbitrary even dimension $d \geq 4$. Visually, the procedure adds new columns to the left side of table 1 with lower values of $\alpha - \beta_t$. One can move down the column through the repeated use of (6.10), and the starting entry in each column is (6.17). In summary, one can find all required integrals starting from a single master integral $I_{0\{111\}}$ which we will evaluate analytically in the following section.

6.4 Evaluation of the master integral

In this section we present a method to evaluate integrals of the form $I_{\nu+1\{\nu\nu\nu\}}$, $\nu \in \mathbb{R}$. In particular, if we choose $\nu = -1$ to evaluate $I_{0\{-1-1-1\}}$, using (6.7) then gives

$$I_{0\{111\}} = p_1^2 p_2^2 p_3^2 I_{0\{-1-1-1\}} \tag{6.20}$$

which is the master integral we will need for the analysis of conserved currents and stress tensors.

In appendix A.6 we also present expressions for $I_{2\{111\}}$ and $I_{1\{000\}}$. In particular, $I_{1\{000\}}$ is convergent and finite and is known in the literature, *e.g.*, [33, 49].

Let us now evaluate the master integral $I_{0\{111\}}$. We will start with the more general problem of evaluating integrals of the form $I_{\nu+1\{\nu\nu\nu\}}$, $\nu \in \mathbb{R}$. To write the results in compact form, we introduce the following variables. First, we define

$$\begin{aligned}\lambda^2 &= (p_1 + p_2 - p_3)(p_1 - p_2 + p_3)(-p_1 + p_2 + p_3)(p_1 + p_2 + p_3) \\ &= -p_1^4 - p_2^4 - p_3^4 + 2p_1^2 p_2^2 + 2p_1^2 p_3^2 + 2p_2^2 p_3^2 \\ &= 4\left[p_1^2 p_2^2 - (\boldsymbol{p}_1 \cdot \boldsymbol{p}_2)^2\right] = 4 \cdot \mathrm{Gram}(\boldsymbol{p}_1, \boldsymbol{p}_2),\end{aligned} \tag{6.21}$$

where Gram is the Gram determinant. For physical momentum configurations obeying the triangle inequalities we have $\lambda^2 \geq 0$, with $\lambda^2 = 0$ holding if and only if the momenta are collinear. Next, we define

$$X = \frac{p_1^2 - p_2^2 - p_3^2 + \sqrt{-\lambda^2}}{2p_2 p_3}, \qquad Y = \frac{p_2^2 - p_1^2 - p_3^2 + \sqrt{-\lambda^2}}{2p_1 p_3}, \tag{6.22}$$

$$Z = \frac{p_3^2 - p_1^2 - p_2^2 - \sqrt{-\lambda^2}}{2p_1 p_2}. \tag{6.23}$$

Note that X and Y are symmetric under $p_1 \leftrightarrow p_2$, while the sign in front of the square root in Z is different to that in X and Y. For physical momentum configurations these variables are complex, although the entire expression for a given triple-K integral remains real.

Starting with the representation (A.5.16) of the triple-K integral, one can use the reduction formulae (A.5.10) - (A.5.13) in order to find

$$\begin{aligned}I_{\nu+1\{\nu\nu\nu\}} = \frac{2^{\nu-2}\Gamma(\nu)\pi}{\sin(\pi\nu)} &\left[\frac{p_3^{2\nu}}{p_1 p_2} Z \cdot F_\nu(Z^2) + \frac{p_2^{2\nu}}{p_1 p_3} Y \cdot F_\nu\left(Z\frac{Y}{X}\right) + \frac{p_1^{2\nu}}{p_2 p_3} X \cdot F_\nu\left(Z\frac{X}{Y}\right)\right] \\ &+ \frac{2^{3\nu-2}\pi^{\frac{3}{2}}\Gamma\left(\nu+\frac{1}{2}\right)}{\sin^2(\pi\nu)} (p_1 p_2 p_3)^{2\nu} (\sqrt{-\lambda^2})^{-(2\nu+1)}.\end{aligned} \tag{6.24}$$

where

$$F_\nu(x) = {}_2F_1(1, \nu+1; 1-\nu; x) \tag{6.25}$$

and the X, Y, Z variables are defined in (6.22, 6.23) while λ^2 is given by (6.21). Note that this particular combination of parameters in the hypergeometric function appears in Legendre functions.

For generic values of ν the expression (6.24) is finite. In order to evaluate $I_{0\{111\}}$, however, we require $\nu = -1$ (see (6.20)) where (6.24) has singularities. In cases such as this, (6.24) may be series expanded in ν. The relevant expansion of the hypergeometric function here is

$$F_{-1+\epsilon}(x) = 1 + F^{(1)}_{-1}(x)\epsilon + F^{(2)}_{-1}(x)\epsilon^2 + O(\epsilon^3), \tag{6.26}$$

where

$$F^{(1)}_{-1}(x) = 1 - \left(1 - \frac{1}{x}\right)\log(1-x), \tag{6.27}$$

$$F^{(2)}_{-1}(x) = 2 + \left(1 - \frac{1}{x}\right)\left[-\log(1-x) + \log^2(1-x) + \mathrm{Li}_2\, x\right]. \tag{6.28}$$

Combining everything we obtain an analytic expression for the integral $I_{0+\epsilon\{-1+\epsilon,-1+\epsilon,-1+\epsilon\}}$. Note however that this result is given in a different regularisation scheme than our usual (5.7). We can easily change the regularisation scheme through a comparison of the local terms in the triple-K integrals in two regularisation schemes. This can be done without actually evaluating the integrals: we simply use (A.4.1, A.4.2) and/or (A.4.7) to expand out two of the three Bessel K functions in a given triple-K integral then use the formulae (A.5.21, A.5.23, A.5.25). In each case, we find only a finite number of terms leading to singularities.

To illustrate this, consider for example the integral $I_{2\{111\}}$. The calculations for $I_{0\{111\}}$ are essentially identical, however, due to the fact that $I_{0+\epsilon\{111\}}$ has a double pole in ϵ, the resulting expressions are much longer. The integrals in the two regularization schemes are given by

$$I_{2+\epsilon\{111\}} = \int_0^\infty \mathrm{d}x\, x^\epsilon \left[1 + O(x^2)\right] p_3 K_1(p_3 x) \tag{6.29}$$

$$I_{2+\epsilon\{1+\epsilon,1+\epsilon,1+\epsilon\}} = \int_0^\infty \mathrm{d}x\, x^{-\epsilon} \left[(4p_3)^\epsilon \Gamma^2(1+\epsilon) + O(x^2)\right] p_3^{1+\epsilon} K_{1+\epsilon}(p_3 x) \tag{6.30}$$

Using the expansion of the Bessel K functions we find that the two integrals differ by local terms:

$$I_{2+\epsilon\{111\}} = I_{2+\epsilon\{1+\epsilon,1+\epsilon,1+\epsilon\}} + \frac{3}{2\epsilon} + \frac{3}{2}(-\gamma_E + \log 2) + O(\epsilon). \tag{6.31}$$

We can obtain $I_{0+\epsilon\{-1-1-1\}}$ from $I_{0+\epsilon\{-1+\epsilon,-1+\epsilon,-1+\epsilon\}}$ in a similar way, and then use (6.20). The exact analytic expression for $I_{0\{111\}}$ is given in appendix A.6. The only special functions appearing in the result are dilogarithms.

7 Worked example: $\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3}\rangle$

Now that our general method is complete, in this section we present a full worked example, the $\langle\!\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3}\rangle\!\rangle$ correlation function. Here we will take J^μ to be a conserved $U(1)$ current; more general results are listed in Part II. This correlator provides a useful test case as, while more complex than the $\langle T^{\mu_1\nu_1} T^{\mu_2\nu_2}\mathcal{O}\rangle$ correlator we used to illustrate the method in earlier sections, it is nonetheless simpler than correlators with more stress-energy tensors.

We will also discuss the complete evaluation of all integrals in both $d = 3$ and $d = 4$ and present a concrete model, free fermions, where these correlators can be explicitly computed by standard Feynman diagrams. These results provide a nontrivial consistency check on our method.

7.1 Primary conformal Ward identities

We start with the analysis of primary CWIs for the $\langle\!\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3} \rangle\!\rangle$ correlation function in general Euclidean dimension d. For the decomposition of the transverse-traceless part of $\langle\!\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3} \rangle\!\rangle$ we follow the analysis of section 3.1. The decomposition consists of four form factors,

$$\begin{aligned}
&\langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1) j^{\mu_2}(\boldsymbol{p}_2) j^{\mu_3}(\boldsymbol{p}_3) \rangle\!\rangle \\
&\quad = \Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1) \pi^{\mu_2}_{\alpha_2}(\boldsymbol{p}_2) \pi^{\mu_3}_{\alpha_3}(\boldsymbol{p}_3) \Big[A_1 p_2^{\alpha_1} p_2^{\beta_1} p_3^{\alpha_2} p_1^{\alpha_3} + A_2 \delta^{\alpha_2\alpha_3} p_2^{\alpha_1} p_2^{\beta_1} \\
&\qquad + A_3 \delta^{\alpha_1\alpha_2} p_2^{\beta_1} p_1^{\alpha_3} + A_3(p_2 \leftrightarrow p_3) \delta^{\alpha_1\alpha_3} p_2^{\beta_1} p_3^{\alpha_2} \\
&\qquad + A_4 \delta^{\alpha_1\alpha_3} \delta^{\alpha_2\beta_1} \Big] .
\end{aligned} \tag{7.1}$$

Here, $p_2 \leftrightarrow p_3$ denotes exchange of the arguments p_2 and p_3, *i.e.*, $A_3(p_2 \leftrightarrow p_3) = A_3(p_1, p_3, p_2)$. If on the other hand no arguments are given then the standard ordering is assumed, *i.e.*, $A_3 = A_3(p_1, p_2, p_3)$. Note that the form factors A_1, A_2 and A_4 are symmetric under $p_2 \leftrightarrow p_3$,

$$A_j(p_1, p_3, p_2) = A_j(p_1, p_2, p_3), \qquad j \in \{1, 2, 4\}, \tag{7.2}$$

while the form factor A_3 does not exhibit any symmetry properties.

Next, the primary CWIs can be extracted by means of the procedure described in section 4.3. These CWIs are

$$\begin{array}{ll}
\mathrm{K}_{12}\, A_1 = 0, & \mathrm{K}_{13}\, A_1 = 0, \\
\mathrm{K}_{12}\, A_2 = -2A_1, & \mathrm{K}_{13}\, A_2 = -2A_1, \\
\mathrm{K}_{12}\, A_3 = 0, & \mathrm{K}_{13}\, A_3 = 4A_1, \\
\mathrm{K}_{12}\, A_4 = 2A_3, & \mathrm{K}_{13}\, A_4 = 2A_3(p_2 \leftrightarrow p_3).
\end{array} \tag{7.3}$$

The solution follows from the analysis of section 5.2,

$$A_1 = \alpha_1 J_{4\{000\}}, \tag{7.4}$$

$$A_2 = \alpha_1 J_{3\{100\}} + \alpha_2 J_{2\{000\}}, \tag{7.5}$$

$$A_3 = 2\alpha_1 J_{3\{001\}} + \alpha_3 J_{2\{000\}}, \tag{7.6}$$

$$A_4 = 2\alpha_1 J_{2\{011\}} + \alpha_3 \left(J_{1\{010\}} + J_{1\{001\}} \right) + \alpha_4 J_{0\{000\}}. \tag{7.7}$$

7.2 Evaluation of secondary conformal Ward identities

The independent secondary CWIs for $\langle\!\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3} \rangle\!\rangle$ are listed in the second part of the paper and read

$$\begin{aligned}
(*)\ & \mathrm{L}_{2,2}\, A_1 + \mathrm{R}_2 \left[A_3 - A_3(p_2 \leftrightarrow p_3) \right] \\
&\quad = 2d \cdot \text{coefficient of } p_2^{\mu_1} p_3^{\mu_2} p_1^{\mu_3} \text{ in } p_{1\nu_1} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2}(\boldsymbol{p}_2) J^{\mu_3}(\boldsymbol{p}_3) \rangle\!\rangle, \qquad (7.8) \\
& \mathrm{L}'_{1,2}\, A_1 + 2\, \mathrm{R}'_1 \left[A_3 - A_2 \right] \\
&\quad = 2d \cdot \text{coefficient of } p_2^{\mu_1} p_2^{\nu_1} p_1^{\mu_3} \text{ in } p_{2\mu_2} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2}(\boldsymbol{p}_2) J^{\mu_3}(\boldsymbol{p}_3) \rangle\!\rangle, \qquad (7.9) \\
& \mathrm{L}_{2,0}\, A_2 - p_1^2 \left[A_3 - A_3(p_2 \leftrightarrow p_3) \right] \\
&\quad = 2d \cdot \text{coefficient of } \delta^{\mu_2\mu_3} p_2^{\mu_1} \text{ in } p_{1\nu_1} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2}(\boldsymbol{p}_2) J^{\mu_3}(\boldsymbol{p}_3) \rangle\!\rangle, \qquad (7.10) \\
& \mathrm{L}_{2,2}\, A_3 - 2\, \mathrm{R}_2\, A_4 \\
&\quad = 4d \cdot \text{coefficient of } \delta^{\mu_1\mu_2} p_1^{\mu_3} \text{ in } p_{1\nu_1} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2}(\boldsymbol{p}_2) J^{\mu_3}(\boldsymbol{p}_3) \rangle\!\rangle, \qquad (7.11)
\end{aligned}$$

where L and R operators are defined in (4.31) and (4.32). They can be obtained by the procedure outlined in section 4.3.2. Note that there are four primary constants and four secondary CWIs. As some of the secondary CWIs are trivially satisfied, however, not all four of the primary

constants are fixed, as we expect from the position space analysis [3]. Secondary CWIs that are trivially satisfied are denoted by asterisks in the second part of the paper (for example (7.8) above is of this type).

Before solving the secondary CWIs, we must simplify the semi-local terms appearing on their right-hand sides. Differentiating (4.48, 4.49, 4.61) we find the following transverse and trace Ward identities,

$$\begin{aligned}
&p_1^{\nu_1}\langle\!\langle T_{\mu_1\nu_1}(\boldsymbol{p}_1)J^{\mu_2}(\boldsymbol{p}_2)J^{\mu_3}(\boldsymbol{p}_3)\rangle\!\rangle = \\
&\quad = p_1^{\nu_1}\langle\!\langle \frac{\delta T_{\mu_1\nu_1}}{\delta A_{\mu_3}}(\boldsymbol{p}_1,\boldsymbol{p}_3)J^{\mu_2}(\boldsymbol{p}_2)\rangle\!\rangle + p_1^{\nu_1}\langle\!\langle \frac{\delta T_{\mu_1\nu_1}}{\delta A_{\mu_2}}(\boldsymbol{p}_1,\boldsymbol{p}_2)J^{\mu_3}(\boldsymbol{p}_3)\rangle\!\rangle \\
&\qquad - p_3^{\mu_1}\langle\!\langle J^{\mu_2}(-\boldsymbol{p}_2)J^{\mu_3}(-\boldsymbol{p}_2)\rangle\!\rangle - p_2^{\mu_1}\langle\!\langle J^{\mu_2}(\boldsymbol{p}_3)J^{\mu_3}(-\boldsymbol{p}_3)\rangle\!\rangle \\
&\qquad + \delta^{\mu_3}_{\mu_1}p_{3\alpha}\langle\!\langle J^{\mu_2}(\boldsymbol{p}_2)J^{\alpha}(-\boldsymbol{p}_2)\rangle\!\rangle + \delta^{\mu_2}_{\mu_1}p_{2\alpha}\langle\!\langle J^{\alpha}(\boldsymbol{p}_3)J^{\mu_3}(-\boldsymbol{p}_3)\rangle\!\rangle, \qquad (7.12)\\
&p_{2\mu_2}\langle\!\langle T_{\mu_1\nu_1}(\boldsymbol{p}_1)J^{\mu_2}(\boldsymbol{p}_2)J^{\mu_3}(\boldsymbol{p}_3)\rangle\!\rangle = \\
&\quad = 2p_{2\mu_2}\langle\!\langle \frac{\delta J^{\mu_2}}{\delta g^{\mu_1\nu_1}}(\boldsymbol{p}_2,\boldsymbol{p}_1)J^{\mu_3}(\boldsymbol{p}_3)\rangle\!\rangle + \delta_{\mu_1\nu_1}p_{1\alpha}\langle\!\langle J^{\alpha}(\boldsymbol{p}_3)J^{\mu_3}(-\boldsymbol{p}_3)\rangle\!\rangle, \qquad (7.13)\\
&\langle\!\langle T(\boldsymbol{p}_1)J^{\mu_2}(\boldsymbol{p}_2)J^{\mu_3}(\boldsymbol{p}_3)\rangle\!\rangle = \langle\!\langle \frac{\delta T}{\delta A_{\mu_2}}(\boldsymbol{p}_1,\boldsymbol{p}_2)J^{\mu_3}(\boldsymbol{p}_3)\rangle\!\rangle + \langle\!\langle \frac{\delta T}{\delta A_{\mu_3}}(\boldsymbol{p}_1,\boldsymbol{p}_3)J^{\mu_2}(\boldsymbol{p}_2)\rangle\!\rangle. \qquad (7.14)
\end{aligned}$$

In the next section we will extract algebraic equations between the primary constants by taking the zero-momentum limit $p_3 \to 0$. The details of this procedure are described in section 5.3. We will find that in the zero-momentum limit the right-hand sides of the secondary CWIs (7.8) - (7.10) are given by

$$\begin{aligned}
&\lim_{\substack{p_3\to 0\\ p_1=p_2=p}} \text{coefficient of } p_2^{\mu_1}p_2^{\nu_1}p_1^{\mu_3} \text{ in } p_{2\mu_2}\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)J^{\mu_2}(\boldsymbol{p}_2)J^{\mu_3}(\boldsymbol{p}_3)\rangle\!\rangle = 0, \qquad (7.15)\\
&\lim_{\substack{p_3\to 0\\ p_1=p_2=p}} \text{coefficient of } p_2^{\mu_1}p_3^{\mu_2}p_1^{\mu_3} \text{ in } p_{1\nu_1}\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)J^{\mu_2}(\boldsymbol{p}_2)J^{\mu_3}(\boldsymbol{p}_3)\rangle\!\rangle = 0, \qquad (7.16)\\
&\lim_{\substack{p_3\to 0\\ p_1=p_2=p}} \text{coefficient of } \delta^{\mu_2\mu_3}p_2^{\mu_1} \text{ in } p_{1\nu_1}\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)J^{\mu_2}(\boldsymbol{p}_2)J^{\mu_3}(\boldsymbol{p}_3)\rangle\!\rangle = \\
&\qquad\qquad = \text{coefficient of } \delta^{\mu_2\mu_3} \text{ in } \langle\!\langle J^{\mu_2}(\boldsymbol{p})J^{\mu_3}(-\boldsymbol{p})\rangle\!\rangle. \qquad (7.17)
\end{aligned}$$

Let us start with the first result (7.15). Due to conformal invariance, the only operators in $\delta J^{\mu_2 a}/\delta g^{\mu_1\nu_1}$ that can give a non-vanishing result under the expectation value with the current is another current $J^{\mu b}$. In general, the descendants of the current can also give a non-vanishing 2-point function with another current. In this case, however, the dilatation degree of $\delta J^{\mu_2 a}/\delta g^{\mu_1\nu_1}$ is $d-1$, and so descendants cannot appear. The most general form of the functional derivative term is therefore

$$\frac{\delta J^{\mu_2}}{\delta g^{\mu_1\nu_1}} = c_1\delta_{\mu_1\nu_1}J^{\mu_2} + c_2\delta^{\mu_2}_{(\mu_1}J_{\nu_1)} + \dots \qquad (7.18)$$

where c_1 and c_2 are numerical constants and the omitted terms may contain operators from different conformal families to that of $J^{\mu b}$. The 2-point function then reads

$$\langle\!\langle \frac{\delta J^{\mu_2}}{\delta g^{\mu_1\nu_1}}(\boldsymbol{p}_2,\boldsymbol{p}_1)J^{\mu_3}(\boldsymbol{p}_3)\rangle\!\rangle = \Big[c_1\delta_{\mu_1\nu_1}\delta^{\mu_2}_{\alpha} + c_2\delta^{\mu_2}_{(\mu_1}\delta_{\nu_1)\alpha}\Big]\,\langle\!\langle J^{\alpha}(\boldsymbol{p}_3)J^{\mu_3}(-\boldsymbol{p}_3)\rangle\!\rangle. \qquad (7.19)$$

In the limit $p_3 \to 0$, however, the 2-point function vanishes, since it behaves as p_3^{d-2} and $d > 2$. The same argument works for the second term in (7.13) and so (7.15) also vanishes.

Let us now establish the remaining formulae (7.16) and (7.17). Following the same argument for the limit $p_3 \to 0$, we can restrict consideration to the following terms in (7.12)

$$p_1^{\nu_1}\langle\!\langle \frac{\delta T_{\mu_1\nu_1}}{\delta A_{\mu_3}}(\boldsymbol{p}_1,\boldsymbol{p}_3)J^{\mu_2}(\boldsymbol{p}_2)\rangle\!\rangle - p_3^{\mu_1}\langle\!\langle J^{\mu_2}(-\boldsymbol{p}_2)J^{\mu_3}(-\boldsymbol{p}_2)\rangle\!\rangle + \delta^{\mu_3}_{\mu_1}p_{3\alpha}\langle\!\langle J^{\mu_2}(\boldsymbol{p}_2)J^{\alpha}(-\boldsymbol{p}_2)\rangle\!\rangle. \qquad (7.20)$$

Using the representation (5.65) it is straightforward to expand the last two terms. As usual, we must use the convention (3.7) for the momenta associated with Lorentz indices, leading to the right-hand sides of (7.16) and (7.17). The remaining task is then to show that there are no contributions from the first term with the functional derivative.

Since the dimension of the stress-energy tensor is d, while that of the conserved current is $d-1$ and that of the source A_μ is 1, the only possible contributions to the first term in (7.12) are

$$T_{\mu\nu} = c_3 \left[A_\mu J_\nu + A_\nu J_\mu\right] + \dots \tag{7.21}$$

where c_3 is a numerical constant and the omitted terms do not contain the current or its descendants. This definition of c_3 applies if the J^μ operator is the unique spin-1 conserved current in theory. If not, we can instead define the constant c_3 through the 2-point function

$$\langle\!\langle \frac{\delta T_{\mu_1\nu_1}}{\delta A_{\mu_2}}(\boldsymbol{p}_1,\boldsymbol{p}_2) J^{\mu_3}(\boldsymbol{p}_3)\rangle\!\rangle = 2c_3 \delta^{\mu_2}_{(\mu_1} \langle\!\langle J_{\nu_1)}(\boldsymbol{p}_3) J^{\mu_3}(-\boldsymbol{p}_3)\rangle\!\rangle. \tag{7.22}$$

After taking the functional derivative one finds that tensors $p_2^{\mu_1} p_3^{\mu_2} p_1^{\mu_3}$ and $\delta^{\mu_2\mu_3}$ are absent in (7.12).

Finally, with the definition of the c_3 constant as in (7.22), the same method can be applied to work out the zero-momentum limit of the right-hand side of the final secondary CWI (7.11), yielding the result

$$\begin{aligned}\lim_{\substack{p_3\to 0\\ p_1=p_2=p}} \text{ coefficient of } \delta^{\mu_1\mu_2} p_1^{\mu_3} \text{ in } p_{1\nu_1} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2}(\boldsymbol{p}_2) J^{\mu_3}(\boldsymbol{p}_3)\rangle\!\rangle = \\ = c_3 \cdot \text{coefficient of } \delta^{\mu_1\mu_2} \text{ in } \langle\!\langle J^{\mu_1}(\boldsymbol{p}) J^{\mu_2}(-\boldsymbol{p})\rangle\!\rangle.\end{aligned} \tag{7.23}$$

7.3 Solutions to secondary conformal Ward identities

Our goal now is to analyse the additional constraints imposed by the secondary CWIs (7.8) - (7.11) on the solution (7.7) of the primary CWIs. We proceed as in sections 5.3.2 and 5.3.3. First, we use solutions (7.7) and take the zero-momentum limit $p_3 \to 0$. From (7.9) and (7.10), we then derive the following equations for the primary constants

$$\alpha_2 = \alpha_3, \tag{7.24}$$

$$d\alpha_1 + \alpha_2 = \frac{2^{3-\frac{d}{2}} s_d c_J}{\Gamma\left(\frac{d}{2}\right)}, \tag{7.25}$$

where

$$s_d = \begin{cases} \frac{1}{\pi}(-1)^{\frac{d-1}{2}} & \text{if } d = 3,5,7,\dots, \\ (-1)^{\frac{d}{2}-1} & \text{if } d = 4,6,8,\dots \end{cases} \tag{7.26}$$

and c_J encodes the normalisation of the 2-point function as given in (5.65). For the right-hand sides we used (7.15) - (7.17)

The situation is more interesting for the last of the secondary CWIs (7.11). Assuming d to be odd, we find this equation exhibits a singularity and the $\epsilon \to 0$ limit cannot be taken. Expanding in powers of ϵ, the zero-momentum limit of the left-hand side is

$$\left(\frac{2}{\epsilon} - \log p^2\right) \cdot \frac{2^{\frac{d}{2}-3} d p^{d-2} \pi}{\sin\left(\frac{d\pi}{2}\right) \Gamma\left(\frac{d}{2}-1\right)} \left[\alpha_4 + (-2+d)\alpha_3\right] + \dots \tag{7.27}$$

where the omitted terms are of order ϵ^0. Since the right-hand side is finite and does not contain logarithms, (7.11) forces

$$\alpha_4 = (2-d)\alpha_3 + O(\epsilon). \tag{7.28}$$

With this value of α_4, the left-hand side of (7.11) is finite, but does not necessarily match the right-hand side. To solve this problem we must consider a first-order correction to α_4, *i.e.*, we write

$$\alpha_4 = (2-d)\alpha_3 + \epsilon\alpha_4^{(1)}. \tag{7.29}$$

Using (7.23), the equation for $\alpha_4^{(1)}$ leads to

$$2d(d-2)\alpha_1 + d\alpha_3 + 2\alpha_4^{(1)} = \frac{2^{5-\frac{d}{2}} s_d c_3 c_J}{\Gamma\left(\frac{d}{2}-1\right)}. \tag{7.30}$$

This equation is valid for even d as well: in this case we find that the left-hand side of (7.11) contains a $1/\epsilon^2$ singularity. As there is no corresponding singularity on the right-hand side we recover the same conditions (7.28) and (7.30) as above.

Summarising, we found that the primary constants in the solution to the primary CWIs (7.7) satisfy

$$\begin{aligned} \alpha_3 &= \alpha_2, \\ d\alpha_1 + \alpha_2 &= \frac{2^{3-\frac{d}{2}} s_d c_J}{\Gamma\left(\frac{d}{2}\right)}, \\ \alpha_4 &= -(d-2)\alpha_2, \\ 2d(d-2)\alpha_1 + d\alpha_2 + 2\alpha_4^{(1)} &= \frac{2^{5-\frac{d}{2}} s_d c_3 c_J}{\Gamma\left(\frac{d}{2}-1\right)}. \end{aligned} \tag{7.31}$$

Through an analysis similar to that of section 5.3.6, we find there are no further constraints on the primary constants.

Our solution of the primary and secondary CWIs above depends on one undetermined primary constant as well as two different 2-point function normalisations. This result is in fact consistent with the position space result of [3] (which involves only a single 2-point function normalisation) by virtue of our different definition for the 3-point function, namely

$$\begin{aligned} \langle T_{\mu_1\nu_1}(\boldsymbol{x}) J^{\mu_2}(\boldsymbol{y}) J^{\mu_3}(\boldsymbol{z})\rangle = {} & \frac{-1}{\sqrt{g(\boldsymbol{z})}}\frac{\delta}{\delta A_{\mu_3}(\boldsymbol{z})}\frac{-1}{\sqrt{g(\boldsymbol{y})}}\frac{\delta}{\delta A_{\mu_2}(\boldsymbol{y})}\frac{-2}{\sqrt{g(\boldsymbol{x})}}\frac{\delta}{\delta g^{\mu_1\nu_1}(\boldsymbol{x})} Z[g^{\mu\nu}, A_\rho] \\ & + \langle \frac{\delta T_{\mu_1\nu_1}(\boldsymbol{x})}{\delta A_{\mu_2}(\boldsymbol{y})} J^{\mu_3}(\boldsymbol{z})\rangle + \langle \frac{\delta T_{\mu_1\nu_1}(\boldsymbol{x})}{\delta A_{\mu_3}(\boldsymbol{z})} J^{\mu_2}(\boldsymbol{y})\rangle. \end{aligned} \tag{7.32}$$

In [3] (and similarly [26, 28]) the semi-local terms on the right-hand side of this formula are absorbed into the definition of the $\langle T_{\mu_1\nu_1} J^{\mu_2} J^{\mu_3}\rangle$ correlator: it is these semi-local terms that are responsible, via (7.22), for the dependence of our solution on the additional normalisation constant c_3.

7.4 General form of $\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3}\rangle$ in $d = 3$

Let us now focus on the special case of $d = 3$. Examining the form of the solution (7.7) to the primary CWIs, we find that all triple-K integrals can be evaluated in terms of elementary integrals using (A.4.4). If an integral diverges, we use the regularisation (5.7). In this way, we

find

$$J_{4\{000\}} = I_{\frac{9}{2}\{\frac{3}{2}\frac{1}{2}\frac{1}{2}\}} = 2\left(\frac{\pi}{2}\right)^{\frac{3}{2}} \frac{4p_1+p_2+p_3}{(p_1+p_2+p_3)^4}, \tag{7.33}$$

$$J_{3\{100\}} = I_{\frac{7}{2}\{\frac{5}{2}\frac{1}{2}\frac{1}{2}\}} = \left(\frac{\pi}{2}\right)^{\frac{3}{2}} \frac{9(p_1p_2+p_1p_3)+6p_2p_3+8p_1^2+3(p_2^2+p_3^2)}{(p_1+p_2+p_3)^3}, \tag{7.34}$$

$$J_{2\{000\}} = I_{\frac{5}{2}\{\frac{3}{2}\frac{1}{2}\frac{1}{2}\}} = \left(\frac{\pi}{2}\right)^{\frac{3}{2}} \frac{2p_1+p_2+p_3}{(p_1+p_2+p_3)^2}, \tag{7.35}$$

$$\begin{aligned} J_{2\{011\}} = I_{\frac{5}{2}\{\frac{3}{2}\frac{3}{2}\frac{3}{2}\}} = &-\left(\frac{\pi}{2}\right)^{\frac{3}{2}} \frac{1}{(p_1+p_2+p_3)^2}\left[2p_1p_2p_3+p_1^3+p_2^3+p_3^3\right. \\ &\left.+2(p_1^2p_2+p_1p_2^2+p_1p_3^2+p_3p_1^2+p_2p_3^2+p_3p_2^2)\right], \end{aligned} \tag{7.36}$$

$$\begin{aligned} J_{1+\epsilon\{010\}} = I_{\frac{3}{2}+\epsilon\{\frac{3}{2}\frac{3}{2}\frac{1}{2}\}} = &\left(\frac{\pi}{2}\right)^{\frac{3}{2}}\left[-\frac{p_3}{\epsilon}+p_3\log(p_1+p_2+p_3)\right. \\ &\left.+\frac{-p_1p_2+(\gamma_E-2)(p_1p_3+p_2p_3)-p_1^2-p_2^2+(\gamma_E-1)p_3^2}{p_1+p_2+p_3}+O(\epsilon)\right], \end{aligned} \tag{7.37}$$

$$\begin{aligned} J_{0+\epsilon\{000\}} = I_{\frac{1}{2}+\epsilon\{\frac{3}{2}\frac{1}{2}\frac{1}{2}\}} = &\left(\frac{\pi}{2}\right)^{\frac{3}{2}}\left[-\frac{p_2+p_3}{\epsilon}+(p_2+p_3)\log(p_1+p_2+p_3)\right. \\ &\left.+(\gamma_E-1)(p_2+p_3)-p_1+O(\epsilon)\right], \end{aligned} \tag{7.38}$$

with similar integrals following from the permutation formula (5.11).

Applying the secondary CWIs (7.31) we then obtain the final result

$$A_1 = \alpha_1 \frac{2(4p_1+p_2+p_3)}{(p_1+p_2+p_3)^4}, \tag{7.39}$$

$$A_2 = \frac{2\alpha_1 p_1^2}{(p_1+p_2+p_3)^3} - \frac{2(2p_1+p_2+p_3)}{(p_1+p_2+p_3)^2}c_J, \tag{7.40}$$

$$A_3 = \frac{\alpha_1}{(p_1+p_2+p_3)^3}\left[-2p_1^2-p_2^2+p_3^2-3p_1p_2+3p_1p_3\right] - \frac{2(2p_1+p_2+p_3)}{(p_1+p_2+p_3)^2}c_J, \tag{7.41}$$

$$\begin{aligned} A_4 = &\alpha_1 \frac{(p_1+p_2-p_3)(p_1-p_2+p_3)(2p_1+p_2+p_3)}{2(p_1+p_2+p_3)^2} \\ &+\left(\frac{2p_1^2}{p_1+p_2+p_3}-p_2-p_3\right)c_J + 2(p_2+p_3)c_3c_J. \end{aligned} \tag{7.42}$$

In these results we rescaled the coefficient α_1 according to $\alpha_1(\pi/2)^{3/2} \mapsto \alpha_1$, so as to remove the awkward factor of $(\pi/2)^{3/2}$.

The form factors build the transverse-traceless part of the correlation function according to (7.1). The full correlation function can then be recovered by means of (3.20) and (3.21). Using the transverse and trace Ward identities (7.12, 7.13, 7.14), we find

$$\begin{aligned} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)J^{\mu_2}(\boldsymbol{p}_2)J^{\mu_3}(\boldsymbol{p}_3)\rangle\!\rangle &= \langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1)j^{\mu_2}(\boldsymbol{p}_2)j^{\mu_3}(\boldsymbol{p}_3)\rangle\!\rangle \\ &+\left[2\mathscr{T}_\alpha^{\mu_1\nu_1}(\boldsymbol{p}_1)\pi^{\mu_3[\alpha}(\boldsymbol{p}_3)p_3^{\beta]}+\frac{p_3^{\mu_3}}{p_3^2}\delta^{\mu_1\nu_1}p_{1\beta}\right]\langle\!\langle J^{\mu_2}(\boldsymbol{p}_2)J_\beta(-\boldsymbol{p}_2)\rangle\!\rangle \\ &+\pi_{\alpha_3}^{\mu_3}(\boldsymbol{p}_3)\left[\mathscr{T}^{\mu_1\nu_1\alpha_1}(\boldsymbol{p}_1)p_1^{\beta_1}+\frac{\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)}{d-1}\delta^{\alpha_1\beta_1}\right]\langle\!\langle\frac{\delta T_{\alpha_1\beta_1}}{\delta A_{\alpha_3}}(\boldsymbol{p}_1,\boldsymbol{p}_3)J^{\mu_2}(\boldsymbol{p}_2)\rangle\!\rangle \\ &+\frac{2p_3^{\mu_3}p_{3\alpha_3}}{p_3^2}\delta^{\mu_1\alpha_1}\delta^{\nu_1\beta_1}\langle\!\langle\frac{\delta J^{\alpha_3}}{\delta g^{\alpha_1\beta_1}}(\boldsymbol{p}_3,\boldsymbol{p}_1)J^{\mu_2}(\boldsymbol{p}_2)\rangle\!\rangle \\ &+\text{ everything with }(\boldsymbol{p}_2,\mu_2)\leftrightarrow(\boldsymbol{p}_3,\mu_3), \end{aligned} \tag{7.43}$$

where $\mathscr{T}_\alpha^{\mu\nu}$ was given in (3.22). Here we assume no scale anomalies are present: if anomalies occur, the additional ultralocal contribution (8.25) should be added to (7.43), as we will discuss in section 8.

The result (7.43) is the most general explicit expression for the $\langle\!\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3} \rangle\!\rangle$ correlation function in the momentum space. As we can see, it depends on one undetermined primary constant plus the normalisations of the 2-point functions.

7.5 Free fermions in $d = 3$

As a cross-check on our calculations we now consider free fermions in $d = 3$ Euclidean dimensions given by the action

$$S = \int \mathrm{d}^3\boldsymbol{x}\, e \left[\bar{\psi} e^\mu_a \gamma^a \overset{\leftrightarrow}{D}_\mu \psi\right], \tag{7.44}$$

where

$$D_\mu = \nabla_\mu - \mathrm{i}A_\mu, \qquad \nabla_\mu = \partial_\mu - \frac{\mathrm{i}}{2}\omega^{ab}_\mu \Sigma_{ab}, \tag{7.45}$$

and ω^{ab}_μ is the spin connection

$$\omega^{ab}_\mu = e^a_\nu \partial_\mu e^{\nu b} + e^a_\nu e^{\sigma b} \Gamma^\nu{}_{\sigma\mu}, \qquad \Sigma^{ab} = \frac{\mathrm{i}}{4}[\gamma^a, \gamma^b]. \tag{7.46}$$

Here $\Gamma^\nu{}_{\sigma\mu}$ is the Christoffel symbol associated with the metric $g_{\mu\nu}$, while e^a_μ are vielbeins satisfying $e^a_\mu e_{\nu a} = g_{\mu\nu}$ and the gamma matrices γ^a satisfy $\gamma^\mu = e^\mu_a \gamma^a$. On flat space, we then have $\{\gamma^a, \gamma^b\} = -2\delta^{ab}$. In $d = 3$, the spin-$\frac{1}{2}$ representation of the group $SO(3)$ is 2-dimensional and $\mathrm{Tr}(\gamma^a\gamma^b) = -2\delta^{ab}$.

Notice that the gauge field A_μ is treated as a source for the conserved current and is not a degree of freedom. The stress-energy tensor and the conserved current in the presence of the sources are

$$T_{\mu\nu} = \frac{2}{\sqrt{g}}\frac{\delta S}{\delta g^{\mu\nu}} = \bar{\psi}\gamma_{(\mu} \overset{\leftrightarrow}{D}_{\nu)} \psi - g_{\mu\nu}\bar{\psi}\gamma^\alpha \overset{\leftrightarrow}{D}_\alpha \psi, \tag{7.47}$$

$$J^\mu = \frac{1}{\sqrt{g}}\frac{\delta S}{\delta A^\mu} = -\mathrm{i}\bar{\psi}\gamma^\mu\psi. \tag{7.48}$$

In this case the current is associated with the $U(1)$ symmetry, therefore we omit the group indices on J^μ. By direct calculation we find

$$\langle\!\langle J^\mu(\boldsymbol{p}) J^\nu(-\boldsymbol{p}) \rangle\!\rangle = -\frac{1}{16} p \pi^{\mu\nu}(\boldsymbol{p}), \tag{7.49}$$

$$\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}) T^{\mu_2\nu_2}(-\boldsymbol{p}) \rangle\!\rangle = \frac{1}{128} p^3 \Pi^{\mu_1\nu_1\mu_2\nu_2}(\boldsymbol{p}). \tag{7.50}$$

The transverse Ward identities can be obtained by differentiation of the equations (4.48, 4.49) and are listed in the second part of the paper. Some terms of the terms involve functional derivatives and may be evaluated directly from expressions (7.47, 7.48),

$$\frac{\delta T_{\mu\nu}(\boldsymbol{x})}{\delta A_\rho(\boldsymbol{y})} = \frac{1}{2}[J_\mu \delta^\rho_\nu + J_\nu \delta^\rho_\mu - 2J^\rho \delta_{\mu\nu}]\delta(\boldsymbol{x} - \boldsymbol{y}), \tag{7.51}$$

$$\frac{\delta J^\mu(\boldsymbol{x})}{\delta g^{\alpha\beta}(\boldsymbol{y})} = \frac{1}{4}[J_\beta \delta^\mu_\alpha + J_\alpha \delta^\mu_\beta]\delta(\boldsymbol{x} - \boldsymbol{y}), \tag{7.52}$$

where the sources are turned off after the derivative is taken. All together, for this particular CFT we find

$$c_J = -\frac{1}{16}, \qquad c_T = \frac{1}{128}, \qquad c_3 = \frac{1}{2}. \tag{7.53}$$

where the 2-point function normalisations c_J and c_T, and the constant c_3, are as defined in (5.65), (5.66) and (7.22) respectively.

The 3-point function can be calculated by the usual Feynman rules. Using the results of section 3.3, one finds

$$A_1 = -\frac{4p_1 + p_2 + p_3}{12\,(p_1+p_2+p_3)^4}, \tag{7.54}$$

$$A_2 = \frac{9(p_1p_2 + p_1p_3) + 6p_2p_3 + 4p_1^2 + 3(p_2^2 + p_3^2)}{24\,(p_1+p_2+p_3)^3}, \tag{7.55}$$

$$A_3 = \frac{6p_1p_2 + 3p_1p_3 + 3p_2p_3 + 4p_1^2 + 2p_2^2 + p_3^2}{12\,(p_1+p_2+p_3)^3}, \tag{7.56}$$

$$A_4 = -\frac{4p_1p_2p_3 + 7(p_1^2p_2 + p_1^2p_3) - 2(p_1p_2^2 + p_1p_3^2) + p_2p_3^2 + p_3p_2^2 + 8p_1^3 - (p_2^3 + p_3^3)}{48\,(p_1+p_2+p_3)^2}. \tag{7.57}$$

The form factors A_j are defined in the decomposition (7.1).

We can compare this result directly with the solution (7.39) - (7.42). Since we know the 2-point function normalisations (7.53) there is only one undetermined constant, α_1. The solution (7.54) - (7.57) then fits perfectly with $\alpha_1 = -\frac{1}{24}$. In fact, the secondary Ward identities provide quite a robust check on the standard QFT calculation of the 3-point function: for example, a mistake leading to the overall rescaling of all form factors in (7.54) - (7.57) by some factor would immediately lead to an inconsistency with the 2-point function normalisation constants (7.53).

7.6 General form of $\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3} \rangle$ in $d = 4$

Using the reduction scheme of section 6, we can write down the most general form of $\langle\!\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3} \rangle\!\rangle$ in $d = 4$. Starting from the solutions (7.7) and (7.31) for the primary and secondary CWIs, using the regularisation scheme (5.7) and the relations in table 1, page 44, after removing divergences we find

$$A_1 = \alpha_1 I_{5\{211\}}, \tag{7.58}$$

$$A_2 = -\left(2c_J + \alpha_1 p_1 \frac{\partial}{\partial p_1}\right) I^{(0)}_{3\{211\}}, \tag{7.59}$$

$$A_3 = -2\left(c_J + \alpha_1 p_3 \frac{\partial}{\partial p_3}\right) I^{(0)}_{3\{211\}}, \tag{7.60}$$

$$A_4 = 2c_J\left[-2 + p_2\frac{\partial}{\partial p_2} + p_3\frac{\partial}{\partial p_3}\right] I^{(0)}_{1\{211\}} + 2\alpha_1 p_2 p_3 \frac{\partial^2}{\partial p_2 \partial p_3} I^{(0)}_{1\{211\}} \tag{7.61}$$

$$+ 4(c_J - c_3 c_J)\left[p_2^2 \log p_2 + p_3^2 \log p_3 - (p_2^2 + p_3^2)\left(\tfrac{1}{2} - \gamma_E + \log 2\right) - \tfrac{1}{2}p_1^2\right], \tag{7.62}$$

where $I^{(0)}_{\alpha\{\beta_j\}}$ denotes the the coefficient of ϵ^0 in the series expansion of the regulated integral $I_{\alpha+\epsilon\{\beta_j\}}$ in ϵ. This is the exact, finite and fully renormalised result for the transverse-traceless part of the correlation function. The triple-K integrals appearing can be straightforwardly evaluated using the reduction scheme in table 1 on page 44 starting from the master integral $I_{0\{111\}}$ given in appendix A.6. (We have nonetheless retained the above form for its compactness.) The transverse-traceless part of the correlation function can be recovered using (7.1), as in the case of $d = 3$. The full 3-point function can then be reconstructed by means of (7.43). For $d = 4$ an anomalous contribution appears, however, due to the addition of counterterms required to render the 3-point function finite that are not of the form assumed in section 4.4. We will return to the treatment of anomalies shortly, in section 8.

7.7 Free fermions in $d = 4$

The momentum space $\langle\!\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3} \rangle\!\rangle$ correlation function for free fermions in $d = 4$ dimensions was discussed in [26, 28]. In this section we will show how to simplify these calculations considerably using our method. We already know that the solution to the primary CWIs is given

by (7.7). We therefore need to calculate explicitly only one primary constant, say α_1, since the remaining constants are determined by the secondary CWIs (7.31).

To evaluate α_1 we can use the standard Feynman parametrisation, which gives

$$\text{coefficient of } p_2^{\mu_1} p_2^{\nu_1} p_3^{\mu_2} p_1^{\mu_3} \text{ in } \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2}(\boldsymbol{p}_2) J^{\mu_3}(\boldsymbol{p}_1)\rangle\!\rangle = -\frac{2}{\pi^2}\int_{[0,1]^3} \mathrm{d}X\, \frac{x_1 x_2 x_3^2}{D}, \tag{7.63}$$

where

$$\mathrm{d}X = \mathrm{d}x_1 \mathrm{d}x_2 \mathrm{d}x_3\, \delta(x_1 + x_2 + x_3 - 1), \tag{7.64}$$

$$D = p_1^2 x_2 x_3 + p_2^2 x_1 x_3 + p_3^2 x_1 x_2. \tag{7.65}$$

On the other hand, using (A.3.23), we find

$$J_{4\{000\}} = I_{5\{211\}} = 96 \cdot \int_{[0,1]^3} \mathrm{d}X\, \frac{x_1 x_2 x_3^2}{D} \tag{7.66}$$

and hence

$$\alpha_1 = -\frac{1}{48\pi^2}. \tag{7.67}$$

The 2-point function normalisation c_J and the constant c_3 are

$$c_J = -\frac{1}{12\pi^2}, \qquad c_3 = \frac{1}{2}, \tag{7.68}$$

and so, from (7.31), we find

$$\alpha_2 = \alpha_3 = \frac{1}{4\pi^2}, \qquad \alpha_4 = -\frac{1}{2\pi^2} - \frac{\epsilon}{6\pi^2}. \tag{7.69}$$

With these primary constants the expressions (7.7) represent a complete and concise solution to the transverse-traceless part of the $\langle\!\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3}\rangle\!\rangle$ correlation function for free fermions in $d = 4$. The full 3-point function can be recovered as in the $d = 3$ case via (7.1) and (7.43).

The above solution can be confirmed by direct calculations. For free field theory we computed the entire $\langle\!\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3}\rangle\!\rangle$ correlator using Passarino-Veltman reduction [50]. The coefficients of the appropriate tensors were then extracted according to section 3.3 and the result compared with the Feynman parametrised integrals. Exact agreement was found, both for the functional form of (7.7) and for the constants in (7.69).

Our result can also be compared with those of [28]. Up to a multiplicative factor, we find that

$$\begin{aligned}\text{coefficient of } p_2^{\mu_1} p_2^{\nu_1} p_3^{\mu_2} p_1^{\mu_3} &\sim \int_{[0,1]^3} \mathrm{d}X \frac{c_9(x_1,x_2) - 2c_8(x_1,x_2) + c_7(x_1,x_2)}{D} \\ &= \int_{[0,1]^3} \mathrm{d}X \frac{4x_1 x_2 x_3^2}{D},\end{aligned} \tag{7.70}$$

where the c_j polynomials are defined in the table 3 of [28]. This expression agrees with (7.66) and indeed represents the form factor A_1.

8 Divergences and anomalies in $d = 4$

We discussed in previous sections the solution of the conformal Ward identities and we have seen that in certain cases the triple-K integrals diverge and need to be regularised and renormalised. These infinities should be removed by means of local covariant counterterms. It turns out, however, that in some cases the counterterms break some of the symmetries and this leads to

anomalies. We have already encountered this issue when discussing the 3-point function of scalar operators in section 2. There, we saw the triple-K integral corresponding to the 3-point function of a dimension two operator in $d = 3$ diverges and the infinity can be removed by adding a local counterterm that is cubic in the source of the operator. The counterterm however is not scale invariant and this implies a trace anomaly. The anomaly then implies that the 4-point function of the stress-energy tensor with three scalar operators contains an ultra-local term that is not scheme-dependent because it is fixed unambiguously by the the anomalous Ward identity. Recall also that finite local counterterms are related to/parametrise scheme-dependence. In this case the same counterterm but with a finite coefficient is related to scheme-dependence.

In this section we would like to like to extend this discussion to correlators of the stress-energy tensor and of symmetry currents. For concreteness, we will focus on the case $d = 4$ but the discussion generalises to all dimensions and/or other operators. In particular, we will analyse the $\langle\!\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3} \rangle\!\rangle$ and $\langle\!\langle T^{\mu_1\nu_1} T^{\mu_2\nu_2} T^{\mu_3\nu_3} \rangle\!\rangle$ correlators. We will show that the divergences in the solutions for the transverse-traceless parts are cancelled by counterterms. These counterterms break scale invariance and lead to trace anomalies. The anomalies that originate from infinities in 2-point functions lead to scheme-independent ultra-local terms in 3-point functions, while those originating in 3-point functions lead to ulta-local scheme-independent terms in 4-point functions. As in the discussion above, the counterterms but with finite coefficients are related to/parametrise scheme-dependence of these correlators.

8.1 Counterterms and anomalies

In $d = 4$ the following counterterms can be introduced

$$S = S_{\rm CFT}[g^{\mu\nu}, A^a_\mu] + \int \mathrm{d}^4\boldsymbol{x}\sqrt{g}\left[\frac{\kappa_0}{4}F^a_{\mu\nu}F^{\mu\nu a} + a_0 E_4 + c_0 W^2 + b_0 R^2 + \tilde{b}_0 \Box R\right], \tag{8.1}$$

where E_4 is Euler density and W^2 is the square of Weyl tensor for the metric $g_{\mu\nu}$,

$$E_4 = R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} - 4R_{\mu\nu}R^{\mu\nu} + R^2, \tag{8.2}$$

$$W^2 = R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} - 2R_{\mu\nu}R^{\mu\nu} + \tfrac{1}{3}R^2. \tag{8.3}$$

The regularisation scheme is a dimensional regularisation with $d = 4-\epsilon$. Our conventions for the Riemann and Ricci tensors follow [51]. The constants in the counterterm action are functions of ϵ and are typically divergent. These divergences are required to cancel the corresponding singularities in the regularised solutions of the primary and secondary CWIs so that a finite limit exists as we send $\epsilon \to 0$. The counterterms (8.1) also necessarily contribute finite pieces, however, leading to trace anomalies. By taking a functional derivative and using the results of [52] one finds the following anomalous contribution to the trace Ward identity (4.61)

$$\langle T\rangle = \epsilon\left[\frac{\kappa_0}{4}F^a_{\mu\nu}F^{\mu\nu a} + a_0\left(E_4 + \frac{2}{3}\Box R\right) + c_0 W^2 + 12 b_0\sqrt{g}\Box R\right]. \tag{8.4}$$

With the usual representation for the anomalous trace of the stress-energy tensor

$$\langle T\rangle = \frac{\kappa}{4}F^a_{\mu\nu}F^{\mu\nu a} + aE_4 + cW^2 + b\Box R, \tag{8.5}$$

we find the anomalies

$$\kappa = \kappa_0^{(-1)}, \qquad a = a_0^{(-1)}, \qquad c = c_0^{(-1)}, \tag{8.6}$$

where $(\ldots)^{(-1)}$ denotes the term of order ϵ^{-1} in the series expansion of the coupling constant in the regulator.

8.2 $\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3} \rangle$

For the $\langle\!\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3} \rangle\!\rangle$ correlator only the second term in (8.1) contributes. The value of the constant κ_0 is fixed by a renormalisation of the 2-point function $\langle\!\langle J^\mu J^\nu \rangle\!\rangle$. Using the dimensional regularisation $d = 4 - \epsilon$, we find the 2-point function

$$\langle\!\langle J^\mu(\boldsymbol{p}) J^\nu(-\boldsymbol{p}) \rangle\!\rangle = \pi^{\mu\nu}(\boldsymbol{p}) p^2 \left(\frac{2c_J}{\epsilon} - c_J \log p^2 - \kappa_0 \right). \tag{8.7}$$

We therefore need

$$\kappa_0 = \frac{2c_J}{\epsilon} + O(\epsilon^0) \tag{8.8}$$

fixing the anomaly coefficient in (8.5) to be

$$\kappa = \kappa_0^{(-1)} = 2c_J. \tag{8.9}$$

With the value of κ_0 fixed, we can check that all divergences in the 3-point function now cancel as well. This is a non-trivial check on our solutions and the singularities of the triple-K integrals. We find that the counterterm contribution to the solution (7.7) is

$$A_1^{\text{anomaly}} = O(\epsilon^0), \tag{8.10}$$

$$A_2^{\text{anomaly}} = 2\kappa_0 = \frac{4c_J}{\epsilon} + O(\epsilon^0), \tag{8.11}$$

$$A_3^{\text{anomaly}} = 2\kappa_0 = \frac{4c_J}{\epsilon} + O(\epsilon^0), \tag{8.12}$$

$$A_4^{\text{anomaly}} = -\kappa_0(p_1^2 - p_2^2 - p_3^2) - 2c_3\kappa_0(p_2^2 + p_3^2) \tag{8.13}$$

$$= -\frac{2c_J}{\epsilon}(p_1^2 - p_2^2 - p_3^2) - \frac{4c_3 c_J}{\epsilon}(p_2^2 + p_3^2) + O(\epsilon^0), \tag{8.14}$$

where the constant c_J is defined in (5.65) and c_3 is defined in (7.22). The appearance of c_3 is due to the our definition of the 3-point function (7.32).

The result (8.10) - (8.14) is very constraining: in particular, the coefficients of the singular terms in the regularised solution must be multiples of c_J and not the undetermined primary constant α_1.

The divergences of the triple-K integrals entering the regularised solution can be evaluated using the method outlined in section 6.4, yielding

$$I_{5\{211\}} = \text{finite}, \tag{8.15}$$

$$I_{3+\epsilon\{211\}} = \frac{2}{\epsilon} + O(\epsilon^0), \tag{8.16}$$

$$I_{1+\epsilon\{211\}} = -\frac{p_2^2 + p_3^2}{\epsilon^2} + \frac{1}{\epsilon}\left[p_2^2 \log p_2 + p_3^2 \log p_3 \right. \tag{8.17}$$

$$\left. + (p_2^2 + p_3^2)\left(-\tfrac{1}{2} + \gamma_E - \log 2\right) - \tfrac{1}{2} p_1^2 \right]. \tag{8.18}$$

Substituting everything into the solution (7.7), one can check that all singularities cancel leaving a finite result. The scheme-dependent terms arise due to the $O(\epsilon^0)$ ambiguity in the definition of κ_0 in (8.8). Looking at (8.10) - (8.14), we see that the scheme-dependence may only change the transverse-traceless part according to

$$A_2 \mapsto A_2 + 2\kappa_0^{(0)}, \tag{8.19}$$

$$A_3 \mapsto A_3 + 2\kappa_0^{(0)}, \tag{8.20}$$

$$A_4 \mapsto A_4 - \kappa_0^{(0)}(p_1^2 - p_2^2 - p_3^2) - 2c_3\kappa_0^{(0)}(p_2^2 + p_3^2), \tag{8.21}$$

where $\kappa_0^{(0)}$ is the $O(\epsilon^0)$ part of κ_0.

Finally, one can look for anomalies in the trace of the stress-energy tensor. Due to the dimensional regularisation $d = 4-\epsilon$, the trace of the stress-energy tensor acquires a contribution from the counterterm

$$\langle T \rangle = \frac{\kappa}{4} F_{\mu\nu} F^{\mu\nu}, \qquad \kappa = \kappa_0^{(-1)} = 2c_J. \tag{8.22}$$

The anomalous term contributes to the trace Ward identity, which acquires a contribution

$$\langle\!\langle T(\boldsymbol{p}_1) J^{\mu_2}(\boldsymbol{p}_2) J^{\mu_3}(\boldsymbol{p}_3) \rangle\!\rangle_{\text{anomaly}} = \kappa \left[p_3^{\mu_2} p_2^{\mu_3} - \tfrac{1}{2}(p_1^2 - p_2^2 - p_3^2)\delta^{\mu_2\mu_3} \right] \tag{8.23}$$

modifying the form of the total 3-point function. In arbitrary dimension, the form of the anomaly is

$$\langle\!\langle T(\boldsymbol{p}_1) J^{\mu_2}(\boldsymbol{p}_2) J^{\mu_3}(\boldsymbol{p}_3) \rangle\!\rangle_{\text{anomaly}} = \pi^{\mu_2}_{\alpha_2}(\boldsymbol{p}_2) \pi^{\mu_3}_{\alpha_3}(\boldsymbol{p}_3) \left[B_1 p_3^{\alpha_2} p_1^{\alpha_3} + B_2 \delta^{\alpha_2\alpha_3} \right] \tag{8.24}$$

where the form factors B_1 and B_2 are functions of the momentum magnitudes. The contribution to the full 3-point function can then be recovered using (7.43), yielding

$$\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2}(\boldsymbol{p}_2) J^{\mu_3}(\boldsymbol{p}_3) \rangle\!\rangle_{\text{anomaly}} = \frac{\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)}{d-1} \pi^{\mu_2}_{\alpha_2}(\boldsymbol{p}_2) \pi^{\mu_3}_{\alpha_3}(\boldsymbol{p}_3) \left[B_1 p_3^{\alpha_2} p_1^{\alpha_3} + B_2 \delta^{\alpha_2\alpha_3} \right]. \tag{8.25}$$

In our example (8.23), in $d = 4$ we find

$$B_1 = -\kappa, \qquad B_2 = -\frac{\kappa}{2}(p_1^2 - p_2^2 - p_3^2). \tag{8.26}$$

A list of all anomalies and their contributions to all correlators is given in appendix A.8.

8.3 $\langle T^{\mu_1\nu_1} T^{\mu_2\nu_2} T^{\mu_3\nu_3} \rangle$

The situation for $\langle\!\langle T^{\mu_1\nu_1} T^{\mu_2\nu_2} T^{\mu_3\nu_3} \rangle\!\rangle$ is very similar to that for $\langle\!\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3} \rangle\!\rangle$. The $\Box R$ and R^2 in the action (8.1) can be omitted from the analysis since the former is a total derivative while the latter does not contribute to the transverse-traceless part of the correlators, and can instead be fixed by the renormalisation of the trace part.

The 2-point function in the presence of the counterterms is

$$\langle\!\langle T^{\mu_1\nu_1}(-\boldsymbol{p}) T^{\mu_2\nu_2}(\boldsymbol{p}) \rangle\!\rangle_{\text{reg}} = \Pi^{\mu_1\nu_1\mu_2\nu_2}(\boldsymbol{p}) p^4 \left[\frac{2c_T}{\epsilon} - c_T \log p^2 - 4c_0 + O(\epsilon) \right], \tag{8.27}$$

where we assume that the trace part was removed by adding the appropriate R^2 term. As we can see, the transverse-traceless part depends on c_0 only and requires

$$c_0 = \frac{c_T}{2\epsilon} + O(\epsilon^0). \tag{8.28}$$

The anomalous contributions to the transverse-traceless part of the 3-point function are then

$$A_1^{\text{anomaly}} = 0, \tag{8.29}$$

$$A_2^{\text{anomaly}} = -16(a_0 + c_0), \tag{8.30}$$

$$A_3^{\text{anomaly}} = 8 \left[c_0(p_1^2 + p_2^2) - a_0 p_3^2 \right], \tag{8.31}$$

$$A_4^{\text{anomaly}} = 8 \left[c_0(p_1^2 + p_2^2 + 3p_3^2) - a_0(p_1^2 + p_2^2 - p_3^2) \right], \tag{8.32}$$

$$A_5^{\text{anomaly}} = -4 \left[a_0 \lambda^2 + c_0(p_1^2 + p_2^2 + p_3^2)^2 + 8 c_g c_0 (p_1^4 + p_2^4 + p_3^4) \right], \tag{8.33}$$

where we used the definition of the 3-point function (4.47). The constant c_g is defined, in the case where $T_{\mu\nu}$ is the unique spin-2 conserved current, as

$$\frac{T_{\mu_1\nu_1}(\boldsymbol{x})}{g^{\mu_2\nu_2}(\boldsymbol{y})} = 4 c_g \delta_{(\mu_1(\mu_2} T_{\nu_1)\nu_2)}(\boldsymbol{x}) \delta(\boldsymbol{x} - \boldsymbol{y}) + \dots \tag{8.34}$$

or more generally

$$\langle\!\langle \frac{\delta T_{\mu_1\nu_1}}{\delta g^{\mu_2\nu_2}}(\boldsymbol{p}_1,\boldsymbol{p}_2)T_{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\!\rangle = 4c_g\delta_{(\mu_1(\mu_2}\langle\!\langle T_{\nu_1)\nu_2)}(\boldsymbol{p}_3)T_{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\!\rangle + \dots \tag{8.35}$$

where the omitted terms do not contain tensors we have listed explicitly.

As in the case of $\langle\!\langle T^{\mu_1\nu_1}J^{\mu_2}J^{\mu_3}\rangle\!\rangle$, we can find the divergences in the regularised form factors coming from the divergences in the triple-K integrals, giving

$$A_1^{\rm reg} = O(\epsilon^0), \tag{8.36}$$

$$A_2^{\rm reg} = \frac{8}{\epsilon}(16\alpha_1+\alpha_2) + O(\epsilon^0), \tag{8.37}$$

$$A_3^{\rm reg} = -\frac{4}{\epsilon}\left[(p_1^2+p_2^2)c_T + p_3^2(c_T-16\alpha_1-\alpha_2)\right] + O(\epsilon^0), \tag{8.38}$$

$$A_4^{\rm reg} = \frac{4}{\epsilon}\left[(p_1^2+p_2^2-p_3^2)(16\alpha_1+\alpha_2-2c_T) - 4c_Tp_3^2\right] + O(\epsilon^0), \tag{8.39}$$

$$A_5^{\rm reg} = \frac{2}{\epsilon}\left[(16\alpha_1+\alpha_2)\lambda^2 + 2c_T(1+4c_g)(p_1^4+p_2^4+p_3^4)\right] + O(\epsilon^0), \tag{8.40}$$

Since the value of c_0 is already fixed by (8.28), we can use one of the form factors, say A_2, to find

$$a_0 = \frac{1}{2\epsilon}(16\alpha_1+\alpha_2-c_T) + O(\epsilon^0). \tag{8.41}$$

The immediate conclusion is that all singularities must appear with coefficients that are multiples of $16\alpha_1+\alpha_2$ or c_T. As we can see, this is indeed the case. Substituting a_0 and c_0 as given by (8.41) and (8.28) into the remaining equations, we obtain an exact cancellation of divergences. The anomaly coefficients in (8.5) are thus

$$a = \frac{1}{2}(16\alpha_1+\alpha_2-c_T), \qquad c = \frac{c_T}{2}. \tag{8.42}$$

Finally, from (8.29) - (8.33), we find the scheme-dependent contributions take the form

$$A_2 \mapsto A_2 - 16(a_0^{(0)}+c_0^{(0)}), \tag{8.43}$$

$$A_3 \mapsto A_3 + 8\left[c_0^{(0)}(p_1^2+p_2^2) - a_0^{(0)}p_3^2\right], \tag{8.44}$$

$$A_4 \mapsto A_4 + 8\left[c_0^{(0)}(p_1^2+p_2^2+3p_3^2) - a_0^{(0)}(p_1^2+p_2^2-p_3^2)\right], \tag{8.45}$$

$$A_5 \mapsto A_5 - 4\left[a_0^{(0)}\lambda^2 + c_0^{(0)}(p_1^2+p_2^2+p_3^2)^2 + 8c_gc_0^{(0)}(p_1^4+p_2^4+p_3^4)\right], \tag{8.46}$$

where $a_0^{(0)}$ and $c_0^{(0)}$ are arbitrary terms of order ϵ^0 in a_0 and c_0.

9 Extensions

In this final section of Part I, we briefly discuss two extensions of the present analysis: how to write the results for tensor correlators in terms of a helicity basis, and the issues that arise when we try to generalise to higher-point correlation functions.

9.1 Helicity formalism

In the helicity formalism, one writes down a basis for the space of transverse and transverse-traceless tensors in terms of polarisation tensors $\xi_\mu^{(s)}$ and $\epsilon_{\mu\nu}^{(s)}$ respectively, where $s=\pm1$. Various

definitions exist for the polarisation tensors, for example

$$\sum_{s=\pm1} \xi_\mu^{(s)}(\boldsymbol{p})\bar{\xi}_\nu^{(s)}(\boldsymbol{p}) = \pi_{\mu\nu}(\boldsymbol{p}), \qquad \xi_\mu^{(s)}(\boldsymbol{p})\bar{\xi}^{(s')\mu}(\boldsymbol{p}) = \delta^{ss'}, \tag{9.1}$$

$$\frac{1}{2}\sum_{s=\pm1} \epsilon_{\mu\nu}^{(s)}(\boldsymbol{p})\bar{\epsilon}_{\rho\sigma}^{(s)}(\boldsymbol{p}) = \Pi_{\mu\nu\rho\sigma}(\boldsymbol{p}), \qquad \epsilon_{\mu\nu}^{(s)}(\boldsymbol{p})\bar{\epsilon}^{(s')\mu\nu}(\boldsymbol{p}) = 2\delta^{ss'}, \tag{9.2}$$

where the bar over a symbol denotes the complex conjugation. The helicity-projected operators are then

$$J^{(s)}(\boldsymbol{p}) = \bar{\xi}_\mu^{(s)}(\boldsymbol{p})J^\mu(\boldsymbol{p}), \qquad T^{(s)}(\boldsymbol{p}) = \frac{1}{2}\bar{\epsilon}_{\mu\nu}^{(s)}(\boldsymbol{p})T^{\mu\nu}(\boldsymbol{p}). \tag{9.3}$$

Correlation functions of the helicity-projected operators can easily be obtained from the transverse-traceless parts of the correlators. First observe that the semi-local parts of any correlation function vanish when contracted with polarisation tensors. Indeed, equations (9.1) and (9.2) imply that

$$\pi_\mu^\nu\bar{\xi}_\nu^{(s)} = \bar{\xi}_\mu^{(s)}, \qquad \Pi_{\mu\nu}^{\rho\sigma}\bar{\epsilon}_{\rho\sigma}^{(s)} = \bar{\epsilon}_{\mu\nu}^{(s)}. \tag{9.4}$$

Then, using equation (4.20) we can write

$$\bar{\xi}_\mu^{(s)} j_{\text{loc}}^\mu = \bar{\xi}_\nu^{(s)}\pi_\mu^\nu j_{\text{loc}}^\mu = 0 \tag{9.5}$$

and similarly $\bar{\epsilon}_{\mu\nu}^{(s)} t_{\text{loc}}^{\mu\nu} = 0$. To obtain correlation functions in the helicity formalism, one can therefore apply helicity projectors to the transverse-traceless parts of correlators only. Due to (9.4), the projectors (3.1) and (3.2) can then be removed as well. Finally, one needs to compute a small number of contractions of the helicity projectors with momenta and with the metric.

As an example, consider the $\langle\!\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}\mathcal{O}\rangle\!\rangle$ correlation function. Applying first the helicity projectors to its decomposition (3.10), we find

$$\begin{aligned}\langle\!\langle T^{(s_1)}(\boldsymbol{p}_1)T^{(s_2)}(\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle\!\rangle &= \frac{1}{4}\bar{\epsilon}_{\mu_1\nu_1}^{(s_1)}(\boldsymbol{p}_1)\bar{\epsilon}_{\mu_2\nu_2}^{(s_2)}(\boldsymbol{p}_2)\langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1)t^{\mu_2\nu_2}(\boldsymbol{p}_2)\mathcal{O}(\boldsymbol{p}_3)\rangle\!\rangle \\ &= \frac{1}{4}\Big[A_1\bar{\epsilon}_{\mu_1\nu_1}^{(s_1)}(\boldsymbol{p}_1)p_2^{\mu_1}p_2^{\nu_1}\bar{\epsilon}_{\mu_2\nu_2}^{(s_2)}(\boldsymbol{p}_2)p_3^{\mu_2}p_3^{\nu_2} + A_2\bar{\epsilon}_{\mu_1\alpha}^{(s_1)}(\boldsymbol{p}_1)\bar{\epsilon}_{\mu_2}^{(s_2)\alpha}(\boldsymbol{p}_2)p_2^{\mu_1}p_3^{\mu_2} \\ &\qquad + A_3\bar{\epsilon}_{\alpha\beta}^{(s_1)}(\boldsymbol{p}_1)\bar{\epsilon}^{(s_2)\alpha\beta}(\boldsymbol{p}_2)\Big].\end{aligned} \tag{9.6}$$

The contractions with helicity tensors depend on the precise definition of the latter and also the overall dimension. Let us consider, for example, the case where $d = 3$ and $\Delta_2 = \Delta_3 = 1$. The required contractions can be found in [17],

$$\bar{\epsilon}_{\mu_1\nu_1}^{(s_1)}(\boldsymbol{p}_1)p_2^{\mu_1}p_2^{\nu_1} = \frac{\lambda^2}{4\sqrt{2}p_1^2}, \tag{9.7}$$

$$\bar{\epsilon}_{\mu_1\alpha}^{(s_1)}(\boldsymbol{p}_1)\bar{\epsilon}_{\mu_2}^{(s_2)\alpha}(\boldsymbol{p}_2)p_2^{\mu_1}p_3^{\mu_2} = \frac{\lambda^2 S_3^{(s_1s_2)}}{16p_1^2p_2^2}, \tag{9.8}$$

$$\bar{\epsilon}_{\alpha\beta}^{(s_1)}(\boldsymbol{p}_1)\bar{\epsilon}^{(s_2)\alpha\beta}(\boldsymbol{p}_2) = \frac{(S_3^{(s_1s_2)})^2}{8p_1^2p_2^2}, \tag{9.9}$$

where λ^2 is defined in (6.21) and

$$S_3^{(s_1s_2)} = p_3^2 - (s_1p_1 + s_2p_2)^2. \tag{9.10}$$

Using (5.79) - (5.81) for the form factors, the most general solution is

$$\begin{aligned}\langle\!\langle T^{(s_1)}(\boldsymbol{p}_1)T^{(s_2)}(\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle &= \alpha_1^I\frac{3p_1p_2}{4p_3}\left(\frac{p_1+p_2-p_3}{p_1+p_2+p_3}\right)^2\delta^{s_1s_2} \\ &+ \frac{c_{\mathcal{O}}S_3^{(s_1s_2)}}{32p_1^2p_2^2p_3}\left[-2c_1^I\lambda^2 + \left((c_1^I-3c_2^I)(p_1^2+p_2^2) + 3(c_1^I+c_2^I)p_3^2\right)S_3^{(s_1s_2)}\right].\end{aligned} \tag{9.11}$$

The constants c_1^J and c_2^J are defined in (5.74) and $c_{\mathcal{O}}$ is the normalisation constant of the 2-point function $\langle\!\langle \mathcal{O}^J \mathcal{O}^I \rangle\!\rangle$ defined in (5.64).

As a check on our results in (7.7), we compared our solution with that obtained in [29] for the $\langle\!\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3} \rangle\!\rangle$ correlator of free scalars and fermions finding perfect agreement.

9.2 Higher-point correlation functions

It would be interesting to apply the present formalism to higher-point correlation functions in momentum space. Unfortunately, this seems to be a much more difficult task. In general, the ideas of the tensor decomposition described in section 3.1 are valid, but much less constraining. For concreteness, consider the $\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) T^{\mu_2\nu_2}(\boldsymbol{p}_2) \mathcal{O}(\boldsymbol{p}_3) \mathcal{O}(\boldsymbol{p}_4) \rangle\!\rangle$ correlation function. As there are now three independent momenta, each transverse or transverse-traceless projector (3.1) and (3.2) can be contracted with either of the two independent transverse momenta to yield a non-vanishing result. The decomposition of the transverse-traceless part of the correlation function under consideration is therefore

$$\begin{aligned}\langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1) t^{\mu_2\nu_2}(\boldsymbol{p}_2) \mathcal{O}(\boldsymbol{p}_3) \mathcal{O}(\boldsymbol{p}_4) \rangle\!\rangle &= \Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1) \Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2) \Bigg[\\ &\sum_{n_{1,2}\in\{2,3\}} \sum_{n_{3,4}\in\{3,4\}} A_{n_1 n_2 n_3 n_4} p_{n_1}^{\alpha_1} p_{n_2}^{\beta_1} p_{n_3}^{\alpha_2} p_{n_4}^{\beta_2} \\ &+ \delta^{\alpha_1\alpha_2} \sum_{n_1\in\{2,3\}} \sum_{n_2\in\{3,4\}} A_{n_1 n_2} p_{n_1}^{\beta_1} p_{n_2}^{\beta_2} + A_0\, \delta^{\alpha_1\alpha_2} \delta^{\beta_1\beta_2} \Bigg], \end{aligned} \tag{9.12}$$

where $A_{n_1 n_2 n_3 n_4}$, $A_{n_1 n_2}$ and A_0 are form factors. Initially, we thus have $2^4 = 16$ tensor structures following from the contractions of the transverse-traceless projectors with momenta, while in case of 3-point functions the corresponding tensor was unique.

The decomposition above is valid as long as $d \geq 4$. In case of $d = 3$, the metric $\delta^{\mu\nu}$ is not an independent tensor according to (3.5). In this case, the decomposition (9.12) can be truncated after the second line.

In the case of the correlator $\langle\!\langle T^{\mu_1\nu_1} T^{\mu_2\nu_2} T^{\mu_3\nu_3} T^{\mu_4\nu_4} \rangle\!\rangle$ our procedure reduces the number of independent tensor structures from the original 47 868 simple tensors built from the metric and three independent momenta down to 382 transverse-traceless tensor structures. This number, however, should be further diminished when the full symmetry group S_4 is imposed. For the 3-point function, the full symmetry group was automatically encoded in the decomposition, since only one momentum could appear under a chosen Lorentz index. In case of the 4-point function, the permutation group mixes various momenta and the fully symmetric structure is not clearly visible. For comparison, in the case of the 3-point function $\langle\!\langle T^{\mu_1\nu_1} T^{\mu_2\nu_2} T^{\mu_3\nu_3} \rangle\!\rangle$, 499 simple tensors built up from the metric and two independent momenta were reduced down to five transverse-traceless tensor structures.

In addition, the form factors are no longer functions of momentum magnitudes only. If we consider an n-point function in a d-dimensional CFT with $d \geq n$, then the scalar products

$$p_{ij} = \boldsymbol{p}_i \cdot \boldsymbol{p}_j, \quad i, j = 1, 2, \ldots, n, \quad i < j \tag{9.13}$$

are independent variables. In this case, the form factors can be regarded as functions of p_{ij}.

It would be interesting to work out the form of the conformal Ward identities for higher-point correlation functions. It would be vital to understand the objects corresponding to conformal ratios in momentum space. So far, we can only count the number of degrees of freedom in an n-point function in a d-dimensional CFT and compare it to the number of independent conformal ratios which is $n(n-3)/2$, assuming $d \geq n$ [36]. Indeed, the number of independent scalar products (9.13) is reduced by the $n-1$ differential equations (4.2) and (4.3) following from the

special conformal Ward identities, plus the additional constraint (4.1) from the dilatation Ward identity, leaving

$$\frac{n(n-1)}{2} - (n-1) - 1 = \frac{n(n-3)}{2} \tag{9.14}$$

in accordance with the number of conformal ratios.

List of results

Definitions

Here we collect together the necessary definitions and notation required to present our main results; further details may be found in the first part of the paper.

Basic conventions

We assume $d \geq 3$ Euclidean dimensions. Vectors are denoted by bold letters, *e.g.*, $\boldsymbol{p}_1$, but all results will be expressed in terms of the magnitudes of the momenta

$$p_j = |\boldsymbol{p}_j| = \sqrt{\boldsymbol{p}_j^2}, \qquad j = 1, 2, 3. \tag{R.1.1}$$

In particular, the form factors $A_j = A_j(p_1, p_2, p_3)$ are functions of the momentum magnitudes. Arrows denote the exchange of arguments, *e.g.*, $A_j(p_1 \leftrightarrow p_2) = A_j(p_2, p_1, p_3)$. If no arguments are given for a particular form factor then the standard ordering is assumed, $A_j(p_1, p_2, p_3)$.

To write the results in compact form, we frequently make use of the following symmetric polynomials in the momentum magnitudes

$$\begin{aligned} &a_{123} = p_1 + p_2 + p_3, \qquad && b_{123} = p_1p_2 + p_1p_3 + p_2p_3, \qquad && c_{123} = p_1p_2p_3, \\ &a_{ij} = p_i + p_j, && b_{ij} = p_ip_j, \end{aligned} \tag{R.1.2}$$

where $i, j = 1, 2, 3$. We also define the useful constant

$$s_d = \begin{cases} \frac{1}{\pi}(-1)^{\frac{d-1}{2}} & \text{if } d = 3, 5, 7, \dots \\ (-1)^{\frac{d}{2}-1} & \text{if } d = 4, 6, 8, \dots \end{cases} \tag{R.1.3}$$

Conformal Ward identities (CWIs) and triple-K integrals

The CWI operators K_j and K_{ij}, $i, j = 1, 2, 3$ are defined in (4.25) and (4.26) by

$$\mathrm{K}_j = \frac{\partial^2}{\partial p_j^2} + \frac{d + 1 - 2\Delta_j}{p_j} \frac{\partial}{\partial p_j}, \tag{R.1.4}$$

$$\mathrm{K}_{ij} = \mathrm{K}_i - \mathrm{K}_j\,. \tag{R.1.5}$$

By Δ_j, $j = 1, 2, 3$ we denote the conformal dimension of the j-th operator in a given 3-point function. For example in $\langle\!\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3} \rangle\!\rangle$ we have $\Delta_1 = d$ and $\Delta_2 = \Delta_3 = d - 1$.

The triple-K integral (5.1) and its reduced version (5.3) are

$$I_{\alpha\{\beta_1\beta_2\beta_3\}}(p_1, p_2, p_3) = \int_0^\infty \mathrm{d}x\, x^\alpha \prod_{j=1}^3 p_j^{\beta_j} K_{\beta_j}(p_j x), \tag{R.1.6}$$

$$J_{N\{k_j\}} = I_{\frac{d}{2}-1+N\{\Delta_j - \frac{d}{2} + k_j\}}, \tag{R.1.7}$$

where K_ν is the Bessel function K (modified Bessel function of the second kind) and we use a shortened notation $\{k_j\} = \{k_1k_2k_3\}$.

Triple-K integrals with half-integer β coefficients can be evaluated directly, while triple-K integrals with integer indices can be evaluated by means of the reduction scheme. These cases are sufficient to write the 3-point functions of conserved currents and stress-energy tensors in any dimension $d \geq 3$. In even dimensions the required integrals can all be evaluated from a single master integral $I_{0\{111\}}$. The result for this master integral is given in appendix A.6 and the reduction scheme for other necessary integrals is presented in table 1 on page 44.

Solutions to the primary CWIs, see section 5.2, are given as linear combinations of reduced triple-K integrals multiplied by constants, denoted by α_j, $j \in \mathbb{N}$ and called primary constants. If a primary constant is not restricted by means of the secondary CWIs, then it is a free parameter depending on the details of the theory.

If a triple-K integral diverges, then it can be regularised by

$$\alpha \mapsto \alpha + \epsilon, \qquad \beta_j \text{ does not change, } j = 1,2,3 \tag{R.1.8}$$

At leading order in ϵ, this scheme is equivalent (up to ultralocal terms) to the momentum space dimensional regularisation $d \mapsto d - \epsilon$, see section 5.3.4. If the regulator ϵ cannot be removed, then both triple-K integrals and primary constants are power series in ϵ. By $x^{(n)}$ we denote the coefficient of ϵ^n in the series expansion of x about $\epsilon = 0$, where the regularisation $\alpha \mapsto \alpha + \epsilon$ is assumed.

The differential operators appearing in the secondary CWIs are defined by (4.31) and (4.32) and read

$$\begin{aligned} \mathrm{L}_{s,N} &= p_1(p_1^2 + p_2^2 - p_3^2)\frac{\partial}{\partial p_1} + 2p_1^2 p_2 \frac{\partial}{\partial p_2} \\ &\quad + \left[(2d - \Delta_1 - 2\Delta_2 + s + N)p_1^2 + (\Delta_1 - 2 + s)(p_3^2 - p_2^2)\right], \end{aligned} \tag{R.1.9}$$

$$\mathrm{R}_s = p_1 \frac{\partial}{\partial p_1} - (\Delta_1 - 2 + s), \tag{R.1.10}$$

$$\mathrm{L}'_{s,N} = \mathrm{L}_{s,N} \text{ with } (p_1 \leftrightarrow p_2) \text{ and } (\Delta_1 \leftrightarrow \Delta_2), \tag{R.1.11}$$

$$\mathrm{R}'_s = \mathrm{R}_s \text{ with } (p_1 \to p_2) \text{ and } (\Delta_1 \to \Delta_2). \tag{R.1.12}$$

The secondary CWIs denoted by an asterisk are redundant, *i.e.*, they do not impose any additional constraints on primary constants, see section 7.2.

Tensor decomposition

Our standard conventions for Lorentz indices are discussed in section 3.1 (see in particular (3.7)) and read

$$\boldsymbol{p}_1, \boldsymbol{p}_2 \text{ for } \mu_1, \nu_1; \ \boldsymbol{p}_2, \boldsymbol{p}_3 \text{ for } \mu_2, \nu_2 \text{ and } \boldsymbol{p}_3, \boldsymbol{p}_1 \text{ for } \mu_3, \nu_3. \tag{R.1.13}$$

The transverse and transverse-traceless projectors (3.1) and (3.2) are

$$\pi^\mu_\alpha(\boldsymbol{p}) = \delta^\mu_\alpha - \frac{p^\mu p_\alpha}{p^2}, \tag{R.1.14}$$

$$\Pi^{\mu\nu}_{\alpha\beta}(\boldsymbol{p}) = \frac{1}{2}\left(\pi^\mu_\alpha(\boldsymbol{p})\pi^\nu_\beta(\boldsymbol{p}) + \pi^\mu_\beta(\boldsymbol{p})\pi^\nu_\alpha(\boldsymbol{p})\right) - \frac{1}{d-1}\pi^{\mu\nu}(\boldsymbol{p})\pi_{\alpha\beta}(\boldsymbol{p}). \tag{R.1.15}$$

The transverse(-traceless) and semi-local parts of the conserved current J^μ the stress-energy tensor $T^{\mu\nu}$ are given by (3.17) and (3.18) and read

$$j^\mu \equiv \pi^\mu_\alpha J^\alpha, \qquad j^\mu_{\text{loc}} \equiv J^\mu - j^\mu, \tag{R.1.16}$$

$$t^{\mu\nu} \equiv \Pi^{\mu\nu}_{\alpha\beta} T^{\alpha\beta}, \qquad t^{\mu\nu}_{\text{loc}} \equiv T^{\mu\nu} - t^{\mu\nu}. \tag{R.1.17}$$

The semi-local parts (denoted with the subscript 'loc') can also be expressed as

$$j^\mu_{\text{loc}} = \frac{p^\mu}{p^2} r, \qquad t^{\mu\nu}_{\text{loc}} = \frac{p^\mu}{p^2}R^\nu + \frac{p^\nu}{p^2}R^\mu - \frac{p^\mu p^\nu}{p^4}R + \frac{1}{d-1}\pi^{\mu\nu}\left(T - \frac{R}{p^2}\right), \tag{R.1.18}$$

where longitudinal and trace parts are

$$r = p_\mu J^\mu, \qquad R^\nu = p_\mu T^{\mu\nu}, \qquad R = p_\nu R^\nu, \qquad T = T^\mu_\mu. \tag{R.1.19}$$

It will also be useful to define the operator $\mathscr{T}^{\mu\nu}_\alpha$ as in (3.22), namely

$$\mathscr{T}^{\mu\nu}_\alpha(\boldsymbol{p}) = \frac{1}{p^2}\left[2p^{(\mu}\delta^{\nu)}_\alpha - \frac{p_\alpha}{d-1}\left(\delta^{\mu\nu} + (d-2)\frac{p^\mu p^\nu}{p^2}\right)\right]. \tag{R.1.20}$$

We also denote $\mathscr{T}^{\mu\nu\alpha} = \delta^{\alpha\beta}\mathscr{T}^{\mu\nu}_\beta$.

Operators in the theory

We assume the CFT contains the following data:

- A symmetry group G. The conserved current $J^{\mu a}$, $a = 1, \ldots, \dim G$, is then the Noether current associated with the symmetry and is sourced by a potential A^a_μ. Currents transform in the adjoint representation and we denote the structure constants as f^{abc}. We assume the Killing form is diagonal, $\mathrm{tr}(T^a T^b) = \frac{1}{2}\delta^{ab}$, where T^a are generators of the group.
- Scalar primary operators $\mathcal{O}^I$ all of the same dimension Δ. They are sourced by ϕ_0^I and transform in a representation R of the symmetry group. The representation matrices are denoted by $(T^a_R)^{IJ}$.
- A stress-energy tensor $T_{\mu\nu}$ sourced by a metric $g^{\mu\nu}$.

The relevant Ward identities in the CFT are discussed in section 4; in particular the transverse Ward identities (and our assumptions for the terms in these containing functional derivatives) are given in section 4.4.

The normalisation constants $c_{\mathcal{O}}$, c_J, c_T of 2-point functions are

$$\langle\!\langle \mathcal{O}^I(\boldsymbol{p})\mathcal{O}^J(-\boldsymbol{p})\rangle\!\rangle = c_{\mathcal{O}}\delta^{IJ} \times \begin{cases} p^{2\Delta-d} & \text{if } 2\Delta \neq d+2n,\ n\in\mathbb{N}, \\ p^{2\Delta-d}\left(-\log p^2 + \text{local}\right) & \text{if } 2\Delta = d+2n,\ n\in\mathbb{N} \end{cases} \tag{R.1.21}$$

$$\langle\!\langle J^{\mu a}(\boldsymbol{p})J^{\nu b}(-\boldsymbol{p})\rangle\!\rangle = c_J\delta^{ab} \times \begin{cases} \pi^{\mu\nu}(\boldsymbol{p})p^{d-2} & \text{if } d = 3,5,7,\ldots \\ \pi^{\mu\nu}(\boldsymbol{p})p^{d-2}\left(-\log p^2 + \text{local}\right) & \text{if } d = 4,6,8,\ldots \end{cases} \tag{R.1.22}$$

$$\langle\!\langle T^{\mu\nu}(\boldsymbol{p})T^{\rho\sigma}(-\boldsymbol{p})\rangle\!\rangle = c_T \times \begin{cases} \Pi^{\mu\nu\rho\sigma}(\boldsymbol{p})p^{d} & \text{if } d = 3,5,7,\ldots \\ \Pi^{\mu\nu\rho\sigma}(\boldsymbol{p})p^{d}\left(-\log p^2 + \text{local}\right) & \text{if } d = 4,6,8,\ldots \end{cases} \tag{R.1.23}$$

In the following, we will illustrate our general results with specific examples in $d = 3$, 4 and 5 dimensions. We consider for these purposes scalar operators both with dimensions $\Delta = d-2$ and with dimension $\Delta = d$. The former may be constructed as $\mathcal{O} = \phi^2$ in a theory of free scalars, where ϕ is the fundamental field, while the latter presents an interesting case being marginal.

$\langle\mathcal{O}\mathcal{O}\mathcal{O}\rangle$

$$\langle\!\langle \mathcal{O}^{I_1}(\boldsymbol{p}_1)\mathcal{O}^{I_2}(\boldsymbol{p}_2)\mathcal{O}^{I_3}(\boldsymbol{p}_3)\rangle\!\rangle = A_1^{I_1I_2I_3}(p_1,p_2,p_3), \tag{R.2.1}$$

The primary CWIs are

$$\mathrm{K}_{ij}\, A_1^{I_1I_2I_3} = 0, \qquad i,j = 1,2,3, \tag{R.2.2}$$

The solution in terms of triple-K integrals (R.1.7) is

$$A_1^{I_1I_2I_3} = \alpha_1^{I_1I_2I_3} J_{0\{000\}}, \tag{R.2.3}$$

where $\alpha_1^{I_1I_2I_3}$ is an arbitrary constant. (Note that primary constants inherit the group structure of the correlation function.) For any permutation σ of the set $\{1,2,3\}$ the A_1 form factor satisfies

$$A_1^{I_{\sigma(1)}I_{\sigma(2)}I_{\sigma(3)}}(p_{\sigma(1)},p_{\sigma(2)},p_{\sigma(3)}) = A_1^{I_1I_2I_3}(p_1,p_2,p_3). \tag{R.2.4}$$

$\langle J^{\mu_1}\mathcal{O}\mathcal{O}\rangle$

Ward identities. The transverse Ward identity is

$$\begin{aligned} &p_{1\mu_1}\langle\!\langle J^{\mu_1 a}(\boldsymbol{p}_1)\mathcal{O}^{I_2}(\boldsymbol{p}_2)\mathcal{O}^{I_3}(\boldsymbol{p}_3)\rangle\!\rangle = \\ &\quad = -(T^a_R)^{KI_3}\langle\!\langle \mathcal{O}^K(\boldsymbol{p}_2)\mathcal{O}^{I_2}(-\boldsymbol{p}_2)\rangle\!\rangle - (T^a_R)^{KI_2}\langle\!\langle \mathcal{O}^K(\boldsymbol{p}_3)\mathcal{O}^{I_3}(-\boldsymbol{p}_3)\rangle\!\rangle. \end{aligned} \tag{R.3.1}$$

Reconstruction formula. The full 3-point function can be reconstructed from the transverse-traceless part as

$$\begin{aligned}\langle\!\langle J^{\mu_1 a}(\boldsymbol{p}_1)\mathcal{O}^{I_2}(\boldsymbol{p}_2)\mathcal{O}^{I_3}(\boldsymbol{p}_3)\rangle\!\rangle &= \langle\!\langle j^{\mu_1 a}(\boldsymbol{p}_1)\mathcal{O}^{I_2}(\boldsymbol{p}_2)\mathcal{O}^{I_3}(\boldsymbol{p}_3)\rangle\!\rangle \\ &- \frac{p_1^{\mu_1}}{p_1^2}\left[(T_R^a)^{KI_3}\langle\!\langle\mathcal{O}^K(\boldsymbol{p}_2)\mathcal{O}^{I_2}(-\boldsymbol{p}_2)\rangle\!\rangle + (T_R^a)^{KI_2}\langle\!\langle\mathcal{O}^K(\boldsymbol{p}_3)\mathcal{O}^{I_3}(-\boldsymbol{p}_3)\rangle\!\rangle\right].\end{aligned} \tag{R.3.2}$$

Decomposition of the 3-point function. The tensor decomposition of the transverse-traceless part is

$$\langle\!\langle j^{\mu_1 a}(\boldsymbol{p}_1)\mathcal{O}^{I_2}(\boldsymbol{p}_2)\mathcal{O}^{I_3}(\boldsymbol{p}_3)\rangle\!\rangle = \pi^{\mu_1}_{\alpha_1}(\boldsymbol{p}_1)\cdot A_1^{aI_2I_3}p_2^{\alpha_1}, \tag{R.3.3}$$

where the form factor A_1 depends on the momentum magnitudes. This form factor is symmetric under $(p_2, I_2) \leftrightarrow (p_3, I_3)$, *i.e.*,

$$A_1^{aI_3I_2}(p_1, p_3, p_2) = A_1^{aI_2I_3}(p_1, p_2, p_3). \tag{R.3.4}$$

This form factor is given by

$$A_1^{aI_2I_3} = \text{coefficient of } p_2^{\mu_1} \text{ in } \langle\!\langle J^{\mu_1 a}(\boldsymbol{p}_1)\mathcal{O}^{I_2}(\boldsymbol{p}_2)\mathcal{O}^{I_3}(\boldsymbol{p}_3)\rangle\!\rangle. \tag{R.3.5}$$

Primary conformal Ward identities. The primary CWIs are

$$\mathrm{K}_{ij}\, A_1^{aI_2I_3} = 0, \qquad i,j = 1,2,3, \tag{R.3.6}$$

The solution in terms of triple-K integrals (R.1.7) is

$$A_1^{aI_2I_3} = \alpha_1^{aI_2I_3} J_{1\{000\}}, \tag{R.3.7}$$

where $\alpha_1^{aI_2I_3}$ is a constant. In particular $\alpha_1^{aI_3I_2} = \alpha_1^{aI_2I_3}$. If the integral diverges, the regularisation (R.1.8) should be used.

Secondary conformal Ward identities. The independent secondary CWI is

$$\mathrm{L}_{1,0}\, A_1^{aI_2I_3} = 2(d-2)\left[p_{1\mu_1}\langle\!\langle J^{\mu_1 a}(\boldsymbol{p}_1)\mathcal{O}^{I_2}(\boldsymbol{p}_2)\mathcal{O}^{I_3}(\boldsymbol{p}_3)\rangle\!\rangle\right], \tag{R.3.8}$$

where $\mathrm{L}_{s,N}$ is given by (R.1.9). Assuming the unitarity bound $\Delta_2 = \Delta_3 = \Delta \geq \frac{d}{2} - 1$ for the dimensions of the scalar operators we find

$$\alpha_1^{aI_2I_3} = -\frac{2^{4-\frac{d}{2}}(T_R^a)^{I_2I_3}c_{\mathcal{O}}}{\Gamma\left(\frac{d}{2}-1\right)}\left(1 - \frac{1}{2}\delta_{\Delta,d+n}\right)\times C, \tag{R.3.9}$$

where

$$C = \begin{cases} \frac{1}{\pi}\sin\left(\frac{\pi}{2}(d-2\Delta)\right) & \text{if } 2\Delta \neq d+2n,\ n\in\mathbb{N}, \\ (-1)^{\Delta-\frac{d}{2}} & \text{if } 2\Delta = d+2n,\ n\in\mathbb{N}. \end{cases} \tag{R.3.10}$$

and $c_{\mathcal{O}}$ represents the normalisation of the 2-point function (R.1.21). The 3-point function $\langle\!\langle J^{\mu_1}\mathcal{O}\mathcal{O}\rangle\!\rangle$ is therefore completely determined in terms of this normalisation.

Examples

For $d=3$ and $\Delta_2=\Delta_3=1$ we find

$$A_1^{aI_2I_3} = -\frac{2(T_R^a)^{I_2I_3}c_{\mathcal{O}}}{b_{23}a_{123}}. \tag{R.3.11}$$

For $d=3$ and $\Delta_2=\Delta_3=3$ we find

$$A_1^{aI_2I_3} = -p_1(T_R^a)^{I_2I_3}c_{\mathcal{O}}\log a_{123} + \frac{c_{\mathcal{O}}(T_R^a)^{I_2I_3}}{a_{123}}\left[p_2^2+p_3^2-p_1^2+p_2p_3+p_1a_{123}(2-\gamma_E)\right]. \tag{R.3.12}$$

For $d=4$ and $\Delta_2=\Delta_3=2$ we find

$$A_1^{aI_2I_3} = 4p_1(T_R^a)^{I_2I_3}c_{\mathcal{O}}\frac{\partial}{\partial p_1}I_{1\{000\}}, \tag{R.3.13}$$

where the $I_{1\{000\}}$ integral is given by (A.6.9).

For $d=4$ and $\Delta_2=\Delta_3=4$ we find

$$A_1^{aI_2I_3} = -2(T_R^a)^{I_2I_3}c_{\mathcal{O}}I^{(0)}_{2\{122\}}, \tag{R.3.14}$$

where $I^{(0)}_{2\{122\}}$ denotes the coefficient of ϵ^0 in the series expansion of the regulated integral $I_{2+\epsilon\{122\}}$. The integral can be obtained from the master integral (A.6.1) via the reduction scheme in table 1, page 44.

For $d=5$ and $\Delta_2=\Delta_3=3$ we find

$$A_1^{aI_2I_3} = 2(T_R^a)^{I_2I_3}c_{\mathcal{O}}\frac{p_1+a_{123}}{a_{123}^2}. \tag{R.3.15}$$

For $d=5$ and $\Delta_2=\Delta_3=5$ we find

$$\begin{aligned}A_1^{aI_2I_3} = -3p_1^3(T_R^a)^{I_2I_3}c_{\mathcal{O}}\log a_{123} + \frac{(T_R^a)^{I_2I_3}c_{\mathcal{O}}}{a_{123}^2}\Big[&p_1^5(4-3\gamma_E)+p_1^4(11-6\gamma_E)a_{23}\\ &- p_1^3\left((-10+3\gamma_E)a_{23}^2+3b_{23}\right)+2a_{23}p_1^2(2a_{23}^2-3b_{23})\\ &+(2p_1+a_{23})(a_{23}^4-3a_{23}^2b_{23}+b_{23}^2)\Big].\end{aligned} \tag{R.3.16}$$

$\langle J^{\mu_1}J^{\mu_2}\mathcal{O}\rangle$

Ward identities. The transverse Ward identity is

$$p_{1\mu_1}\langle\!\langle J^{\mu_1a_1}(\boldsymbol{p}_1)J^{\mu_2a_2}(\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle = p_{1\mu_1}\langle\!\langle\frac{\delta J^{\mu_1a_1}}{\delta A^{a_2}_{\mu_2}}(\boldsymbol{p}_1,\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle. \tag{R.4.1}$$

Reconstruction formula. The full 3-point function can be reconstructed from the transverse-traceless part as

$$\begin{aligned}\langle\!\langle J^{\mu_1a_1}(\boldsymbol{p}_1)J^{\mu_2a_2}(\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle &= \langle\!\langle j^{\mu_1a_1}(\boldsymbol{p}_1)j^{\mu_2a_2}(\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle\\ &+ \frac{p_1^{\mu_1}p_{1\alpha}}{p_1^2}\langle\!\langle\frac{\delta J^{\alpha a_1}}{\delta A^{a_2}_{\mu_2}}(\boldsymbol{p}_1,\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle + \frac{p_2^{\mu_2}p_{2\beta}}{p_2^2}\langle\!\langle\frac{\delta J^{\beta a_2}}{\delta A^{a_1}_{\mu_1}}(\boldsymbol{p}_2,\boldsymbol{p}_1)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle\\ &- \frac{p_1^{\mu_1}p_2^{\mu_2}p_{1\alpha}p_{2\beta}}{p_1^2p_2^2}\langle\!\langle\frac{\delta J^{\alpha a_1}}{\delta A^{a_2}_{\beta}}(\boldsymbol{p}_1,\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle.\end{aligned} \tag{R.4.2}$$

Decomposition of the 3-point function. The tensor decomposition of the transverse-traceless part is

$$\langle\!\langle j^{\mu_1 a_1}(\boldsymbol{p}_1) j^{\mu_2 a_2}(\boldsymbol{p}_2) \mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle = \pi^{\mu_1}_{\alpha_1}(\boldsymbol{p}_1)\pi^{\mu_2}_{\alpha_2}(\boldsymbol{p}_2)\left[A_1^{a_1 a_2 I} p_2^{\alpha_1} p_3^{\alpha_2} + A_2^{a_1 a_2 I}\delta^{\alpha_1\alpha_2}\right]. \tag{R.4.3}$$

The form factors A_1 and A_2 are functions of the momentum magnitudes. Both form factors are symmetric under $(p_1, a_1) \leftrightarrow (p_2, a_2)$, *i.e.*, they satisfy

$$A_j^{a_2 a_1 I}(p_2, p_1, p_3) = A_j^{a_1 a_2 I}(p_1, p_2, p_3), \qquad j = 1, 2. \tag{R.4.4}$$

These form factors can be calculated as follows

$$A_1^{a_1 a_2 I} = \text{coefficient of } p_2^{\mu_1} p_3^{\mu_2}, \tag{R.4.5}$$

$$A_2^{a_1 a_2 I} = \text{coefficient of } \delta^{\mu_1\mu_2} \tag{R.4.6}$$

in $\langle\!\langle J^{\mu_1 a_1}(\boldsymbol{p}_1) J^{\mu_2 a_2}(\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle$.

Primary conformal Ward identities. The primary CWIs are

$$\begin{aligned} &\mathrm{K}_{12}\, A_1^{a_1 a_2 I} = 0, && \mathrm{K}_{13}\, A_1^{a_1 a_2 I} = 0, \\ &\mathrm{K}_{12}\, A_2^{a_1 a_2 I} = 0, && \mathrm{K}_{13}\, A_2^{a_1 a_2 I} = 2A_1^{a_1 a_2 I}, \end{aligned} \tag{R.4.7}$$

The solution in terms of triple-K integrals (R.1.7) is

$$A_1^{a_1 a_2 I} = \alpha_1^{a_1 a_2 I} J_{2\{000\}}, \tag{R.4.8}$$

$$A_2^{a_1 a_2 I} = \alpha_1^{a_1 a_2 I} J_{1\{001\}} + \alpha_2^{a_1 a_2 I} J_{0\{000\}}, \tag{R.4.9}$$

where $\alpha_j^{a I_2 I_3}$, $j = 1, 2$ are constants. In particular $\alpha_j^{a_2 a_1 I} = \alpha_j^{a_1 a_2 I}$ for $j = 1, 2$. If the integrals diverge, the regularisation (R.1.8) should be used.

Secondary conformal Ward identities. The independent secondary CWI is

$$\begin{aligned} &\mathrm{L}_{1,0}\, A_1^{a_1 a_2 I} + 2\,\mathrm{R}_1\, A_2^{a_1 a_2 I} = \\ &\quad = 2(d-2)\cdot \text{coefficient of } p_3^{\mu_2} \text{ in } p_{1\mu_1}\langle\!\langle J^{\mu_1 a_1}(\boldsymbol{p}_1) J^{\mu_2 a_2}(\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle, \end{aligned} \tag{R.4.10}$$

where L and R operators are given by (R.1.9) and (R.1.10). This leads to

$$\alpha_2^{(0) a_1 a_2 I} = \frac{1}{2}\Delta_3(d - \Delta_3 - 2)\alpha_1^{a_1 a_2 I}, \tag{R.4.11}$$

$$\alpha_2^{(1) a_1 a_2 I} = \begin{cases} \frac{1}{2}(2\Delta_3 - d - 2)\alpha_1^{a_1 a_2 I} & \text{if } \Delta_3 \neq d - 2 - 2n,\ n \in \mathbb{N}, \\ -\frac{1}{2}(2\Delta_3 - d + 2)\alpha_1 + C c^{a_1 a_2 I} c_{\mathcal{O}} & \text{if } \Delta_3 = d - 2 - 2n,\ n \in \mathbb{N}, \end{cases} \tag{R.4.12}$$

$$\alpha_2^{(2) a_1 a_2 I} = -\frac{1}{2}\alpha_1^{a_1 a_2 I}, \tag{R.4.13}$$

where Δ_3 is the conformal dimension of the scalar operator satisfying the unitarity bound and $c_{\mathcal{O}}$ is the normalisation of the 2-point function (R.1.21). By $\alpha_j^{(n)}$ we mean the coefficient of ϵ^n in the series expansion of α_j, and we assume that the constant α_1 does not depend on the regulator, *i.e.*, $\alpha_1^{a_1 a_2 I} = \alpha_1^{(0) a_1 a_2 I}$. We denote

$$C = \frac{(-1)^{\frac{d-\Delta_3}{2}} 2^{2+\Delta_3-\frac{d}{2}}(d-2)\Gamma\left(\frac{d-\Delta_3}{2}\right)\Gamma\left(\frac{\Delta_3+1}{2}\right)}{\sqrt{\pi}\,\Gamma\left(\frac{\Delta_3}{2}\right)\Gamma\left(\frac{d+\Delta_3}{2}\right)} \times \begin{cases} \frac{1}{\Gamma\left(\frac{d-2\Delta_3}{2}\right)} & \text{if } 2\Delta_3 \neq d + 2n, \\ -\Gamma\left(\Delta_3 - \frac{d}{2} + 1\right) & \text{if } 2\Delta_3 = d + 2n, \end{cases} \tag{R.4.14}$$

and we define the $c^{a_1a_2K}$ constant as

$$p_{1\mu_1}\langle\!\langle \frac{\delta J^{\mu_1 a_1}}{\delta A^{a_2}_{\mu_2}}(\boldsymbol{p}_1,\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle\Big|_{p_1=p_2=p} = p_3^{\mu_2}\cdot c^{a_1a_2K}p^{d-\Delta_3-2}\langle\!\langle \mathcal{O}^K(\boldsymbol{p}_3)\mathcal{O}^I(-\boldsymbol{p}_3)\rangle\!\rangle + \dots \quad \text{(R.4.15)}$$

What we mean here is the following: we first write down the most general tensor decomposition of $p_{1\mu_1}\langle\!\langle \frac{\delta J^{\mu_1 a_1}}{\delta A^{a_2}_{\mu_2}}(\boldsymbol{p}_1,\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle$ and take the coefficient of $p_3^{\mu_2}$. We then set $p_1 = p_2 = p$ in the expression for this coefficient. If the regulator is present, the limit $\epsilon \to 0$ can be taken after the primary constants and triple-K integrals are substituted into the primary CWIs (R.4.8) - (R.4.9).

To summarise, the 3-point function $\langle\!\langle J^{\mu_1}J^{\mu_2}\mathcal{O}\rangle\!\rangle$ depends on the 2-point function normalisations $c_{\mathcal{O}}$ and $c^{a_1a_2K}$, and on one undetermined primary constant $\alpha_1^{a_1a_2I}$.

Examples

For $d = 3$ and $\Delta_3 = 1$ we find

$$A_1^{a_1a_2I} = \frac{\alpha_1^{a_1a_2I}}{p_3 a_{123}^2}, \quad \text{(R.4.16)}$$

$$A_2^{a_1a_2I} = \alpha_1^{a_1a_2I}\left(\frac{1}{a_{123}} - \frac{1}{2p_3}\right) - \frac{c^{a_1a_2I}c_{\mathcal{O}}}{p_3}. \quad \text{(R.4.17)}$$

For $d = 3$ and $\Delta_3 = 3$ we find

$$A_1^{a_1a_2I} = \alpha_1^{a_1a_2I}\frac{a_{123}+p_3}{a_{123}^2}, \quad \text{(R.4.18)}$$

$$A_2^{a_1a_2I} = -\alpha_1^{a_1a_2I}\frac{-2p_3^2 + p_3a_{12} + a_{12}^2}{2a_{123}}. \quad \text{(R.4.19)}$$

For $d = 4$ and $\Delta_3 = 2$ we find

$$A_1^{a_1a_2I} = \alpha_1^{a_1a_2I}p_1p_2\frac{\partial^2}{\partial p_1 p_2}I_{1\{000\}}, \quad \text{(R.4.20)}$$

$$A_2^{a_1a_2I} = \alpha_1^{a_1a_2I}I^{(0)}_{2\{111\}} + (\alpha_1^{a_1a_2I} - 2c^{a_1a_2I}c_{\mathcal{O}})(\log p_3 - \log 2 + \gamma_E), \quad \text{(R.4.21)}$$

where $I^{(0)}_{\alpha\{\beta_j\}}$ denotes the coefficient of ϵ^0 in the series expansion of the regulated integral $I_{\alpha+\epsilon\{\beta_j\}}$. Both $I_{1\{000\}}$ and $I_{2\{111\}}$ integrals are given in appendix (A.6).

For $d = 4$ and $\Delta_3 = 4$ we find

$$A_1^{a_1a_2I} = \alpha_1^{a_1a_2I}I^{(0)}_{3\{112\}}, \quad \text{(R.4.22)}$$

$$A_2^{a_1a_2I} = \alpha_1^{a_1a_2I}\Big[-p_3\frac{\partial}{\partial p_3}I^{(0)}_{1\{112\}} + p_1^2\log p_1 + p_2^2\log p_2 - (p_1^2+p_2^2)\left(\tfrac{1}{2} - \gamma_E + \log 2\right) - \tfrac{1}{2}p_3^2\Big], \quad \text{(R.4.23)}$$

where $I^{(0)}_{\alpha\{\beta_j\}}$ denotes the coefficient of ϵ^0 in the series expansion of the regulated integral $I_{\alpha+\epsilon\{\beta_j\}}$. The integrals can be obtained from the master integral (A.6.1) via the reduction scheme in table 1, page 44.

For $d = 5$ and $\Delta_3 = 3$ we find

$$A_1^{a_1a_2I} = \alpha_1^{a_1a_2I} \frac{p_3^2 + 3p_3a_{12} + 2(a_{12}^2 + b_{12})}{a_{123}^3}, \tag{R.4.24}$$

$$A_2^{a_1a_2I} = \alpha_1^{a_1a_2I} \frac{p_3^3 + 2p_3^2a_{12} + p_3(-a_{12}^2 + 4b_{12}) + 2a_{13}(-a_{12}^2 + b_{12})}{2a_{123}^2} - p_3 c^{a_1a_2I} c_{\mathcal{O}}. \tag{R.4.25}$$

For $d = 5$ and $\Delta_3 = 5$ we find

$$A_1^{a_1a_2I} = \frac{\alpha_1^{a_1a_2I}}{a_{123}^3} \left[-2p_3^4 - 6p_3^3a_{12} - 2p_3^2(5a_{12}^2 - 4b_{12}) - 3a_{12}(a_{12} + 3p_3)(a_{12}^2 - b_{12})\right], \tag{R.4.26}$$

$$\begin{aligned} A_2^{a_1a_2I} = \frac{\alpha_1^{a_1a_2I}}{2a_{123}^2} \big[&-2p_3^5 - 4p_3^4a_{12} - 4p_3^3(a_{12}^2 - b_{12}) + p_3^2a_{12}(a_{12}^2 - 7b_{12}) \\ &+ 3a_{12}^2(2p_3 + a_{12})(a_{12}^2 - 3b_{12})\big]. \end{aligned} \tag{R.4.27}$$

$\left\langle J^{\mu_1} J^{\mu_2} J^{\mu_3} \right\rangle$

Ward identities. The transverse Ward identity is

$$\begin{aligned} &p_{1\mu_1} \langle\!\langle J^{\mu_1 a_1}(\boldsymbol{p}_1) J^{\mu_2 a_2}(\boldsymbol{p}_2) J^{\mu_3 a_3}(\boldsymbol{p}_3) \rangle\!\rangle = \\ &\quad = \mathrm{i} f^{a_1 b a_3} \langle\!\langle J^{\mu_3 b}(\boldsymbol{p}_2) J^{\mu_2 a_2}(-\boldsymbol{p}_2) \rangle\!\rangle - \mathrm{i} f^{a_1 a_2 b} \langle\!\langle J^{\mu_2 b}(\boldsymbol{p}_3) J^{\mu_3 a_3}(-\boldsymbol{p}_3) \rangle\!\rangle, \end{aligned} \tag{R.5.1}$$

where f^{abc} are the structure constants of the symmetry group.

Reconstruction formula. The full 3-point function can be reconstructed from the transverse-traceless part as

$$\begin{aligned} \langle\!\langle J^{\mu_1 a_1}(\boldsymbol{p}_1) J^{\mu_2 a_2}(\boldsymbol{p}_2) J^{\mu_3 a_3}(\boldsymbol{p}_3) \rangle\!\rangle &= \langle\!\langle j^{\mu_1 a_1}(\boldsymbol{p}_1) j^{\mu_2 a_2}(\boldsymbol{p}_2) j^{\mu_3 a_3}(\boldsymbol{p}_3) \rangle\!\rangle \\ &+ \left[\frac{p_1^{\mu_1}}{p_1^2} \left(\mathrm{i} f^{a_1 b a_3} \langle\!\langle J^{\mu_3 b}(\boldsymbol{p}_2) J^{\mu_2 a_2}(-\boldsymbol{p}_2) \rangle\!\rangle - \mathrm{i} f^{a_1 a_2 b} \langle\!\langle J^{\mu_2 b}(\boldsymbol{p}_3) J^{\mu_3 a_3}(-\boldsymbol{p}_3) \rangle\!\rangle\right)\right] \\ &\quad + [(\mu_1, a_1, \boldsymbol{p}_1) \leftrightarrow (\mu_2, b_2, \boldsymbol{p}_2)] + [(\mu_1, a_1, \boldsymbol{p}_1) \leftrightarrow (\mu_3, a_3, \boldsymbol{p}_3)] \\ &+ \left[\frac{p_1^{\mu_1} p_2^{\mu_2}}{p_1^2 p_2^2} \mathrm{i} f^{a_1 a_2 b} p_{2\alpha} \langle\!\langle J^{\alpha b}(\boldsymbol{p}_3) J^{\mu_3 a_3}(-\boldsymbol{p}_3) \rangle\!\rangle\right] \\ &\quad + [(\mu_1, a_1, \boldsymbol{p}_1) \leftrightarrow (\mu_3, a_3, \boldsymbol{p}_3)] + [(\mu_2, a_2, \boldsymbol{p}_2) \leftrightarrow (\mu_3, a_3, \boldsymbol{p}_3)]. \end{aligned} \tag{R.5.2}$$

Decomposition of the 3-point function. The tensor decomposition of the transverse-traceless part is

$$\begin{aligned} \langle\!\langle j^{\mu_1 a_1}(\boldsymbol{p}_1) j^{\mu_2 a_2}(\boldsymbol{p}_2) j^{\mu_3 a_3}(\boldsymbol{p}_3) \rangle\!\rangle = \pi_{\alpha_1}^{\mu_1}(\boldsymbol{p}_1) \pi_{\alpha_2}^{\mu_2}(\boldsymbol{p}_2) \pi_{\alpha_3}^{\mu_3}(\boldsymbol{p}_3) \big[&A_1^{a_1a_2a_3} p_2^{\alpha_1} p_3^{\alpha_2} p_1^{\alpha_3} \\ &+ A_2^{a_1a_2a_3} \delta^{\alpha_1\alpha_2} p_1^{\alpha_3} + A_2^{a_1a_3a_2}(p_2 \leftrightarrow p_3) \delta^{\alpha_1\alpha_3} p_3^{\alpha_2} \\ &+ A_2^{a_3a_2a_1}(p_1 \leftrightarrow p_3) \delta^{\alpha_2\alpha_3} p_2^{\alpha_1}\big]. \end{aligned} \tag{R.5.3}$$

The form factors A_1 and A_2 are functions of the momentum magnitudes. If no arguments are given, then we assume the standard ordering, $A_j = A_j(p_1, p_2, p_3)$, while by $p_i \leftrightarrow p_j$ we denote the exchange of the two momenta, *e.g.*, $A_2(p_1 \leftrightarrow p_3) = A_2(p_3, p_2, p_1)$.

The A_1 factor is completely antisymmetric, *i.e.*, for any permutation σ of the set $\{1, 2, 3\}$ it satisfies

$$A_1^{a_{\sigma(1)}a_{\sigma(2)}a_{\sigma(3)}}(p_{\sigma(1)}, p_{\sigma(2)}, p_{\sigma(3)}) = (-1)^\sigma A_1^{a_1a_2a_3}(p_1, p_2, p_3), \tag{R.5.4}$$

where $(-1)^\sigma$ denotes the sign of the permutation σ. The form factors A_2 is antisymmetric under $(p_1, a_1) \leftrightarrow (p_2, a_2)$, *i.e.*,

$$A_2^{a_2 a_1 a_3}(p_2, p_1, p_3) = -A_2^{a_1 a_2 a_3}(p_1, p_2, p_3). \tag{R.5.5}$$

The form factors can be calculated as follows

$$A_1^{a_1 a_2 a_3} = \text{coefficient of } p_2^{\mu_1} p_3^{\mu_2} p_1^{\mu_3}, \tag{R.5.6}$$
$$A_2^{a_1 a_2 a_3} = \text{coefficient of } \delta^{\mu_1 \mu_2} p_1^{\mu_3} \tag{R.5.7}$$

in $\langle\!\langle J^{\mu_1 a_1}(\boldsymbol{p}_1) J^{\mu_2 a_2}(\boldsymbol{p}_2) J^{\mu_3 a_3}(\boldsymbol{p}_3) \rangle\!\rangle$.

Primary conformal Ward identities. The primary CWIs are

$$\begin{aligned} &\mathrm{K}_{12}\, A_1^{a_1 a_2 a_3} = 0, && \mathrm{K}_{13}\, A_1^{a_1 a_2 a_3} = 0, \\ &\mathrm{K}_{12}\, A_2^{a_1 a_2 a_3} = 0, && \mathrm{K}_{13}\, A_2^{a_1 a_2 a_3} = 2A_1^{a_1 a_2 a_3}. \end{aligned} \tag{R.5.8}$$

The solution in terms of triple-K integrals (R.1.7) is

$$A_1^{a_1 a_2 a_3} = \alpha_1^{a_1 a_2 a_3} J_{3\{000\}}, \tag{R.5.9}$$
$$A_2^{a_1 a_2 a_3} = \alpha_1^{a_1 a_2 a_3} J_{2\{001\}} + \alpha_2^{a_1 a_2 a_3} J_{1\{000\}}, \tag{R.5.10}$$

where $\alpha_j^{a_1 a_2 a_3}$, $j = 1, 2$ are constants. If the integrals diverge, the regularisation (R.1.8) should be used.

Secondary conformal Ward identities. The independent secondary CWIs are

$$\begin{aligned} (*)\ & \mathrm{L}_{1,2}\, A_1^{a_1 a_2 a_3} + 2\,\mathrm{R}_1\, [A_2^{a_1 a_2 a_3} - A_2^{a_1 a_3 a_2}(p_2 \leftrightarrow p_3)] = \\ &\quad = 2(d-2) \cdot \text{coefficient of } p_3^{\mu_2} p_1^{\mu_3} \text{ in } p_{1\mu_1} \langle\!\langle J^{\mu_1 a_1}(\boldsymbol{p}_1) J^{\mu_2 a_2}(\boldsymbol{p}_2) J^{\mu_3 a_3}(\boldsymbol{p}_3) \rangle\!\rangle, \end{aligned} \tag{R.5.11}$$
$$\begin{aligned} &\mathrm{L}_{1,0}\, [A_2^{a_3 a_2 a_1}(p_1 \leftrightarrow p_3)] + 2p_1^2\, [A_2^{a_1 a_3 a_2}(p_2 \leftrightarrow p_3) - A_2^{a_1 a_2 a_3}] = \\ &\quad = 2(d-2) \cdot \text{coefficient of } \delta^{\mu_2 \mu_3} \text{ in } p_{1\mu_1} \langle\!\langle J^{\mu_1 a_1}(\boldsymbol{p}_1) J^{\mu_2 a_2}(\boldsymbol{p}_2) J^{\mu_3 a_3}(\boldsymbol{p}_3) \rangle\!\rangle, \end{aligned} \tag{R.5.12}$$

where L and R operators are given by (4.31) and (4.32). The identity denoted by the asterisk is redundant, *i.e.*, it is trivially satisfied in all cases and does not impose any additional conditions on primary constants. The secondary CWIs lead to

$$\alpha_2^{a_1 a_2 a_3} + (d-2)\alpha_1^{a_1 a_2 a_3} = \frac{2^{4-\frac{d}{2}} s_d \cdot \mathrm{i} f^{a_1 a_2 a_3} c_J}{\Gamma\left(\frac{d}{2} - 1\right)}, \tag{R.5.13}$$

where c_J is the 2-point function normalisation (R.1.22) and the constant s_d is defined in (R.1.3). If the regulator is present, the limit $\epsilon \to 0$ can be taken after the primary constants and triple-K integrals are substituted into the primary CWIs (R.5.9) - (R.5.10).

The 3-point function $\langle\!\langle J^{\mu_1} J^{\mu_2} J^{\mu_3} \rangle\!\rangle$ therefore depends on the 2-point function normalisation c_J and on one undetermined primary constant $\alpha_1^{a_1 a_2 a_3}$.

Examples

For $d = 3$ we find

$$A_1^{a_1 a_2 a_3} = \frac{2\alpha_1^{a_1 a_2 a_3}}{a_{123}^3}, \tag{R.5.14}$$
$$A_2^{a_1 a_2 a_3} = \alpha_1^{a_1 a_2 a_3} \frac{p_3}{a_{123}^2} - \frac{2\mathrm{i} f^{a_1 a_2 a_3} c_J}{a_{123}}. \tag{R.5.15}$$

For $d = 4$ we find

$$A_1^{a_1a_2a_3} = \alpha_1^{a_1a_2a_3} I_{4\{111\}}, \tag{R.5.16}$$

$$A_2^{a_1a_2a_3} = -\left(\alpha_1^{a_1a_2a_3} p_3 \frac{\partial}{\partial p_3} + 4\mathrm{i} f^{a_1a_2a_3} c_J\right) I^{(0)}_{2\{111\}}, \tag{R.5.17}$$

where $I^{(0)}_{2\{111\}}$ is the coefficient of ϵ^0 in the series expansion of the regulated integral $I_{2+\epsilon\{111\}}$. The integrals can be obtained from the master integral (A.6.1) via the reduction scheme in table 1, page 44.

For $d = 5$ we find

$$A_1^{a_1a_2a_3} = \frac{2\alpha_1^{a_1a_2a_3}}{a_{123}^4}\left[a_{123}^3 + a_{123}b_{123} + 3c_{123}\right], \tag{R.5.18}$$

$$A_2^{a_1a_2a_3} = \frac{\alpha_1^{a_1a_2a_3} p_3^2}{a_{123}^3}\left[p_3^2 + 3p_3 a_{12} + 2(a_{12}^2 + b_{12})\right] - \frac{2\mathrm{i} f^{a_1a_2a_3} c_J}{a_{123}^2}\left[a_{123}^3 - a_{123}b_{123} - c_{123}\right]. \tag{R.5.19}$$

$\langle T^{\mu_1\nu_1}\mathcal{O}\mathcal{O}\rangle$

Ward identities. The transverse and trace Ward identities are

$$p_1^{\nu_1}\langle\!\langle T_{\mu_1\nu_1}(\boldsymbol{p}_1)\mathcal{O}^{I_2}(\boldsymbol{p}_2)\mathcal{O}^{I_3}(\boldsymbol{p}_3)\rangle\!\rangle = = p_{3\mu_1}\langle\!\langle \mathcal{O}^{I_2}(\boldsymbol{p}_3)\mathcal{O}^{I_3}(-\boldsymbol{p}_3)\rangle\!\rangle + p_{2\mu_1}\langle\!\langle \mathcal{O}^{I_2}(\boldsymbol{p}_2)\mathcal{O}^{I_3}(-\boldsymbol{p}_2)\rangle\!\rangle, \tag{R.6.1}$$

$$\langle\!\langle T(\boldsymbol{p}_1)\mathcal{O}^{I_2}(\boldsymbol{p}_2)\mathcal{O}^{I_3}(\boldsymbol{p}_3)\rangle\!\rangle = -\Delta_3\left[\langle\!\langle \mathcal{O}^{I_2}(\boldsymbol{p}_3)\mathcal{O}^{I_3}(-\boldsymbol{p}_3)\rangle\!\rangle + \langle\!\langle \mathcal{O}^{I_2}(\boldsymbol{p}_2)\mathcal{O}^{I_3}(-\boldsymbol{p}_2)\rangle\!\rangle\right]. \tag{R.6.2}$$

Reconstruction formula. The full 3-point function can be reconstructed from the transverse-traceless part as

$$\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)\mathcal{O}^{I_2}(\boldsymbol{p}_2)\mathcal{O}^{I_3}(\boldsymbol{p}_3)\rangle\!\rangle = \langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1)\mathcal{O}^{I_2}(\boldsymbol{p}_2)\mathcal{O}^{I_3}(\boldsymbol{p}_3)\rangle\!\rangle + \left[p_2^\alpha \mathscr{T}_\alpha^{\mu_1\nu_1}(\boldsymbol{p}_1) - \frac{\Delta_3}{d-1}\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)\right]\langle\!\langle \mathcal{O}^{I_2}(\boldsymbol{p}_2)\mathcal{O}^{I_3}(-\boldsymbol{p}_2)\rangle\!\rangle + [\boldsymbol{p}_2 \leftrightarrow \boldsymbol{p}_3], \tag{R.6.3}$$

where $\mathscr{T}_\alpha^{\mu_1\nu_1}$ is defined in (R.1.20).

Decomposition of the 3-point function. The tensor decomposition of the transverse-traceless part is

$$\langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1)\mathcal{O}^{I_2}(\boldsymbol{p}_2)\mathcal{O}^{I_3}(\boldsymbol{p}_3)\rangle\!\rangle = \Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1)\cdot A_1^{I_2I_3} p_2^{\alpha_1} p_2^{\beta_1}, \tag{R.6.4}$$

where A_1 is a form factor depending on the momentum magnitudes. This form factor is symmetric under $(p_2, I_2) \leftrightarrow (p_3, I_3)$, *i.e.*,

$$A_1^{I_3I_2}(p_1, p_3, p_2) = A_1^{I_2I_3}(p_1, p_2, p_3) \tag{R.6.5}$$

and may be calculated as

$$A_1^{I_2I_3} = \text{coefficient of } p_2^{\mu_1} p_2^{\nu_1} \text{ in } \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)\mathcal{O}^{I_2}(\boldsymbol{p}_2)\mathcal{O}^{I_3}(\boldsymbol{p}_3)\rangle\!\rangle. \tag{R.6.6}$$

Primary conformal Ward identities. The primary CWIs are

$$\mathrm{K}_{ij}\, A_1^{I_2I_3} = 0, \qquad i, j = 1, 2, 3. \tag{R.6.7}$$

The solution in terms of triple-K integrals (R.1.7) is

$$A_1^{I_2I_3} = \alpha_1^{I_2I_3} J_{2\{000\}}, \tag{R.6.8}$$

where $\alpha_1^{I_2I_3}$ is a constant (note $\alpha_1^{I_3I_2} = \alpha_1^{I_2I_3}$). If the integral diverges, the regularisation (R.1.8) should be used.

Secondary conformal Ward identities. The independent secondary CWI is

$$\mathrm{L}_{2,0}\, A_1^{I_2I_3} = 2d \cdot \text{coefficient of } p_2^{\mu_1} \text{ in } p_{1\nu_1} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)\mathcal{O}^{I_2}(\boldsymbol{p}_2)\mathcal{O}^{I_3}(\boldsymbol{p}_3)\rangle\!\rangle, \tag{R.6.9}$$

where $\mathrm{L}_{s,N}$ is defined in (R.1.9). Assuming the unitarity bound for the conformal dimension of the scalar operator $\Delta_2 = \Delta_3 = \Delta \geq \frac{d}{2} - 1$ we find

$$\alpha_1^{I_2I_3} = \frac{2^{3-\frac{d}{2}} c_{\mathcal{O}} \delta^{I_2I_3}}{\Gamma\left(\frac{d}{2}\right)} \left(1 - \frac{1}{2}\delta_{\Delta,d+n+1}\right) \times C, \tag{R.6.10}$$

where

$$C = \begin{cases} \frac{1}{\pi}\sin\left(\frac{\pi}{2}(d-2\Delta)\right) & \text{if } 2\Delta \neq d+2n,\ n \in \mathbb{N}, \\ (-1)^{\Delta-\frac{d}{2}} & \text{if } 2\Delta = d+2n,\ n \in \mathbb{N} \end{cases} \tag{R.6.11}$$

and $c_{\mathcal{O}}$ is the normalisation of the 2-point function (5.64).

The 3-point function $\langle\!\langle T^{\mu_1\nu_1}\mathcal{O}\mathcal{O}\rangle\!\rangle$ is thus uniquely determined in terms of the 2-point function normalisation $c_{\mathcal{O}}$.

Examples

For $d = 3$ and $\Delta_2 = \Delta_3 = 1$ we find

$$A_1^{I_2I_3} = -2c_{\mathcal{O}}\delta^{I_2I_3}\frac{p_1 + a_{123}}{b_{23}a_{123}^2}. \tag{R.6.12}$$

For $d = 3$ and $\Delta_2 = \Delta_3 = 3$ we find

$$A_1^{I_2I_3} = 2c_{\mathcal{O}}\delta^{I_2I_3}\frac{p_1^3 + 2p_1^2a_{23} + (2p_1 + a_{23})(a_{23}^2 - b_{23})}{a_{123}^2}. \tag{R.6.13}$$

For $d = 4$ and $\Delta_2 = \Delta_3 = 2$ we find

$$A_1^{I_2I_3} = -2c_{\mathcal{O}}\delta^{I_2I_3}\left(p_1^2\frac{\partial^2}{\partial p_1^2} - p_1\frac{\partial}{\partial p_1}\right) I_{1\{000\}}, \tag{R.6.14}$$

where the $I_{1\{000\}}$ integral is given by (A.6.9).

For $d = 4$ and $\Delta_2 = \Delta_3 = 4$ we find

$$A_1^{I_2I_3} = -2c_{\mathcal{O}}\delta^{I_2I_3} I^{(0)}_{3\{222\}}, \tag{R.6.15}$$

where $I^{(0)}_{3\{222\}}$ is the coefficient of ϵ^0 in the series expansion of the regulated integral $I_{3+\epsilon\{222\}}$. The integral can be obtained from the master integral (A.6.1) via the reduction scheme in table 1, page 44.

For $d = 5$ and $\Delta_2 = \Delta_3 = 3$ we find

$$A_1^{I_2 I_3} = \tfrac{2}{3} c_{\mathcal{O}} \delta^{I_2 I_3} \frac{8p_1^2 + 9p_1 a_{23} + 3a_{23}^2}{a_{123}^3}. \tag{R.6.16}$$

For $d = 5$ and $\Delta_2 = \Delta_3 = 5$ we find

$$\begin{aligned} A_1^{I_2 I_3} = \frac{2c_{\mathcal{O}} \delta^{I_2 I_3}}{3a_{123}^3} \big[& 3p_1^6 + 9p_1^5 a_{23} + 6p_1^4 (2a_{23}^2 - b_{23}) + 6p_1^3 a_{23}(2a_{23}^2 - 3b_{23}) \\ & + 2p_1^2 (6a_{23}^4 - 15 a_{23}^2 b_{23} + 4 b_{23}^2) \\ & + 3a_{23}(3p_1 + a_{23})(a_{23}^4 - 3a_{23}^2 b_{23} + b_{23}^2) \big]. \end{aligned} \tag{R.6.17}$$

$\langle T^{\mu_1\nu_1} J^{\mu_2} \mathcal{O} \rangle$

The transverse-traceless part of $\langle\!\langle T^{\mu_1\nu_1} J^{\mu_2} \mathcal{O} \rangle\!\rangle$ vanishes. We present the detailed analysis of this case in appendix A.7.

Ward identities. The transverse and trace Ward identities are

$$p_1^{\nu_1} \langle\!\langle T_{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2 a}(\boldsymbol{p}_2) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle = p_1^{\nu_1} \langle\!\langle \frac{\delta T_{\mu_1\nu_1}}{\delta A^a_{\mu_2}}(\boldsymbol{p}_1, \boldsymbol{p}_2) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle, \tag{R.7.1}$$

$$p_{2\mu_2} \langle\!\langle T_{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2 a}(\boldsymbol{p}_2) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle = 2p_{2\mu_2} \langle\!\langle \frac{\delta J^{\mu_2 a}}{\delta g^{\mu_1\nu_1}}(\boldsymbol{p}_2, \boldsymbol{p}_1) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle, \tag{R.7.2}$$

$$\langle\!\langle T(\boldsymbol{p}_1) J^{\mu_2 a}(\boldsymbol{p}_2) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle = 0. \tag{R.7.3}$$

Reconstruction formula. The full 3-point function can be reconstructed from the transverse-traceless part as

$$\begin{aligned} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2 a}(\boldsymbol{p}_2) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle &= \langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1) j^{\mu_2 a}(\boldsymbol{p}_2) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle \\ &+ \frac{\pi^{\mu_2}_{\alpha_2}(\boldsymbol{p}_2) p_1^{\beta_1}}{p_1^2} \mathscr{T}^{\mu_1\nu_1\alpha_1}(\boldsymbol{p}_1) \langle\!\langle \frac{\delta T_{\alpha_1\beta_1}}{\delta A^a_{\alpha_2}}(\boldsymbol{p}_1, \boldsymbol{p}_2) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle \\ &+ \frac{2p_2^{\mu_2} p_{2\alpha_2}}{p_2^2} \delta^{\mu_1\alpha_1} \delta^{\nu_1\beta_1} \langle\!\langle \frac{\delta J^{\alpha_2 a}}{\delta g^{\alpha_1\beta_1}}(\boldsymbol{p}_2, \boldsymbol{p}_1) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle, \end{aligned} \tag{R.7.4}$$

where $\mathscr{T}_\alpha^{\mu_1\nu_1}$ is defined in (R.1.20).

Decomposition of the 3-point function. The tensor decomposition of the transverse-traceless part is

$$\langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1) j^{\mu_2 a}(\boldsymbol{p}_2) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle = \Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1) \pi^{\mu_2}_{\alpha_2}(\boldsymbol{p}_2) \left[A_1^{aI} p_2^{\alpha_1} p_2^{\beta_1} p_3^{\alpha_2} + A_2^{aI} \delta^{\alpha_1\alpha_2} p_2^{\beta_1} \right]. \tag{R.7.5}$$

The form factors A_1 and A_2 depend on the momentum magnitudes, and may be calculated as follows

$$A_1^{aI} = \text{coefficient of } p_2^{\mu_1} p_2^{\nu_1} p_3^{\mu_2}, \tag{R.7.6}$$

$$A_2^{aI} = 2 \cdot \text{coefficient of } \delta^{\mu_1\mu_2} p_2^{\nu_1} \tag{R.7.7}$$

in $\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2 a}(\boldsymbol{p}_2) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle$. These form factors do not exhibit any symmetry properties.

Primary conformal Ward identities. The primary CWIs are

$$\begin{aligned} \mathrm{K}_{12}\, A_1^{aI} &= 0, & \mathrm{K}_{13}\, A_1^{aI} &= 0, \\ \mathrm{K}_{12}\, A_2^{aI} &= 0, & \mathrm{K}_{13}\, A_2^{aI} &= 4A_1^{aI}. \end{aligned} \tag{R.7.8}$$

The solution in terms of triple-K integrals (R.1.7) is

$$A_1^{aI} = \alpha_1^{aI} J_{3\{000\}}, \tag{R.7.9}$$
$$A_2^{aI} = 2\alpha_1^{aI} J_{2\{001\}} + \alpha_2^{aI} J_{1\{000\}}, \tag{R.7.10}$$

where α_j^{aI}, $j = 1, 2$ are constants. If the integrals diverge, the regularisation (R.1.8) should be used.

Secondary conformal Ward identities. The independent secondary CWIs are

$$\mathrm{L}_{2,0}\, A_1^{aI} + \mathrm{R}_2\, A_2^{aI} = 2d \cdot \text{coefficient of } p_2^{\mu_1} p_3^{\mu_2} \text{ in } p_{1\nu_1} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2 a}(\boldsymbol{p}_2) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle, \tag{R.7.11}$$
$$\mathrm{L}'_{1,0}\, A_1^{aI} + 2\,\mathrm{R}'_1\, A_2^{aI} =$$
$$= -2(d-2) \cdot \text{coefficient of } p_2^{\mu_1} p_2^{\nu_1} \text{ in } p_{2\mu_2} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2 a}(\boldsymbol{p}_2) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle, \tag{R.7.12}$$
$$\mathrm{L}_{2,0}\, A_2^{aI} = 4d \cdot \text{coefficient of } \delta^{\mu_1\mu_2} \text{ in } p_{1\nu_1} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2 a}(\boldsymbol{p}_2) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle, \tag{R.7.13}$$

where the L and R operators are given by (R.1.9) and (R.1.10). This leads to the trivial solution

$$\alpha_1^{aI} = \alpha_2^{aI} = 0. \tag{R.7.14}$$

$\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3} \rangle$

Ward identities. The transverse and trace Ward identities are

$$p_1^{\nu_1} \langle\!\langle T_{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2 a_2}(\boldsymbol{p}_2) J^{\mu_3 a_3}(\boldsymbol{p}_3) \rangle\!\rangle =$$
$$= p_1^{\nu_1} \langle\!\langle \frac{\delta T_{\mu_1\nu_1}}{\delta A^{a_3}_{\mu_3}}(\boldsymbol{p}_1, \boldsymbol{p}_3) J^{\mu_2 a_2}(\boldsymbol{p}_2) \rangle\!\rangle + p_1^{\nu_1} \langle\!\langle \frac{\delta T_{\mu_1\nu_1}}{\delta A^{a_2}_{\mu_2}}(\boldsymbol{p}_1, \boldsymbol{p}_2) J^{\mu_3 a_3}(\boldsymbol{p}_3) \rangle\!\rangle$$
$$+\ 2\delta^{\mu_3[\mu_1} p_3^{\alpha]} \langle\!\langle J^{\mu_2 a_2}(\boldsymbol{p}_2) J^{a_3}_{\alpha}(-\boldsymbol{p}_2) \rangle\!\rangle + 2\delta^{\mu_2[\mu_1} p_2^{\alpha]} \langle\!\langle J^{a_2}_{\alpha}(\boldsymbol{p}_3) J^{\mu_3 a_3}(-\boldsymbol{p}_3) \rangle\!\rangle, \tag{R.8.1}$$
$$p_{2\mu_2} \langle\!\langle T_{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2 a_2}(\boldsymbol{p}_2) J^{\mu_3 a_3}(\boldsymbol{p}_3) \rangle\!\rangle =$$
$$= 2p_{2\mu_2} \langle\!\langle \frac{\delta J^{\mu_2 a_2}}{\delta g^{\mu_1\nu_1}}(\boldsymbol{p}_2, \boldsymbol{p}_1) J^{\mu_3 a_3}(\boldsymbol{p}_3) \rangle\!\rangle + \delta_{\mu_1\nu_1} p_{1\alpha} \langle\!\langle J^{\alpha a_2}(\boldsymbol{p}_3) J^{\mu_3 a_3}(-\boldsymbol{p}_3) \rangle\!\rangle, \tag{R.8.2}$$
$$\langle\!\langle T(\boldsymbol{p}_1) J^{\mu_2 a_2}(\boldsymbol{p}_2) J^{\mu_3 a_3}(\boldsymbol{p}_3) \rangle\!\rangle =$$
$$= \langle\!\langle \frac{\delta T}{\delta A^{a_2}_{\mu_2}}(\boldsymbol{p}_1, \boldsymbol{p}_2) J^{\mu_3 a_3}(\boldsymbol{p}_3) \rangle\!\rangle + \langle\!\langle \frac{\delta T}{\delta A^{a_3}_{\mu_3}}(\boldsymbol{p}_1, \boldsymbol{p}_3) J^{\mu_2 a_2}(\boldsymbol{p}_2) \rangle\!\rangle. \tag{R.8.3}$$

Reconstruction formula. The full 3-point function can be reconstructed from the transverse-traceless part as

$$\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2 a_2}(\boldsymbol{p}_2) J^{\mu_3 a_3}(\boldsymbol{p}_3) \rangle\!\rangle = \langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1) j^{\mu_2 a_2}(\boldsymbol{p}_2) j^{\mu_3 a_3}(\boldsymbol{p}_3) \rangle\!\rangle$$
$$+ \left[2\mathscr{T}_{\alpha}^{\mu_1\nu_1}(\boldsymbol{p}_1) \pi^{\mu_3[\alpha}(\boldsymbol{p}_3) p_3^{\beta]} + \frac{p_3^{\mu_3}}{p_3^2} \delta^{\mu_1\nu_1} p_{1\beta} \right] \langle\!\langle J^{\mu_2 a_2}(\boldsymbol{p}_2) J^{a_3}_{\beta}(-\boldsymbol{p}_2) \rangle\!\rangle$$
$$+ \pi^{\mu_3}_{\alpha_3}(\boldsymbol{p}_3) \left[\mathscr{T}^{\mu_1\nu_1\alpha_1}(\boldsymbol{p}_1) p_1^{\beta_1} + \frac{\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)}{d-1} \delta^{\alpha_1\beta_1} \right] \langle\!\langle \frac{\delta T_{\alpha_1\beta_1}}{\delta A^{a_3}_{\alpha_3}}(\boldsymbol{p}_1, \boldsymbol{p}_3) J^{\mu_2 a_2}(\boldsymbol{p}_2) \rangle\!\rangle$$
$$+ \frac{2p_3^{\mu_3} p_{3\alpha_3}}{p_3^2} \delta^{\mu_1\alpha_1} \delta^{\nu_1\beta_1} \langle\!\langle \frac{\delta J^{\alpha_3 a_3}}{\delta g^{\alpha_1\beta_1}}(\boldsymbol{p}_3, \boldsymbol{p}_1) J^{\mu_2 a_2}(\boldsymbol{p}_2) \rangle\!\rangle$$
$$+ \text{ everything with } (\boldsymbol{p}_2, a_2, \mu_2) \leftrightarrow (\boldsymbol{p}_3, a_3, \mu_3), \tag{R.8.4}$$

where $\mathscr{T}_{\alpha}^{\mu_1\nu_1}$ is defined in (R.1.20).

Decomposition of the 3-point function. The tensor decomposition of the transverse-traceless part is

$$\begin{aligned}
&\langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1) j^{\mu_2 a_2}(\boldsymbol{p}_2) j^{\mu_3 a_3}(\boldsymbol{p}_3)\rangle\!\rangle \\
&\quad = \Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1)\pi^{\mu_2}_{\alpha_2}(\boldsymbol{p}_2)\pi^{\mu_3}_{\alpha_3}(\boldsymbol{p}_3)\Big[A_1^{a_2a_3} p_2^{\alpha_1} p_2^{\beta_1} p_3^{\alpha_2} p_1^{\alpha_3} + A_2^{a_2a_3}\delta^{\alpha_2\alpha_3} p_2^{\alpha_1} p_2^{\beta_1} \\
&\qquad + A_3^{a_2a_3}\delta^{\alpha_1\alpha_2} p_2^{\beta_1} p_1^{\alpha_3} + A_3^{a_3a_2}(p_2 \leftrightarrow p_3)\delta^{\alpha_1\alpha_3} p_2^{\beta_1} p_3^{\alpha_2} \\
&\qquad + A_4^{a_2a_3}\delta^{\alpha_1\alpha_3}\delta^{\alpha_2\beta_1}\Big]. \qquad \text{(R.8.5)}
\end{aligned}$$

The form factors A_j, $j = 1,2,3,4$ are functions of the momentum magnitudes. If no arguments are specified then the standard ordering is assumed, $A_j = A_j(p_1,p_2,p_3)$, while by $p_i \leftrightarrow p_j$ we denote the exchange of the two momenta, *e.g.*, $A_3(p_2 \leftrightarrow p_3) = A_3(p_1,p_3,p_2)$.

The form factors A_1, A_2 and A_4 are symmetric under $(p_2,a_2) \leftrightarrow (p_3,a_3)$, *i.e.*, they satisfy,

$$A_j^{a_3a_2}(p_1,p_3,p_2) = A_j^{a_2a_3}(p_1,p_2,p_3), \qquad j \in \{1,2,4\}, \qquad \text{(R.8.6)}$$

while the form factor A_3 does not exhibit any symmetry properties.

The form factors may be determined as follows

$$\begin{aligned}
A_1^{a_2a_3} &= \text{coefficient of } p_2^{\mu_1} p_2^{\nu_1} p_3^{\mu_2} p_1^{\mu_3}, && \text{(R.8.7)}\\
A_2^{a_2a_3} &= \text{coefficient of } \delta^{\mu_2\mu_3} p_2^{\mu_1} p_2^{\nu_1}, && \text{(R.8.8)}\\
A_3^{a_2a_3} &= 2\cdot\text{coefficient of } \delta^{\mu_1\mu_2} p_2^{\nu_1} p_1^{\mu_3}, && \text{(R.8.9)}\\
A_4^{a_2a_3} &= 2\cdot\text{coefficient of } \delta^{\mu_1\mu_2}\delta^{\mu_3\nu_1}, && \text{(R.8.10)}
\end{aligned}$$

in $\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2 a_2}(\boldsymbol{p}_2) J^{\mu_3 a_3}(\boldsymbol{p}_3)\rangle\!\rangle$.

Primary conformal Ward identities. The primary CWIs are

$$\begin{aligned}
&\mathrm{K}_{12}\, A_1^{a_2a_3} = 0, && \mathrm{K}_{13}\, A_1^{a_2a_3} = 0,\\
&\mathrm{K}_{12}\, A_2^{a_2a_3} = -2A_1^{a_2a_3}, && \mathrm{K}_{13}\, A_2^{a_2a_3} = -2A_1^{a_2a_3},\\
&\mathrm{K}_{12}\, A_3^{a_2a_3} = 0, && \mathrm{K}_{13}\, A_3^{a_2a_3} = 4A_1^{a_2a_3},\\
&\mathrm{K}_{12}\, A_4^{a_2a_3} = 2A_3^{a_2a_3}, && \mathrm{K}_{13}\, A_4^{a_2a_3} = 2A_3^{a_3a_2}(p_2 \leftrightarrow p_3),
\end{aligned} \qquad \text{(R.8.11)}$$

The solution in terms of triple-K integrals (R.1.7) is

$$\begin{aligned}
A_1^{a_2a_3} &= \alpha_1^{a_2a_3} J_{4\{000\}}, && \text{(R.8.12)}\\
A_2^{a_2a_3} &= \alpha_1^{a_2a_3} J_{3\{100\}} + \alpha_2^{a_2a_3} J_{2\{000\}}, && \text{(R.8.13)}\\
A_3^{a_2a_3} &= 2\alpha_1^{a_2a_3} J_{3\{001\}} + \alpha_3^{a_2a_3} J_{2\{000\}}, && \text{(R.8.14)}\\
A_4^{a_2a_3} &= 2\alpha_1^{a_2a_3} J_{2\{011\}} + \alpha_3^{a_2a_3}\left(J_{1\{010\}} + J_{1\{001\}}\right) + \alpha_4^{a_2a_3} J_{0\{000\}}, && \text{(R.8.15)}
\end{aligned}$$

where $\alpha_j^{a_2a_3}$, $j = 1,2,3,4$ are constants. In particular all constants are symmetric in the group indices, $\alpha_j^{a_3a_2} = \alpha_j^{a_2a_3}$, $j = 1,2,3,4$. If the integrals diverge, the regularisation (R.1.8) should be used.

Secondary conformal Ward identities. The independent secondary CWIs are

$$\begin{aligned}
&(*)\ \mathrm{L}_{2,2}\, A_1^{a_2a_3} + \mathrm{R}_2\,[A_3^{a_2a_3} - A_3^{a_2a_3}(p_2 \leftrightarrow p_3)] =\\
&\qquad = 2d\cdot\text{coefficient of } p_2^{\mu_1} p_3^{\mu_2} p_1^{\mu_3} \text{ in } p_{1\nu_1}\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2 a_2}(\boldsymbol{p}_2) J^{\mu_3 a_3}(\boldsymbol{p}_3)\rangle\!\rangle, && \text{(R.8.16)}\\
&\mathrm{L}'_{1,2}\, A_1^{a_2a_3} + 2\,\mathrm{R}'_1\,[A_3^{a_2a_3} - A_2^{a_2a_3}] =\\
&\qquad = 2d\cdot\text{coefficient of } p_2^{\mu_1} p_2^{\nu_1} p_1^{\mu_3} \text{ in } p_{2\mu_2}\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2 a_2}(\boldsymbol{p}_2) J^{\mu_3 a_3}(\boldsymbol{p}_3)\rangle\!\rangle, && \text{(R.8.17)}\\
&\mathrm{L}_{2,0}\, A_2^{a_2a_3} - p_1^2\,[A_3^{a_2a_3} - A_3^{a_2a_3}(p_2 \leftrightarrow p_3)] =\\
&\qquad = 2d\cdot\text{coefficient of } \delta^{\mu_2\mu_3} p_2^{\mu_1} \text{ in } p_{1\nu_1}\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2 a_2}(\boldsymbol{p}_2) J^{\mu_3 a_3}(\boldsymbol{p}_3)\rangle\!\rangle, && \text{(R.8.18)}\\
&\mathrm{L}_{2,2}\, A_3^{a_2a_3} - 2\,\mathrm{R}_2\, A_4^{a_2a_3} =\\
&\qquad = 4d\cdot\text{coefficient of } \delta^{\mu_1\mu_2} p_1^{\mu_3} \text{ in } p_{1\nu_1}\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2 a_2}(\boldsymbol{p}_2) J^{\mu_3 a_3}(\boldsymbol{p}_3))\rangle\!\rangle, && \text{(R.8.19)}
\end{aligned}$$

where L and R are given by (R.1.9) and (R.1.10). The identity denoted by the asterisk is redundant, *i.e.*, it is trivially satisfied in all cases and does not impose any additional conditions on primary constants. The secondary CWIs lead to the following relations,

$$\alpha_3^{a_2a_3} = \alpha_2^{a_2a_3}, \tag{R.8.20}$$

$$d\alpha_1^{a_2a_3} + \alpha_2^{a_2a_3} = \frac{2^{3-\frac{d}{2}} s_d c_J \delta^{a_2a_3}}{\Gamma\left(\frac{d}{2}\right)}, \tag{R.8.21}$$

$$\alpha_4^{(0)a_2a_3} = -(d-2)\alpha_2^{a_2a_3}, \tag{R.8.22}$$

$$2d(d-2)\alpha_1^{a_2a_3} + d\alpha_3^{a_2a_3} + 2\alpha_4^{(1)a_2a_3} = \frac{2^{5-\frac{d}{2}} s_d c^{a_2a_3} c_J}{\Gamma\left(\frac{d}{2}-1\right)}, \tag{R.8.23}$$

where c_J is the normalisation of the 2-point function (R.1.22), the constant s_d is defined in (R.1.3) while the constant c^{ab} is defined as

$$\langle\!\langle \frac{\delta T_{\mu_1\nu_1}}{\delta A^a_{\mu_2}}(\boldsymbol{p}_1, \boldsymbol{p}_2) J^{\mu_3 b}(\boldsymbol{p}_3) \rangle\!\rangle = 2c^{ab}\delta^{\mu_2}_{(\mu_1} \langle\!\langle J^a_{\nu_1)}(\boldsymbol{p}_3) J^{\mu_3 b}(-\boldsymbol{p}_3) \rangle\!\rangle. \tag{R.8.24}$$

By $\alpha_4^{(n)a_2a_3}$ we mean the coefficient of ϵ^n in the series expansion of $\alpha_4^{a_2a_3}$ in the regulator ϵ. We assume that the α_1 coefficient is independent of ϵ, *i.e.*, $\alpha_1^{a_2a_3} = \alpha_1^{(0)a_2a_3}$. If the regulator is present, the limit $\epsilon \to 0$ can be taken after the primary constants and triple-K integrals are substituted into the primary CWIs (R.8.12) - (R.8.15).

The 3-point function $\langle\!\langle T^{\mu_1\nu_1} J^{\mu_2} J^{\mu_3} \rangle\!\rangle$ thus depends on the 2-point function normalisations c_J and c^{ab} and one undetermined primary constant $\alpha_1^{a_2a_3}$. The dependence of this correlator on two 2-point function normalisations rather than the one found in [3] is related to our definition of this correlator, as discussed above (7.32).

Examples

For $d = 3$ we find

$$A_1^{a_2a_3} = \alpha_1^{a_2a_3} \frac{2(4p_1 + a_{23})}{a_{123}^4}, \tag{R.8.25}$$

$$A_2^{a_2a_3} = \alpha_1^{a_2a_3} \frac{2p_1^2}{a_{123}^3} - \frac{2(2p_1 + a_{23})}{a_{123}^2} c_J \delta^{a_2a_3}, \tag{R.8.26}$$

$$A_3^{a_2a_3} = \frac{\alpha_1^{a_2a_3}}{a_{123}^3}\left[-2p_1^2 - p_2^2 + p_3^2 - 3p_1p_2 + 3p_1p_3\right] - \frac{2(2p_1 + a_{23})}{a_{123}^2} c_J \delta^{a_2a_3}, \tag{R.8.27}$$

$$\begin{aligned} A_4^{a_2a_3} = {} & \alpha_1^{a_2a_3} \frac{(2p_1 + a_{23})(p_1^2 - a_{23}^2 + 4b_{23})}{2a_{123}^2} \\ & + \left(\frac{2p_1^2}{a_{123}} - a_{23}\right) c_J \delta^{a_2a_3} + 2a_{23} c^{a_3a_2} c_J. \end{aligned} \tag{R.8.28}$$

For $d = 4$ we find

$$A_1^{a_2a_3} = \alpha_1^{a_2a_3} I_{5\{211\}}, \tag{R.8.29}$$

$$A_2^{a_2a_3} = -\left(2c_J\delta^{a_2a_3} + \alpha_1^{a_2a_3} p_1 \frac{\partial}{\partial p_1}\right) I^{(0)}_{3\{211\}}, \tag{R.8.30}$$

$$A_3^{a_2a_3} = -2\left(c_J\delta^{a_2a_3} + \alpha_1^{a_2a_3} p_3 \frac{\partial}{\partial p_3}\right) I^{(0)}_{3\{211\}}, \tag{R.8.31}$$

$$\begin{aligned} A_4^{a_2a_3} = {} & 2c_J\delta^{a_2a_3}\left[-2 + p_2\frac{\partial}{\partial p_2} + p_3\frac{\partial}{\partial p_3}\right] I^{(0)}_{1\{211\}} + 2\alpha_1^{a_2a_3} p_2p_3 \frac{\partial^2}{\partial p_2\partial p_3} I^{(0)}_{1\{211\}} \\ & + 4(c_J\delta^{a_2a_3} - c^{a_3a_2}c_J)\left[p_2^2\log p_2 + p_3^2\log p_3\right. \\ & \left. - (p_2^2+p_3^2)\left(\tfrac{1}{2} - \gamma_E + \log 2\right) - \tfrac{1}{2}p_1^2\right], \end{aligned} \tag{R.8.32}$$

where $I^{(0)}_{\alpha\{\beta_j\}}$ denotes the the coefficient of ϵ^0 in the series expansion of the regulated integral $I_{\alpha+\epsilon\{\beta_j\}}$ in ϵ. The integrals can be obtained from the master integral (A.6.1) via the reduction scheme in table 1, page 44.

In case of $d = 4$ the trace Ward identity is anomalous,

$$\langle T\rangle = \frac{\kappa}{4} F^a_{\mu\nu}F^{\mu\nu a}, \qquad\qquad \kappa = 2c_J. \tag{R.8.33}$$

It leads to the anomalous contribution,

$$\langle\!\langle T(\boldsymbol{p}_1)J^{\mu_2a_2}(\boldsymbol{p}_2)J^{\mu_3a_3}(\boldsymbol{p}_3)\rangle\!\rangle_{\text{anomaly}} = -\kappa\delta^{a_2a_3}\left[p_3^{\mu_2}p_2^{\mu_3} + \tfrac{1}{2}(p_1^2 - p_2^2 - p_3^2)\delta^{\mu_2\mu_3}\right]. \tag{R.8.34}$$

Then the anomalous contribution to the full 3-point function is

$$\begin{aligned} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)J^{\mu_2a_2}(\boldsymbol{p}_2)J^{\mu_3a_3}(\boldsymbol{p}_3)\rangle\!\rangle_{\text{anomaly}} = {} & \frac{\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)}{d-1}\pi^{\mu_2}_{\alpha_2}(\boldsymbol{p}_2)\pi^{\mu_3}_{\alpha_3}(\boldsymbol{p}_3)\delta^{a_2a_3}\times \\ \times\ \kappa & \left[-p_3^{\alpha_2}p_1^{\alpha_3} - \tfrac{1}{2}(p_1^2 - p_2^2 - p_3^2)\delta^{\alpha_2\alpha_3}\right] \end{aligned} \tag{R.8.35}$$

and should be added to the right hand side of (R.8.4).

The scheme dependence of the solution (R.8.29) - (R.8.32) due to the counterterm is

$$\begin{aligned} & A_2^{a_2a_3} \mapsto A_2^{a_2a_3} + 2\kappa_0\delta^{a_2a_3}, \\ & A_3^{a_2a_3} \mapsto A_3^{a_2a_3} + 2\kappa_0\delta^{a_2a_3}, \\ & A_4^{a_2a_3} \mapsto A_4^{a_2a_3} - \kappa_0(p_1^2 - p_2^2 - p_3^2)\delta^{a_2a_3} - 2c^{a_3a_2}\kappa_0(p_2^2 + p_3^2), \end{aligned} \tag{R.8.36}$$

where κ_0 is an arbitrary constant. See section 8.2 for details.

For $d = 5$ we find

$$\begin{aligned} A_1^{a_2a_3} = {} & \frac{2\alpha_1^{a_2a_3}}{a_{123}^5}\left[4p_1^4 + 20p_1^3a_{23} + 4p_1^2(7a_{23}^2 + 6b_{23})\right. \\ & \left. + 15p_1a_{23}(a_{23}^2 + b_{23}) + 3a_{23}^2(a_{23}^2 + b_{23})\right], \end{aligned} \tag{R.8.37}$$

$$\begin{aligned} A_2^{a_2a_3} = {} & \frac{2\alpha_1^{a_2a_3}p_1^2}{a_{123}^4}\left[a_{123}^3 + a_{123}b_{123} + 3c_{123}\right] \\ & - \frac{2c_J\delta^{a_2a_3}}{3a_{123}^3}\left[2p_1^4 + 6p_1^3a_{23} + 2p_1^2(5a_{23}^2 - 4b_{23})\right. \\ & \left. + 9p_1(a_{23}^3 - a_{23}b_{23}) + 3a_{23}^2(a_{23}^2 - b_{23})\right], \end{aligned} \tag{R.8.38}$$

$$
\begin{aligned}
A_3^{a_2a_3} = \frac{\alpha_1^{a_2a_3}}{a_{123}^4}\,&\big[-2p_1^5 - 8p_1^4(p_2+p_3) - 8p_1^3p_2(2p_2+3p_3)\\
&+\ p_1^2(-19p_2^3 - 40p_2^2p_3 + 24p_2p_3^2 + 15p_3^3)\\
&- 3(4p_1+p_2+p_3)(p_2^2-p_3^2)(p_2^2+3p_2p_3+p_3^2)\big]\\
-\ \frac{2c_J\delta^{a_2a_3}}{3a_{123}^3}\,&\big[2p_1^4 + 6p_1^3a_{23} + 2p_1^2(5a_{23}^2 - 4b_{23})\\
&+ 9p_1a_{23}(a_{23}^2 - b_{23}) + 3a_{23}^2(a_{23}^2 - b_{23})\big]\,,
\end{aligned}
\tag{R.8.39}
$$

$$
\begin{aligned}
A_4^{a_2a_3} = \frac{\alpha_1^{a_2a_3}}{2a_{123}^3}\,&\big[2p_1^6 + 6p_1^5a_{23} + 4p_1^4(2a_{23}^2 - b_{23}) + p_1^2(a_{23}^2+b_{23})(3p_1a_{23} - 7a_{23}^2 + 32b_{23})\\
&- 3a_{23}(3p_1+a_{23})(a_{23}^2 - 4b_{23})(a_{23}^2+b_{23})\big]\\
+\ \frac{c_J\delta^{a_2a_3}}{3a_{123}^2}\,&\big[2p_1^5 + 4p_1^4a_{23} + 4p_1^3(a_{23}^2 - b_{23}) - p_1^2a_{23}(a_{23}^2 - 7b_{23})\\
&- 6p_1a_{23}^2(a_{23}^2 - 3b_{23}) - 3a_{23}^3(a_{23}^2 - 3b_{23})\big]\\
+\ 6(p_2^3+p_3^3)&c^{a_3a_2}c_J.
\end{aligned}
\tag{R.8.40}
$$

$\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}\mathcal{O}\rangle$

Ward identities. The transverse and trace Ward identities are

$$
p_1^{\nu_1}\langle\!\langle T_{\mu_1\nu_1}(\boldsymbol{p}_1)T_{\mu_2\nu_2}(\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle = 2p_1^{\nu_1}\langle\!\langle\frac{\delta T_{\mu_1\nu_1}}{\delta g^{\mu_2\nu_2}}(\boldsymbol{p}_1,\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle,
\tag{R.9.1}
$$

$$
\langle\!\langle T(\boldsymbol{p}_1)T_{\mu_2\nu_2}(\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle = 2\langle\!\langle\frac{\delta T}{\delta g^{\mu_2\nu_2}}(\boldsymbol{p}_1,\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle.
\tag{R.9.2}
$$

Reconstruction formula. The full 3-point function can be reconstructed from the transverse-traceless part as

$$
\begin{aligned}
\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle &= \langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1)t^{\mu_2\nu_2}(\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle\\
&+ 2\left[\mathscr{T}^{\mu_1\nu_1\alpha_1}(\boldsymbol{p}_1)p_1^{\beta_1} + \frac{\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)}{d-1}\delta^{\alpha_1\beta_1}\right]\delta^{\mu_2\alpha_2}\delta^{\nu_2\beta_2}\langle\!\langle\frac{\delta T_{\alpha_1\beta_1}}{\delta g^{\alpha_2\beta_2}}(\boldsymbol{p}_1,\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle\\
&+ 2[(\mu_1,\nu_1,\boldsymbol{p}_1)\leftrightarrow(\mu_2,\nu_2,\boldsymbol{p}_2)]\\
&- 4\left[\mathscr{T}^{\mu_1\nu_1\alpha_1}(\boldsymbol{p}_1)p_1^{\beta_1} + \frac{\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)}{d-1}\delta^{\alpha_1\beta_1}\right]\left[\mathscr{T}^{\mu_2\nu_2\alpha_2}(\boldsymbol{p}_2)p_2^{\beta_2} + \frac{\pi^{\mu_2\nu_2}(\boldsymbol{p}_2)}{d-1}\delta^{\alpha_2\beta_2}\right]\times\\
&\qquad\times\ \langle\!\langle\frac{\delta T_{\alpha_1\beta_1}}{\delta g^{\alpha_2\beta_2}}(\boldsymbol{p}_1,\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle,
\end{aligned}
\tag{R.9.3}
$$

where $\mathscr{T}_\alpha^{\mu\nu}$ is defined in (R.1.20).

Decomposition of the 3-point function. The tensor decomposition of the transverse-traceless part is

$$
\begin{aligned}
\langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1)t^{\mu_2\nu_2}(\boldsymbol{p}_2)\mathcal{O}^I(p_3)\rangle\!\rangle = \Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1)\Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)\,&\Big[A_1^I p_2^{\alpha_1}p_2^{\beta_1}p_3^{\alpha_2}p_3^{\beta_2}\\
&+ A_2^I\delta^{\alpha_1\alpha_2}p_2^{\beta_1}p_3^{\beta_2} + A_3^I\delta^{\alpha_1\alpha_2}\delta^{\beta_1\beta_2}\Big].
\end{aligned}
\tag{R.9.4}
$$

The form factors A_j, $j=1,2,3$ are functions of the momentum magnitudes. All form factors are symmetric under $p_1\leftrightarrow p_2$, *i.e.*, they satisfy

$$
A_j^I(p_2,p_1,p_3) = A_j^I(p_1,p_2,p_3), \qquad j=1,2,3.
\tag{R.9.5}
$$

These form factors may be calculated using

$$A_1^I = \text{coefficient of } p_2^{\mu_1} p_2^{\nu_1} p_3^{\mu_2} p_3^{\nu_2}, \tag{R.9.6}$$
$$A_2^I = 4 \cdot \text{coefficient of } \delta^{\mu_1\mu_2} p_2^{\nu_1} p_3^{\nu_2}, \tag{R.9.7}$$
$$A_3^I = 2 \cdot \text{coefficient of } \delta^{\mu_1\mu_2} \delta^{\nu_1\nu_2}, \tag{R.9.8}$$

in $\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) T^{\mu_2\nu_2}(\boldsymbol{p}_2) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle$.

Primary conformal Ward identities. The primary CWIs are

$$\begin{array}{ll} \mathrm{K}_{12}\, A_1^I = 0, & \mathrm{K}_{13}\, A_1^I = 0, \\ \mathrm{K}_{12}\, A_2^I = 0, & \mathrm{K}_{13}\, A_2^I = 8A_1^I, \\ \mathrm{K}_{12}\, A_3^I = 0, & \mathrm{K}_{13}\, A_3^I = 2A_2^I, \end{array} \tag{R.9.9}$$

The solution in terms of triple-K integrals (R.1.7) is

$$A_1^I = \alpha_1^I J_{4\{000\}}, \tag{R.9.10}$$
$$A_2^I = 4\alpha_1^I J_{3\{001\}} + \alpha_2^I J_{2\{000\}}, \tag{R.9.11}$$
$$A_3^I = 2\alpha_1^I J_{2\{002\}} + \alpha_2^I J_{1\{001\}} + \alpha_3^I J_{0\{000\}}, \tag{R.9.12}$$

where α_j^I, $j = 1, 2, 3$ are constants. If the integrals diverge, the regularisation (R.1.8) should be used.

Secondary conformal Ward identities. The independent secondary CWIs are

$$\begin{aligned} &\mathrm{L}_{2,0}\, A_1^I + \mathrm{R}_2\, A_2^I = \\ &\quad = 2d \cdot \text{coefficient of } p_2^{\mu_1} p_3^{\mu_2} p_3^{\nu_2} \text{ in } p_{1\nu_1} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) T^{\mu_2\nu_2}(\boldsymbol{p}_2) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle, \end{aligned} \tag{R.9.13}$$
$$\begin{aligned} &\mathrm{L}_{2,0}\, A_2^I + 4\,\mathrm{R}_2\, A_3^I = \\ &\quad = 8d \cdot \text{coefficient of } \delta^{\mu_1\mu_2} p_3^{\mu_2} \text{ in } p_{1\nu_1} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) T^{\mu_2\nu_2}(\boldsymbol{p}_2) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle. \end{aligned} \tag{R.9.14}$$

where L and R are defined in (R.1.9) and (R.1.10). They lead to the following relations

$$\alpha_2^{(0)I} = (\Delta_3 + 2)(d - \Delta_3 - 2)\alpha_1^I, \tag{R.9.15}$$
$$\alpha_2^{(1)I} = \begin{cases} (2\Delta_3 - d - 4)\alpha_1^I & \text{if } \Delta_3 \neq d - 2 - 2n,\ n \in \mathbb{N}, \\ -(2\Delta_3 - d + 4)\alpha_1^I + C_1 c_1^I c_{\mathcal{O}} & \text{if } \Delta_3 \neq d - 2 - 2n,\ n \in \mathbb{N}, \end{cases} \tag{R.9.16}$$
$$\alpha_2^{(2)I} = -\alpha_1^I, \tag{R.9.17}$$
$$\alpha_3^{(0)I} = \frac{1}{4}\Delta_3(\Delta_3 + 2)(d - \Delta_3)(d - \Delta_3 - 2)\alpha_1^I, \tag{R.9.18}$$
$$\alpha_3^{(1)I} = \begin{cases} -\frac{1}{2}(6d + d^2 - 12\Delta_3 - 2d\Delta_3 + d^2\Delta_3 + 2\Delta_3^2 - 3d\Delta_3^2 + 2\Delta_3^3)\alpha_1^I \\ \qquad \text{if } \Delta_3 \neq d - 2n,\ n \in \mathbb{N}, \\ \frac{1}{4}\left[(2\Delta_3 - d)(\Delta_3 - d + 2)(\Delta_3 + 2)\alpha_1^I - \Delta_3(\Delta_3 - d)\alpha_2^{(1)}\right] + C_2 c_2^I c_{\mathcal{O}} \\ \qquad \text{if } \Delta_3 = d - 2n,\ n \in \mathbb{N}, \end{cases} \tag{R.9.19}$$
$$\alpha_3^{(2)I} = \frac{1}{4}(4 + 2d + d^2 - 4\Delta_3 - 6d\Delta_3 + 6\Delta_3^2)\alpha_1^I. \tag{R.9.20}$$

Here, $c_{\mathcal{O}}$ denotes the normalisation of the 2-point function (5.64) and the constants C_1 and C_2

are

$$C_1 = \frac{(-1)^{\frac{d-\Delta_3}{2}} 2^{4+\Delta_3-\frac{d}{2}} d\,\Gamma\left(\frac{d-\Delta_3}{2}\right)\Gamma\left(\frac{\Delta_3+3}{2}\right)}{\sqrt{\pi}\Gamma\left(\frac{\Delta_3+2}{2}\right)\Gamma\left(\frac{\Delta_3+d+4}{2}\right)}\times$$
$$\times\begin{cases}\frac{1}{\Gamma\left(\frac{d-2\Delta_3}{2}\right)} & \text{if } 2\Delta_3 \neq d+2n,\ n\in\mathbb{N},\\ -\Gamma\left(\Delta_3-\frac{d}{2}+1\right) & \text{if } 2\Delta_3 = d+2n,\ n\in\mathbb{N},\end{cases} \tag{R.9.21}$$

$$C_2 = \frac{(-1)^{\frac{d-\Delta_3+2}{2}} 2^{4+\Delta_3-\frac{d}{2}} d\,\Gamma\left(\frac{d-\Delta_3+2}{2}\right)\Gamma\left(\frac{\Delta_3+1}{2}\right)}{\sqrt{\pi}\Gamma\left(\frac{\Delta_3}{2}\right)\Gamma\left(\frac{\Delta_3+d+2}{2}\right)}\times$$
$$\times\begin{cases}\frac{1}{\Gamma\left(\frac{d-2\Delta_3}{2}\right)} & \text{if } 2\Delta_3 \neq d+2n,\ n\in\mathbb{N},\\ -\Gamma\left(\Delta_3-\frac{d}{2}+1\right) & \text{if } 2\Delta_3 = d+2n,\ n\in\mathbb{N}.\end{cases} \tag{R.9.22}$$

Moreover the c_1^K and c_2^K constants are defined as

$$p_1^{\nu_1}\langle\!\langle \frac{\delta T_{\mu_1\nu_1}}{\delta g^{\mu_2\nu_2}}(\boldsymbol{p}_1,\boldsymbol{p}_2)\mathcal{O}^K(\boldsymbol{p})_3\rangle\!\rangle\Big|_{p_1=p_2=p} = c_1^K p_2^{\mu_1} p_3^{\mu_2} p_3^{\nu_2} p^{d-\Delta_3-2}\langle\!\langle \mathcal{O}^I(\boldsymbol{p}_3)\mathcal{O}^K(-\boldsymbol{p}_3)\rangle\!\rangle$$
$$+\ c_2^K \delta^{\mu_1\mu_2} p_3^{\nu_2} p^{d-\Delta_3}\langle\!\langle \mathcal{O}^I(\boldsymbol{p}_3)\mathcal{O}^K(-\boldsymbol{p}_3)\rangle\!\rangle + \dots \tag{R.9.23}$$

Here, we mean the following: first, write down the most general tensor decomposition of $p_1^{\nu_1}\langle\!\langle \frac{\delta T_{\mu_1\nu_1}}{\delta g^{\mu_2\nu_2}}(\boldsymbol{p}_1,\boldsymbol{p}_2)\mathcal{O}^K(\boldsymbol{p})_3\rangle\!\rangle$ and take the coefficient of $p_2^{\mu_1}p_3^{\mu_2}p_3^{\nu_2}$. We then set $p_1 = p_2 = p$ in the expression for this coefficient.

The primary constants are function of the regulator ϵ and by $\alpha_j^{(n)}$ we denote the coefficient of ϵ^n in the series expansion of α_j in ϵ. We assume that the constant α_1 does not depend on the regulator. If the regulator is present, the limit $\epsilon \to 0$ can be taken after the primary constants and triple-K integrals are substituted into the primary CWIs (R.9.10) - (R.9.12).

The 3-point function $\langle\!\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}\mathcal{O}\rangle\!\rangle$ thus depends on the 2-point function normalisations $c_\mathcal{O}$, c_1^K and c_2^K and one undetermined primary constant α_1^I.

Examples

For $d = 3$ and $\Delta_3 = 1$ we find

$$A_1^I = \frac{\alpha_1^I}{p_3 a_{123}^4}\left[p_3^2 + 4p_3 a_{12} + 3(a_{12}^2 + 2b_{12})\right], \tag{R.9.24}$$

$$A_2^I = \frac{\alpha_1^I}{p_3 a_{123}^3}\left[p_3^3 + 3p_2^2 a_{12} + p_3(-a_{12}^2 + 8b_{12}) - 3a_{12}^3\right] - \frac{4c_1^I c_\mathcal{O}}{p_3}, \tag{R.9.25}$$

$$A_3^I = \frac{\alpha_1^I(a_{12}-p_3)}{4p_3 a_{123}^2}\left[-p_3^3 - 3p_3^2 a_{12} + p_3(a_{12}^2 - 10b_{12}) + 3a_{12}(a_{12}^2 - 2b_{12})\right]$$
$$+\frac{c_\mathcal{O}}{p_3}\left[(c_1^I - 3c_2^I)(p_1^2 + p_2^2) + 3(c_1^I + c_2^I)p_3^2\right]. \tag{R.9.26}$$

For $d = 3$ and $\Delta_3 = 3$ we find

$$A_1^I = \frac{2\alpha_1^I}{a_{123}^4}\left[a_{123}^3 + a_{123}b_{123} + 3c_{123}\right], \tag{R.9.27}$$

$$A_2^I = \frac{2\alpha_1^I}{a_{123}^3}\left[p_3^4 + 3p_3^3 a_{12} + 6p_3^2 b_{12} + 3p_3 a_{12}(-a_{12}^2 + b_{12}) + a_{12}^2(b_{12} - a_{12}^2)\right], \tag{R.9.28}$$

$$\begin{aligned} A_3^I = \frac{\alpha_1^I}{2a_{123}^2}\big[&p_3^5 + 2p_3^4 a_{12} + p_3^3(-3a_{12}^2 + 8b_{12}) - p_3^2(a_{12}^3 + 5a_{12}b_{12}) \\ &+ 6p_3(a_{12}^4 - 3a_{12}^2 b_{12}) + 3a_{12}^3(a_{12}^2 - 3b_{12})\big] \\ &- 4c_2^I c_{\mathcal{O}}(p_1^3 + p_2^3 + p_3^3). \end{aligned} \tag{R.9.29}$$

For $d = 4$ and $\Delta_3 = 2$ we find

$$A_1^I = \alpha_1^I I_{5\{220\}}, \tag{R.9.30}$$

$$A_2^I = 4\alpha_1^I I_{4\{221\}}^{(0)} + 8(2\alpha_1^I - c_1^I c_{\mathcal{O}})(\log p_3 + \gamma_E - \log 2), \tag{R.9.31}$$

$$\begin{aligned} A_3^I = 2\alpha_1^I I_{3\{222\}}^{(0)} &+ 2(-2\alpha_1^I + c_1^I c_{\mathcal{O}})\left[(p_1^2 + p_2^2 + p_3^2)\log p_3\right. \\ &\left. - (p_1^2 + p_2^2 + p_3^2)(1 - \gamma_E + \log 2) + p_3^2\right] \\ &+ \tfrac{16}{3} c_2^I c_{\mathcal{O}}\left[(p_3^2 - p_1^2 - p_2^2)\log p_3 + (p_3^2 - p_1^2 - p_2^2)(\gamma_E - \log 2) - p_3^2\right], \end{aligned} \tag{R.9.32}$$

where $I_{\alpha\{\beta_j\}}^{(0)}$ is the coefficient of ϵ^0 in the series expansion of the regulated integral $I_{\alpha+\epsilon\{\beta_j\}}$. The integral can be obtained from the master integral (A.6.1) via the reduction scheme in table 1, page 44.

For $d = 4$ and $\Delta_3 = 4$ we find

$$A_1^I = \alpha_1^I I_{5\{222\}}^{(0)}, \tag{R.9.33}$$

$$A_2^I = 4\alpha_1^I \left(1 - p_3 \frac{\partial}{\partial p_3}\right) I_{3\{222\}}^{(0)}, \tag{R.9.34}$$

$$\begin{aligned} A_3^I = -2\alpha_1^I p_3 \frac{\partial}{\partial p_3} I_{2\{223\}}^{(0)} &- 2(3a_1 + 4c_2^I c_{\mathcal{O}})\left[\left(p_1^4 \log p_1 - p_2^2 p_3^2 - p_1^4(\tfrac{3}{4} - \gamma_E + \log 2)\right)\right. \\ &\left. + (p_1 \leftrightarrow p_2) + (p_1 \leftrightarrow p_3)\right], \end{aligned} \tag{R.9.35}$$

where $I_{\alpha\{\beta_j\}}^{(0)}$ is the coefficient of ϵ^0 in the series expansion of the regulated integral $I_{\alpha+\epsilon\{\beta_j\}}$. The integral can be obtained from the master integral (A.6.1) via the reduction scheme in table 1, page 44.

For $d = 5$ and $\Delta_3 = 3$ we find

$$\begin{aligned}
A_1^I &= \frac{3\alpha_1^I}{a_{123}^5}\left[3p_3^4 + 15p_3^3 a_{12} + p_3^2(29a_{12}^2 + 2b_{12}) + 5p_3 a_{12}(5a_{12}^2 + 2b_{12})\right. \\
&\qquad \left. + 8(a_{12}^4 + a_{12}^2 b_{12} + b_{12}^2)\right], \qquad \text{(R.9.36)} \\
A_2^I &= \frac{\alpha_1^I}{a_{123}^4}\left[9p_3^5 + 36p_3^4 a_{12} + 6p_3^3(7a_{12}^2 + 4b_{12}) - 12p_3^2 a_{12}(a_{12}^2 - 8b_{12})\right. \\
&\qquad \left. + p_3(-51a_{12}^4 + 96a_{12}^2 b_{12} + 32b_{12}^2) + 8a_{12}(-3a_{12}^4 + 3a_{12}^2 b_{12} + b_{12}^2)\right] \\
&\quad - 4p_3 c_1^I c_{\mathcal{O}}, \qquad \text{(R.9.37)} \\
A_3^I &= \frac{\alpha_1^I}{4a_{123}^3}\left[9p_3^6 + 27p_3^5 a_{12} + 6p_3^4(a_{12}^2 + 7b_{12}) + 18p_3^3 a_{12}(-3a_{12}^2 + 7b_{12})\right. \\
&\qquad + p_3^2(-39a_{12}^4 + 30a_{12}^2 b_{12} + 64b_{12}^2) + 9p_3 a_{12}(a_{12}^2 - 4b_{12})(3a_{12}^2 - 2b_{12}) \\
&\qquad \left. + 24a_{12}^2(a_{12}^4 - 3a_{12}^2 b_{12} + b_{12}^2)\right] \\
&\quad + \frac{p_3 c_{\mathcal{O}}}{6}\left[3(2c_1^I - 5c_2^I)(p_1^2 + p_2^2) + (2c_1^I + 15c_2^I)p_3^2\right]. \qquad \text{(R.9.38)}
\end{aligned}$$

For $d = 5$ and $\Delta_3 = 5$ we find

$$\begin{aligned}
A_1^I &= \frac{6\alpha_1^I}{a_{123}^5}\left[-3a_{123}^6 + 3a_{123}^4 b_{123} + a_{123}^2 b_{123}^2 + a_{123}^3 c_{123} + 3a_{123} b_{123} c_{123} + 4c_{123}^2\right], \qquad \text{(R.9.39)} \\
A_2^I &= \frac{2\alpha_1^I}{a_{123}^4}\left[-9p_3^7 - 36p_3^6 a_{12} - 3p_3^5(17a_{12}^2 + 2b_{12}) - 24p_3^4 a_{12}(a_{12}^2 + b_{12})\right. \\
&\qquad + 8p_3^3(3a_{12}^4 - 12a_{12}^2 b_{12} + 5b_{12}^2) + p_3^2(51a_{12}^5 - 159a_{12}^3 b_{12} + 55a_{12} b_{12}^2) \\
&\qquad \left. + 9a_{12}^2(a_{12} + 4p_3)(a_{12}^4 - 3a_{12}^2 b_{12} + b_{12}^2)\right], \qquad \text{(R.9.40)} \\
A_3^I &= \frac{\alpha_1^I}{2a_{123}^3}\left[-9p_3^8 - 27p_3^7 a_{12} - 3p_3^6(5a_{12}^2 + 8b_{12}) + 9p_3^5(3a_{12}^3 - 8a_{12} b_{12})\right. \\
&\qquad + 8p_3^4(6a_{12}^4 - 15a_{12}^2 b_{12} + 4b_{12}^2) + 9p_3^3(a_{12}^5 + 3a_{12}^3 b_{12} - 11a_{12} b_{12}^2) \\
&\qquad \left. + p_3^2(-69a_{12}^6 + 369a_{12}^4 b_{12} - 393a_{12}^2 b_{12}^2) - 27a_{12}^3(a_{12} + 3p_3)(a_{12}^4 - 5a_{12}^2 b_{12} + 5b_{12}^2)\right] \\
&\quad - 4c_2^I c_{\mathcal{O}}(p_1^5 + p_2^5 + p_3^5). \qquad \text{(R.9.41)}
\end{aligned}$$

$\left\langle T^{\mu_1\nu_1} T^{\mu_2\nu_2} J^{\mu_3}\right\rangle$

This correlation function is at most semi-local, as was proved in [6] through a position space analysis. Our result confirms the triviality of this correlator through independent calculations in momentum space. In appendix A.7 we discuss the triviality of $\langle\langle T^{\mu_1\nu_1} J^{\mu_2}\mathcal{O}\rangle\rangle$, which is very similar to $\langle\langle T^{\mu_1\nu_1} T^{\mu_2\nu_2} J^{\mu_3}\rangle\rangle$.

Ward identities. The transverse and trace Ward identities are

$$\begin{aligned}
&p_1^{\nu_1}\langle\langle T_{\mu_1\nu_1}(\boldsymbol{p}_1) T_{\mu_2\nu_2}(\boldsymbol{p}_2) J^{\mu_3 a}(\boldsymbol{p}_3)\rangle\rangle = 2p_1^{\nu_1}\langle\langle \frac{\delta T_{\mu_1\nu_1}}{\delta g^{\mu_2\nu_2}}(\boldsymbol{p}_1, \boldsymbol{p}_2) J^{\mu_3 a}(\boldsymbol{p}_3)\rangle\rangle, \qquad \text{(R.10.1)} \\
&p_{3\mu_3}\langle\langle T_{\mu_1\nu_1}(\boldsymbol{p}_1) T_{\mu_2\nu_2}(\boldsymbol{p}_2) J^{\mu_3 a}(\boldsymbol{p}_3)\rangle\rangle = 0, \qquad \text{(R.10.2)} \\
&\langle\langle T(\boldsymbol{p}_1) T_{\mu_2\nu_2}(\boldsymbol{p}_2) J^{\mu_3 a}(\boldsymbol{p}_3)\rangle\rangle = 2\langle\langle \frac{\delta T}{\delta g^{\mu_2\nu_2}}(\boldsymbol{p}_1, \boldsymbol{p}_2) J^{\mu_3 a}(\boldsymbol{p}_3)\rangle\rangle. \qquad \text{(R.10.3)}
\end{aligned}$$

Reconstruction formula. The full 3-point function can be reconstructed from the transverse-

traceless part as

$$\begin{aligned}\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)J^{\mu_3 a}(\boldsymbol{p}_3)\rangle\!\rangle &= \langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1)t^{\mu_2\nu_2}(\boldsymbol{p}_2)j^{\mu_3 a}(\boldsymbol{p}_3)\rangle\!\rangle \\ &+ 2\left[\mathscr{T}^{\mu_1\nu_1\alpha_1}(\boldsymbol{p}_1)p_1^{\beta_1} + \frac{\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)}{d-1}\delta_{\alpha_1\beta_1}\right]\delta^{\mu_2\alpha_2}\delta^{\nu_2\beta_2}\langle\!\langle\frac{\delta T_{\alpha_1\beta_1}}{\delta g^{\alpha_2\beta_2}}(\boldsymbol{p}_1,\boldsymbol{p}_2)J^{\mu_3 a}(\boldsymbol{p}_3)\rangle\!\rangle \\ &+ 2[(\mu_1,\nu_1,\boldsymbol{p}_1)\leftrightarrow(\mu_2,\nu_2,\boldsymbol{p}_2)] \\ &- 4\left[\mathscr{T}^{\mu_1\nu_1\alpha_1}(\boldsymbol{p}_1)p_1^{\beta_1} + \frac{\pi^{\mu_1\nu_2}(\boldsymbol{p}_1)}{d-1}\delta_{\alpha_1\beta_1}\right]\left[\mathscr{T}^{\mu_2\nu_2\alpha_2}(\boldsymbol{p}_2)p_2^{\beta_2} + \frac{\pi^{\mu_2\nu_2}(\boldsymbol{p}_2)}{d-1}\delta_{\alpha_2\beta_2}\right]\times \\ &\qquad\times\ \langle\!\langle\frac{\delta T_{\alpha_1\beta_1}}{\delta g^{\alpha_2\beta_2}}(\boldsymbol{p}_1,\boldsymbol{p}_2)J^{\mu_3 a}(\boldsymbol{p}_3)\rangle\!\rangle, \end{aligned} \tag{R.10.4}$$

where $\mathscr{T}^{\mu\nu\alpha}$ is defined in (R.1.20).

Decomposition of the 3-point function. The tensor decomposition of the transverse-traceless part is

$$\begin{aligned}\langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1)t^{\mu_2\nu_2}(\boldsymbol{p}_2)j^{\mu_3 a}(\boldsymbol{p}_3)\rangle\!\rangle &= \Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1)\Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)\pi^{\mu_3}_{\alpha_3}(\boldsymbol{p}_3)\Big[A_1^a p_2^{\alpha_1}p_2^{\beta_1}p_3^{\alpha_2}p_3^{\beta_2}p_1^{\alpha_3} \\ &+ A_2^a\delta^{\beta_1\beta_2}p_2^{\alpha_1}p_3^{\alpha_2}p_1^{\alpha_3} \\ &+ A_3^a\delta^{\alpha_1\alpha_3}p_2^{\beta_1}p_3^{\alpha_2}p_3^{\beta_2} - A_3^a(p_1\leftrightarrow p_2)\delta^{\alpha_2\alpha_3}p_2^{\alpha_1}p_2^{\beta_1}p_3^{\beta_2} \\ &+ A_4^a\delta^{\alpha_1\alpha_2}\delta^{\beta_1\beta_2}p_1^{\alpha_3} \\ &+ A_5^a\delta^{\alpha_1\alpha_2}\delta^{\alpha_3\beta_2}p_2^{\beta_1} - A_5(p_1\leftrightarrow p_2)\delta^{\alpha_1\alpha_2}\delta^{\alpha_3\beta_1}p_3^{\beta_2}\Big]. \end{aligned} \tag{R.10.5}$$

The form factors A_j, $j=1,\ldots,5$ are functions of the momentum magnitudes. If no arguments are specified then the standard ordering is assumed, $A_j = A_j(p_1,p_2,p_3)$, while by $p_i\leftrightarrow p_j$ we denote the exchange of the two momenta, *e.g.*, $A_3(p_1\leftrightarrow p_2) = A_3(p_2,p_1,p_3)$.

The form factors A_1, A_2 and A_4 are antisymmetric under $p_1\leftrightarrow p_2$, *i.e.*, they satisfy

$$A_j^a(p_2,p_1,p_3) = -A_j^a(p_1,p_2,p_3), \qquad j\in\{1,2,4\}. \tag{R.10.6}$$

The remaining form factors A_3 and A_5 do not exhibit any symmetry properties.

The form factors can be calculated as follows

$$A_1^a = \text{coefficient of } p_2^{\mu_1}p_2^{\nu_1}p_3^{\mu_2}p_3^{\nu_2}p_1^{\mu_3}, \tag{R.10.7}$$
$$A_2^a = 4\cdot\text{coefficient of } \delta^{\nu_1\nu_2}p_2^{\mu_1}p_3^{\mu_2}p_1^{\mu_3}, \tag{R.10.8}$$
$$A_3^a = 2\cdot\text{coefficient of } \delta^{\mu_1\mu_3}p_2^{\nu_1}p_3^{\mu_2}p_3^{\nu_2}, \tag{R.10.9}$$
$$A_4^a = 2\cdot\text{coefficient of } \delta^{\mu_1\mu_2}\delta^{\nu_1\nu_2}p_1^{\mu_3}, \tag{R.10.10}$$
$$A_5^a = 4\cdot\text{coefficient of } \delta^{\mu_1\mu_2}\delta^{\mu_3\nu_2}p_2^{\nu_1} \tag{R.10.11}$$

in $\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)J^{\mu_3 a}(\boldsymbol{p}_3)\rangle\!\rangle$.

Primary conformal Ward identities. The primary CWIs are

$$\begin{aligned}&\mathrm{K}_{12}\,A_1^a = 0, && \mathrm{K}_{13}\,A_1^a = 0,\\ &\mathrm{K}_{12}\,A_2^a = 0, && \mathrm{K}_{13}\,A_2^a = 2A_1^a,\\ &\mathrm{K}_{12}\,A_3^a = 2A_1^a, && \mathrm{K}_{13}\,A_3^a = 0,\\ &\mathrm{K}_{12}\,A_4^a = 0, && \mathrm{K}_{13}\,A_4^a = 4A_2^a,\\ &\mathrm{K}_{12}\,A_5^a = -2A_2^a, && \mathrm{K}_{13}\,A_5^a = -2\left[A_2^a + A_3^a(p_1\leftrightarrow p_2)\right].\end{aligned} \tag{R.10.12}$$

The solution in terms of triple-K integrals (R.1.7) is

$$A_1^a = \alpha_1^a J_{5\{000\}}, \tag{R.10.13}$$
$$A_2^a = \alpha_1^a J_{4\{001\}} + \alpha_2^a J_{3\{000\}}, \tag{R.10.14}$$
$$A_3^a = \alpha_1^a J_{4\{010\}} + \alpha_3^a J_{3\{000\}}, \tag{R.10.15}$$
$$A_4^a = \alpha_1^a J_{3\{002\}} + 2\alpha_2^a J_{2\{001\}} + \alpha_4^a I_{1\{000\}}, \tag{R.10.16}$$
$$A_5^a = \alpha_1^a J_{3\{101\}} + \alpha_2^a J_{2\{100\}} + \alpha_3^a I_{2\{001\}} + \alpha_5^a J_{1\{000\}}, \tag{R.10.17}$$

where α_j^a, $j = 1, \ldots, 5$ are constants. If the integrals diverge, the regularisation (R.1.8) should be used.

Secondary conformal Ward identities. The independent secondary CWIs are

$$\begin{aligned}
&\mathrm{L}_{2,2}\, A_1^a + \mathrm{R}_2\,[A_2^a - A_3^a] = \\
&\quad = 2d \cdot \text{coefficient of } p_2^{\mu_1} p_3^{\mu_2} p_3^{\nu_2} p_1^{\mu_3} \text{ in } p_{1\nu_1} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) T^{\mu_2\nu_2}(\boldsymbol{p}_2) J^{\mu_3 a}(\boldsymbol{p}_3) \rangle\!\rangle, \qquad \text{(R.10.18)} \\
&\mathrm{L}_{2,2}\, A_2^a + 2\,\mathrm{R}_2\,[2A_4^a + A_5^a(p_1 \leftrightarrow p_2)] = \\
&\quad = 4d \cdot \text{coefficient of } \delta^{\mu_1\mu_3} p_3^{\mu_2} p_3^{\nu_2} \text{ in } p_{1\nu_1} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) T^{\mu_2\nu_2}(\boldsymbol{p}_2) J^{\mu_3 a}(\boldsymbol{p}_3) \rangle\!\rangle, \qquad \text{(R.10.19)} \\
&\mathrm{L}_{2,-2}\, A_3^a - 2\,\mathrm{R}_2\,[A_5^a(p_1 \leftrightarrow p_2)] = \\
&\quad = 8d \cdot \text{coefficient of } \delta^{\mu_1\mu_2} p_3^{\nu_2} p_1^{\mu_3} \text{ in } p_{1\nu_1} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) T^{\mu_2\nu_2}(\boldsymbol{p}_2) J^{\mu_3 a}(\boldsymbol{p}_3) \rangle\!\rangle, \qquad \text{(R.10.20)} \\
&\mathrm{L}_{2,0}\, A_5^a - 2p_1^2\,[2A_4^a + A_5^a(p_1 \leftrightarrow p_2)] = \\
&\quad = 8d \cdot \text{coefficient of } \delta^{\mu_2\mu_3}\delta^{\mu_1\nu_2} \text{ in } p_{1\nu_1} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) T^{\mu_2\nu_2}(\boldsymbol{p}_2) J^{\mu_3 a}(\boldsymbol{p}_3) \rangle\!\rangle, \qquad \text{(R.10.21)}
\end{aligned}$$

where L and R are given by (R.1.10) and (R.1.10). They lead to

$$\alpha_1^a = \alpha_2^a = \alpha_3^a = \alpha_4^a = \alpha_5^{(0)a} = 0, \tag{R.10.22}$$
$$\alpha_5^{(1)a} = -\frac{2^{4-\frac{d}{2}} c^a s_d}{\Gamma\left(\frac{d}{2}\right)}, \tag{R.10.23}$$

where

$$\begin{aligned}
c^a = \text{coefficient of } \delta^{\mu_1\mu_3} p_3^{\mu_2} p_3^{\nu_2} \left\{ \begin{array}{ll} p_3^{d-2} & \text{for odd } d \\ -p_3^{d-2} \log p_3^2 & \text{for even } d \end{array} \right\} \\
\text{in } p_1^{\nu_1} \langle\!\langle \frac{\delta T_{\mu_1\nu_1}}{\delta g^{\mu_2\nu_2}}(\boldsymbol{p}_1, \boldsymbol{p}_2) J^{\mu_3 a}(\boldsymbol{p}_3) \rangle\!\rangle. \qquad \text{(R.10.24)}
\end{aligned}$$

By $\alpha_5^{(n)a}$ we denote the coefficient of ϵ^n in the series expansion of α_5^a in the regulator ϵ. We assume that the α_1 coefficient is independent of ϵ.

$\langle T^{\mu_1\nu_1} T^{\mu_2\nu_2} T^{\mu_3\nu_3} \rangle$

Ward identities. The transverse and trace Ward identities are

$$\begin{aligned}
&p_1^{\nu_1} \langle\!\langle T_{\mu_1\nu_1}(\boldsymbol{p}_1) T_{\mu_2\nu_2}(\boldsymbol{p}_2) T_{\mu_3\nu_3}(\boldsymbol{p}_3) \rangle\!\rangle = \\
&\quad = 2p_1^{\nu_1} \langle\!\langle \frac{\delta T_{\mu_1\nu_1}}{\delta g^{\mu_3\nu_3}}(\boldsymbol{p}_1, \boldsymbol{p}_3) T_{\mu_2\nu_2}(\boldsymbol{p}_2) \rangle\!\rangle + 2p_1^{\nu_1} \langle\!\langle \frac{\delta T_{\mu_1\nu_1}}{\delta g^{\mu_2\nu_2}}(\boldsymbol{p}_1, \boldsymbol{p}_2) T_{\mu_3\nu_3}(\boldsymbol{p}_3) \rangle\!\rangle \\
&\quad + \; 2p_{1(\mu_3} \langle\!\langle T_{\nu_3)\mu_1}(\boldsymbol{p}_2) T_{\mu_2\nu_2}(-\boldsymbol{p}_2) \rangle\!\rangle + 2p_{1(\mu_2} \langle\!\langle T_{\nu_2)\mu_1}(\boldsymbol{p}_3) T_{\mu_3\nu_3}(-\boldsymbol{p}_3) \rangle\!\rangle \\
&\quad + \; \delta_{\mu_3\nu_3} p_3^{\alpha} \langle\!\langle T_{\alpha\mu_1}(\boldsymbol{p}_2) T_{\mu_2\nu_2}(-\boldsymbol{p}_2) \rangle\!\rangle + \delta_{\mu_2\nu_2} p_2^{\alpha} \langle\!\langle T_{\alpha\mu_1}(\boldsymbol{p}_3) T_{\mu_3\nu_3}(-\boldsymbol{p}_3) \rangle\!\rangle \\
&\quad - \; p_{3\mu_1} \langle\!\langle T_{\mu_2\nu_2}(\boldsymbol{p}_2) T_{\mu_3\nu_3}(-\boldsymbol{p}_2) \rangle\!\rangle - p_{2\mu_1} \langle\!\langle T_{\mu_2\nu_2}(\boldsymbol{p}_3) T_{\mu_3\nu_3}(-\boldsymbol{p}_3) \rangle\!\rangle, \qquad \text{(R.11.1)} \\
&\langle\!\langle T(\boldsymbol{p}_1) T_{\mu_2\nu_2}(\boldsymbol{p}_2) T_{\mu_3\nu_3}(\boldsymbol{p}_3) \rangle\!\rangle = \\
&\quad = 2\langle\!\langle \frac{\delta T}{\delta g^{\mu_2\nu_2}}(\boldsymbol{p}_1, \boldsymbol{p}_2) T_{\mu_3\nu_3}(\boldsymbol{p}_3) \rangle\!\rangle + 2\langle\!\langle \frac{\delta T}{\delta g^{\mu_3\nu_3}}(\boldsymbol{p}_1, \boldsymbol{p}_3) T_{\mu_2\nu_2}(\boldsymbol{p}_2) \rangle\!\rangle. \qquad \text{(R.11.2)}
\end{aligned}$$

Reconstruction formula. Define

$$\begin{aligned}
&\mathcal{L}^{\mu_1\nu_1\mu_2\nu_2\mu_3\nu_3}(\boldsymbol{p}_1,\boldsymbol{p}_2,\boldsymbol{p}_3) = \\
&\quad = 2\left[\mathscr{T}^{\mu_1\nu_1\alpha_1}(\boldsymbol{p}_1)p_1^{\beta_1} + \frac{\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)}{d-1}\delta^{\alpha_1\beta_1}\right]\delta^{\mu_3\alpha_3}\delta^{\nu_3\beta_3}\langle\!\langle\frac{\delta T_{\alpha_1\beta_1}}{\delta g^{\alpha_3\beta_3}}(\boldsymbol{p}_1,\boldsymbol{p}_3)T^{\mu_2\nu_2}(\boldsymbol{p}_2)\rangle\!\rangle \\
&\qquad + \Big[\mathscr{T}^{\mu_1\nu_1\beta_3}(\boldsymbol{p}_1)(2p_1^{(\mu_3}\delta^{\nu_3)\alpha_3} + p_3^{\alpha_3}\delta^{\mu_3\nu_3}) - p_3^{\alpha}\mathscr{T}_{\alpha}^{\mu_1\nu_1}(\boldsymbol{p}_1)\delta^{\mu_3\alpha_3}\delta^{\nu_3\beta_3} \\
&\qquad\qquad + \frac{2\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)}{d-1}\delta^{\mu_3\alpha_3}\delta^{\nu_3\beta_3}\Big]\langle\!\langle T_{\alpha_3\beta_3}(\boldsymbol{p}_2)T^{\mu_2\nu_2}(-\boldsymbol{p}_2)\rangle\!\rangle, \qquad \text{(R.11.3)}
\end{aligned}$$

where $\mathscr{T}^{\mu\nu\alpha}$ is defined in (R.1.20). The full 3-point function can be reconstructed from the transverse-traceless part as

$$\begin{aligned}
&\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)T^{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\!\rangle = \langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1)t^{\mu_2\nu_2}(\boldsymbol{p}_2)t^{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\!\rangle \\
&\quad + \sum_{\sigma}\mathcal{L}^{\mu_{\sigma(1)}\nu_{\sigma(1)}\mu_{\sigma(2)}\nu_{\sigma(2)}\mu_{\sigma(3)}\nu_{\sigma(3)}}(\boldsymbol{p}_{\sigma(1)},\boldsymbol{p}_{\sigma(2)},\boldsymbol{p}_{\sigma(3)}) \\
&\quad - \left[\mathscr{T}_{\alpha_3}^{\mu_3\nu_3}(\boldsymbol{p}_3)p_{3\beta_3} + \frac{\pi^{\mu_3\nu_3}(\boldsymbol{p}_3)}{d-1}\delta_{\alpha_3\beta_3}\right]\mathcal{L}^{\mu_1\nu_1\mu_2\nu_2\alpha_3\beta_3}(\boldsymbol{p}_1,\boldsymbol{p}_2,\boldsymbol{p}_3) \\
&\quad - [(\mu_1,\nu_1,\boldsymbol{p}_1)\mapsto(\mu_2,\nu_2,\boldsymbol{p}_2)\mapsto(\mu_3,\nu_3,\boldsymbol{p}_3)\mapsto(\mu_1,\nu_1,\boldsymbol{p}_1)] \\
&\quad - [(\mu_1,\nu_1,\boldsymbol{p}_1)\mapsto(\mu_3,\nu_3,\boldsymbol{p}_3)\mapsto(\mu_2,\nu_2,\boldsymbol{p}_2)\mapsto(\mu_1,\nu_1,\boldsymbol{p}_1)], \qquad \text{(R.11.4)}
\end{aligned}$$

where the sum is taken over all six permutations σ of the set $\{1,2,3\}$.

Decomposition of the 3-point function. The tensor decomposition of the transverse-traceless part is

$$\begin{aligned}
&\langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1)t^{\mu_2\nu_2}(\boldsymbol{p}_2)t^{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\!\rangle \\
&\quad = \Pi_{\alpha_1\beta_1}^{\mu_1\nu_1}(\boldsymbol{p}_1)\Pi_{\alpha_2\beta_2}^{\mu_2\nu_2}(\boldsymbol{p}_2)\Pi_{\alpha_3\beta_3}^{\mu_3\nu_3}(\boldsymbol{p}_3)\Big[A_1 p_2^{\alpha_1}p_2^{\beta_1}p_3^{\alpha_2}p_3^{\beta_2}p_1^{\alpha_3}p_1^{\beta_3} \\
&\qquad + A_2\delta^{\beta_1\beta_2}p_2^{\alpha_1}p_3^{\alpha_2}p_1^{\alpha_3}p_1^{\beta_3} + A_2(p_1\leftrightarrow p_3)\delta^{\beta_2\beta_3}p_2^{\alpha_1}p_2^{\beta_1}p_3^{\alpha_2}p_1^{\alpha_3} \\
&\qquad\qquad + A_2(p_2\leftrightarrow p_3)\delta^{\beta_1\beta_3}p_2^{\alpha_1}p_3^{\alpha_2}p_3^{\beta_2}p_1^{\alpha_3} \\
&\qquad + A_3\delta^{\alpha_1\alpha_2}\delta^{\beta_1\beta_2}p_1^{\alpha_3}p_1^{\beta_3} + A_3(p_1\leftrightarrow p_3)\delta^{\alpha_2\alpha_3}\delta^{\beta_2\beta_3}p_2^{\alpha_1}p_2^{\beta_1} \\
&\qquad\qquad + A_3(p_2\leftrightarrow p_3)\delta^{\alpha_1\alpha_3}\delta^{\beta_1\beta_3}p_3^{\alpha_2}p_3^{\beta_2} \\
&\qquad + A_4\delta^{\alpha_1\alpha_3}\delta^{\alpha_2\beta_3}p_2^{\beta_1}p_3^{\beta_2} + A_4(p_1\leftrightarrow p_3)\delta^{\alpha_1\alpha_3}\delta^{\alpha_2\beta_1}p_3^{\beta_2}p_1^{\beta_3} \\
&\qquad\qquad + A_4(p_2\leftrightarrow p_3)\delta^{\alpha_1\alpha_2}\delta^{\alpha_3\beta_2}p_2^{\beta_1}p_1^{\beta_3} \\
&\qquad + A_5\delta^{\alpha_1\beta_2}\delta^{\alpha_2\beta_3}\delta^{\alpha_3\beta_1}\Big]. \qquad \text{(R.11.5)}
\end{aligned}$$

The form factors A_j, $j=1,\dots,5$ are functions of the momentum magnitudes. If no arguments are specified then the standard ordering is assumed, $A_j = A_j(p_1,p_2,p_3)$, while by $p_i\leftrightarrow p_j$ we denote the exchange of the two momenta, *e.g.*, $A_1(p_1\leftrightarrow p_3) = A_2(p_3,p_2,p_1)$.

The form factors A_1 and A_5 are symmetric under any permutation of momenta, *i.e.*, for any permutation σ of the set $\{1,2,3\}$,

$$A_j(p_{\sigma(1)},p_{\sigma(2)},p_{\sigma(3)}) = A_j(p_1,p_2,p_3), \qquad j\in\{1,5\}. \qquad \text{(R.11.6)}$$

The remaining form factors are symmetric under $p_1\leftrightarrow p_2$, *i.e.*, they satisfy

$$A_j(p_2,p_1,p_3) = A_j(p_1,p_2,p_3), \qquad j\in\{2,3,4\}. \qquad \text{(R.11.7)}$$

The form factors can be calculated as

$$A_1 = \text{coefficient of } p_2^{\mu_1} p_2^{\nu_1} p_3^{\mu_2} p_3^{\nu_2} p_1^{\mu_3} p_1^{\nu_3}, \tag{R.11.8}$$

$$A_2 = 4 \cdot \text{coefficient of } \delta^{\nu_1\nu_2} p_2^{\mu_1} p_3^{\mu_2} p_1^{\mu_3} p_1^{\nu_3}, \tag{R.11.9}$$

$$A_3 = 2 \cdot \text{coefficient of } \delta^{\mu_1\mu_2}\delta^{\nu_1\nu_2} p_1^{\mu_3} p_1^{\nu_3}, \tag{R.11.10}$$

$$A_4 = 8 \cdot \text{coefficient of } \delta^{\mu_1\mu_3}\delta^{\mu_2\nu_3} p_2^{\nu_1} p_3^{\nu_2}, \tag{R.11.11}$$

$$A_5 = 8 \cdot \text{coefficient of } \delta^{\mu_1\nu_2}\delta^{\mu_2\nu_3}\delta^{\mu_3\nu_1}, \tag{R.11.12}$$

in $\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) T^{\mu_2\nu_2}(\boldsymbol{p}_2) T^{\mu_3\nu_3}(\boldsymbol{p}_3) \rangle\!\rangle$.

Primary conformal Ward identities. The primary CWIs are

$$\begin{aligned}
&\mathrm{K}_{12}\, A_1 = 0, && \mathrm{K}_{13}\, A_1 = 0,\\
&\mathrm{K}_{12}\, A_2 = 0, && \mathrm{K}_{13}\, A_2 = 8A_1,\\
&\mathrm{K}_{12}\, A_3 = 0, && \mathrm{K}_{13}\, A_3 = 2A_2,\\
&\mathrm{K}_{12}\, A_4 = 4\left[A_2(p_1 \leftrightarrow p_3) - A_2(p_2 \leftrightarrow p_3)\right], && \mathrm{K}_{13}\, A_4 = -4A_2(p_2 \leftrightarrow p_3),\\
&\mathrm{K}_{12}\, A_5 = 2\left[A_4(p_2 \leftrightarrow p_3) - A_4(p_1 \leftrightarrow p_3)\right], && \mathrm{K}_{13}\, A_5 = 2\left[A_4 - A_4(p_1 \leftrightarrow p_3)\right].
\end{aligned} \tag{R.11.13}$$

The solution in terms of triple-K integrals (R.1.7) is

$$A_1 = \alpha_1 J_{6\{000\}}, \tag{R.11.14}$$

$$A_2 = 4\alpha_1 J_{5\{001\}} + \alpha_2 J_{4\{000\}}, \tag{R.11.15}$$

$$A_3 = 2\alpha_1 J_{4\{002\}} + \alpha_2 J_{3\{001\}} + \alpha_3 J_{2\{000\}}, \tag{R.11.16}$$

$$A_4 = 8\alpha_1 J_{4\{110\}} - 2\alpha_2 J_{3\{001\}} + \alpha_4 J_{2\{000\}}, \tag{R.11.17}$$

$$A_5 = 8\alpha_1 J_{3\{111\}} + 2\alpha_2 \left(J_{2\{110\}} + J_{2\{101\}} + J_{2\{011\}}\right) + \alpha_5 J_{0\{000\}}, \tag{R.11.18}$$

where α_j, $j = 1, \ldots, 5$ are constants. If the integrals diverge, the regularisation (R.1.8) should be used.

Secondary conformal Ward identities. The independent secondary CWIs are

$$\begin{aligned}
&(*)\ \mathrm{L}_{2,4}\, A_1 + \mathrm{R}_2\left[A_2 - A_2(p_2 \leftrightarrow p_3)\right] =\\
&\quad = 2d \cdot \text{coeff. of } p_2^{\mu_1} p_3^{\mu_2} p_3^{\nu_2} p_1^{\mu_3} p_1^{\nu_3} \text{ in } p_{1\nu_1} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) T^{\mu_2\nu_2}(\boldsymbol{p}_2) T^{\mu_3\nu_3}(\boldsymbol{p}_3) \rangle\!\rangle,
\end{aligned} \tag{R.11.19}$$

$$\begin{aligned}
&\mathrm{L}_{2,4}\, A_2 + 2\,\mathrm{R}_2\left[2A_3 - A_4(p_1 \leftrightarrow p_3)\right] =\\
&\quad = 8d \cdot \text{coefficient of } \delta^{\mu_1\mu_2} p_3^{\nu_2} p_1^{\mu_3} p_1^{\nu_3} \text{ in } p_{1\nu_1} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) T^{\mu_2\nu_2}(\boldsymbol{p}_2) T^{\mu_3\nu_3}(\boldsymbol{p}_3) \rangle\!\rangle,
\end{aligned} \tag{R.11.20}$$

$$\begin{aligned}
&(*)\ \mathrm{L}_{2,2}\left[A_2(p_1 \leftrightarrow p_3)\right] + \mathrm{R}_2\left[A_4(p_2 \leftrightarrow p_3) - A_4\right] + 2p_1^2\left[A_2(p_2 \leftrightarrow p_3) - A_2\right] =\\
&\quad = 8d \cdot \text{coefficient of } \delta^{\mu_2\mu_3} p_2^{\mu_1} p_3^{\nu_2} p_1^{\nu_3} \text{ in } p_{1\nu_1} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) T^{\mu_2\nu_2}(\boldsymbol{p}_2) T^{\mu_3\nu_3}(\boldsymbol{p}_3) \rangle\!\rangle,
\end{aligned} \tag{R.11.21}$$

$$\begin{aligned}
&\mathrm{L}_{2,2}\left[A_4(p_2 \leftrightarrow p_3)\right] - 2\,\mathrm{R}_2\, A_5 + 2p_1^2\left[A_4(p_1 \leftrightarrow p_3) - 4A_3\right] =\\
&\quad = 16d \cdot \text{coefficient of } \delta^{\mu_1\mu_2}\delta^{\mu_3\nu_2} p_1^{\nu_3} \text{ in } p_{1\nu_1} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) T^{\mu_2\nu_2}(\boldsymbol{p}_2) T^{\mu_3\nu_3}(\boldsymbol{p}_3) \rangle\!\rangle,
\end{aligned} \tag{R.11.22}$$

$$\begin{aligned}
&\mathrm{L}_{2,0}\left[A_3(p_1 \leftrightarrow p_3)\right] + p_1^2\left[A_4 - A_4(p_2 \leftrightarrow p_3)\right] =\\
&\quad = 4d \cdot \text{coefficient of } \delta^{\mu_2\mu_3}\delta^{\nu_2\nu_3} p_2^{\mu_1} \text{ in } p_{1\nu_1} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) T^{\mu_2\nu_2}(\boldsymbol{p}_2) T^{\mu_3\nu_3}(\boldsymbol{p}_3) \rangle\!\rangle,
\end{aligned} \tag{R.11.23}$$

where the operators L and R are defined in (R.1.9) and (R.1.10). The identities denoted by asterisks are redundant, *i.e.*, they are trivially satisfied in all cases and do not impose any additional conditions on primary constants. The secondary CWIs lead to which lead to

$$\alpha_4 = (3d+2)\alpha_2 + 2\alpha_3, \tag{R.11.24}$$

$$\alpha_5^{(0)} = -2d^2\alpha_2, \tag{R.11.25}$$

$$2d(d+2)\alpha_1 + d\alpha_2 + \alpha_3 = -\frac{2^{3-\frac{d}{2}} s_d c_T}{\Gamma\left(\frac{d}{2}\right)}, \tag{R.11.26}$$

$$d(d+2)(8d\alpha_1 + 3\alpha_2) + 2d\alpha_3 + 2\alpha_5^{(1)} = -\frac{2^{5-\frac{d}{2}} s_d d}{\Gamma\left(\frac{d}{2}\right)}\left[c_T + 2c_g c_T\right]. \tag{R.11.27}$$

By $\alpha_5^{(n)}$ we mean the coefficient of ϵ^n in the series expansion of α_5 in the regulator ϵ. We assume that the α_1 coefficient is independent of ϵ, *i.e.*, $\alpha_1 = \alpha_1^{(0)}$. The constant c_T is the 2-point function normalisation (R.1.23), while s_d is defined in (R.1.3) and the constant c_g is defined as

$$\langle\!\langle \frac{\delta T_{\mu_1\nu_1}}{\delta g^{\mu_2\nu_2}}(\boldsymbol{p}_1,\boldsymbol{p}_2) T_{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\!\rangle = 4c_g \delta_{(\mu_1(\mu_2}\langle\!\langle T_{\nu_1)\nu_2)}(\boldsymbol{p}_3) T_{\mu_3\nu_3}(-\boldsymbol{p}_3)\rangle\!\rangle + \dots \tag{R.11.28}$$

The omitted terms do not contain the tensor structures listed explicitly. If the regulator is present, the limit $\epsilon \to 0$ can be taken after the primary constants and triple-K integrals are substituted into the primary CWIs (R.11.14) - (R.11.18).

The 3-point function $\langle\!\langle T^{\mu_1\nu_1} T^{\mu_2\nu_2} T^{\mu_3\nu_3}\rangle\!\rangle$ therefore depends on the 2-point function normalisations c_T and c_g and two undetermined primary constants α_1 and α_2. The dependence of this correlator on two 2-point function normalisations rather than only one as found in [3] is related to the definition (4.47) we adopt for this correlator. Our definition differs by the semi-local terms on the right-hand side of (4.47), and it is these terms that produce the dependence of our solution on c_g through (R.11.28). (Similar considerations also apply for $\langle\!\langle T_{\mu_1\nu_1} J^{\mu_2} J^{\mu_3}\rangle\!\rangle$ as discussed above (7.32).)

An additional effect in dimension $d = 3$ is that the tensor decomposition becomes degenerate meaning there are only two instead of the usual five form factors. In consequence, the stress-energy tensor 3-point function in $d = 3$ only depends on the primary constant α_1, rather than on both α_1 and α_2. Along with the two 2-point function normalisations, this makes three parameters in total (or two using the definition of the 3-point function in [3]). We present a discussion of this degeneracy in appendix A.2, the results of which we make use of below.

Examples

For $d = 3$ we find

$$A_1 = \frac{8\alpha_1}{a_{123}^6}\left[a_{123}^3 + 3a_{123}b_{123} + 15c_{123}\right], \tag{R.11.29}$$

$$\begin{aligned} A_2 = {} & \frac{8\alpha_1}{a_{123}^5}\left[4p_3^4 + 20p_3^3 a_{12} + 4p_3^2(7a_{12}^2 + 6b_{12}) + 15p_3 a_{12}(a_{12}^2 + b_{12}) + 3a_{12}^2(a_{12}^2 + b_{12})\right] \\ & + \frac{2\alpha_2}{a_{123}^4}\left[a_{123}^3 + a_{123}b_{123} + 3c_{123}\right], \end{aligned} \tag{R.11.30}$$

$$\begin{aligned} A_3 = {} & \frac{2\alpha_1 p_3^2}{a_{123}^4}\left[7p_3^3 + 28p_3^2 a_{12} + 3p_3(11a_{12}^2 + 6b_{12}) + 12a_{12}(a_{12}^2 + b_{12})\right] \\ & + \frac{\alpha_2 p_3^2}{a_{123}^3}\left[p_3^2 + 3p_3 a_{12} + 2(a_{12}^2 + b_{12})\right] - \frac{2c_T}{a_{123}^2}\left[a_{123}^3 - a_{123}b_{123} - c_{123}\right], \end{aligned} \tag{R.11.31}$$

$$\begin{aligned} A_4 = {} & \frac{4\alpha_1}{a_{123}^4}\left[-3p_3^5 - 12p_3^4 a_{12} - 9p_3^3(a_{12}^2 + 2b_{12}) + 9p_3^2 a_{12}(a_{12}^2 - 3b_{12})\right. \\ & \left. + (4p_3 + a_{12})(3a_{12}^4 - 3a_{12}^2 b_{12} + 4b_{12}^2)\right] \\ & + \frac{\alpha_2}{a_{123}^3}\left[-p_3^4 - 3p_3^3 a_{12} - 6p_3^2 b_{12} + a_{12}(a_{12}^2 - b_{12})(3p_3 + a_{12})\right] \\ & - \frac{4c_T}{a_{123}^2}\left[a_{123}^3 - a_{123}b_{123} - c_{123}\right], \end{aligned} \tag{R.11.32}$$

$$A_5 = \frac{2\alpha_1}{a_{123}^3}\left[-3a_{123}^6 + 9a_{123}^4 b_{123} + 12a_{123}^2 b_{123}^2 - 33a_{123}^3 c_{123} + 12a_{123}b_{123}c_{123} + 8c_{123}^2\right]$$
$$+ \frac{\alpha_2}{2a_{123}^2}\left[-a_{123}^5 + 3a_{123}^3 b_{123} + 4a_{123}b_{123}^2 - 11a_{123}^2 c_{123} + 4b_{123}c_{123}\right]$$
$$+ 2(c_T + 4c_g c_T)(p_1^3 + p_2^3 + p_3^3). \tag{R.11.33}$$

For $d = 4$ we find

$$A_1 = \alpha_1 I_{7\{222\}}, \tag{R.11.34}$$
$$A_2 = \left(16\alpha_1 + \alpha_2 - 4\alpha_1 p_1 \frac{\partial}{\partial p_1}\right) I^{(0)}_{5\{222\}}, \tag{R.11.35}$$
$$A_3 = \left(2c_T - (18\alpha_1 + \alpha_2)p_3 \frac{\partial}{\partial p_3} + 2\alpha_1 p_3^2 \frac{\partial^2}{\partial p_3^2}\right) I^{(0)}_{3\{222\}}, \tag{R.11.36}$$
$$A_4 = 2\left(16\alpha_1 - \alpha_2 + 2c_T - 16\alpha_1 p_1 \frac{\partial}{\partial p_1} - 16\alpha_1 p_2 \frac{\partial}{\partial p_2} + \alpha_2 p_3 \frac{\partial}{\partial p_3}\right.$$
$$\left. + 4\alpha_1 p_1 p_2 \frac{\partial^2}{\partial p_1 \partial p_2}\right) I^{(0)}_{3\{222\}}, \tag{R.11.37}$$
$$A_5 = 2\left[32(8\alpha_1 + \alpha_2) - 8(8\alpha_1 + \alpha_2)\left(p_1 \frac{\partial}{\partial p_1} + p_2 \frac{\partial}{\partial p_2} + p_3 \frac{\partial}{\partial p_3}\right)\right.$$
$$+ (16\alpha_1 + \alpha_2)\left(p_1 p_2 \frac{\partial^2}{\partial p_1 \partial p_2} + p_1 p_3 \frac{\partial^2}{\partial p_1 \partial p_3} + p_2 p_3 \frac{\partial^2}{\partial p_2 \partial p_3}\right)$$
$$\left. - 4\alpha_1 p_1 p_2 p_3 \frac{\partial^3}{\partial p_1 \partial p_2 \partial p_3}\right] I^{(0)}_{1\{222\}}$$
$$+ 2\left[48\alpha_1 + 5\alpha_2 - 2(c_T + 4c_g c_T)\right] \times \left[\left(p_1^4 \log p_1 - p_2^2 p_3^2 - p_1^4(\tfrac{3}{4} - \gamma_E + \log 2)\right)\right.$$
$$\left. + (p_1 \leftrightarrow p_2) + (p_1 \leftrightarrow p_3)\right], \tag{R.11.38}$$

where $I^{(0)}_{\alpha\{\beta_j\}}$ is the coefficient of ϵ^0 in the series expansion of the regulated integral $I_{\alpha+\epsilon\{\beta_j\}}$. The integrals can be obtained from the master integral (A.6.1) via the reduction scheme in table 1, page 44.

In case of $d = 4$ the trace Ward identity is anomalous,

$$\langle T \rangle = aE_4 + cW^2, \tag{R.11.39}$$

where E_4 is the Euler density and W^2 square of the Weyl tensor. It leads to the anomalous contribution,

$$\langle\langle T(\boldsymbol{p}_1) T^{\mu_2\nu_2}(\boldsymbol{p}_2) T^{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\rangle_{\text{anomaly}} = \Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)\Pi^{\mu_3\nu_3}_{\alpha_3\beta_3}(\boldsymbol{p}_3)\left[B_1 p_3^{\alpha_2} p_3^{\beta_2} p_1^{\alpha_3} p_1^{\beta_3}\right.$$
$$\left. + B_2 \delta^{\beta_2\beta_3} p_3^{\alpha_2} p_1^{\alpha_3} + B_3 \delta^{\alpha_2\alpha_3}\delta^{\beta_2\beta_3}\right]$$
$$+ \pi^{\mu_2\nu_2}(\boldsymbol{p}_2)\Pi^{\mu_3\nu_3}_{\alpha_3\beta_3}(\boldsymbol{p}_3) p_1^{\alpha_3} p_1^{\beta_3} B_4 + \Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)\pi^{\mu_3\nu_3}(\boldsymbol{p}_3) p_3^{\alpha_2} p_3^{\beta_2} B_4(p_2 \leftrightarrow p_3)$$
$$+ \pi^{\mu_2\nu_2}(\boldsymbol{p}_2)\pi^{\mu_3\nu_3}(\boldsymbol{p}_3) B_5, \tag{R.11.40}$$

where

$$B_1 = 8(a + c), \tag{R.11.41}$$
$$B_2 = 8(p_1^2 - p_2^2 - p_3^2)(a + c), \tag{R.11.42}$$
$$B_3 = -2\lambda^2(a + c) + 4p_2^2 p_3^2 c, \tag{R.11.43}$$
$$B_4 = -\frac{8}{3} a p_3^2, \tag{R.11.44}$$
$$B_5 = \frac{4}{9} a \lambda^2 \tag{R.11.45}$$

and $\lambda^2 = (p_1+p_2+p_3)(p_1+p_2-p_3)(p_1-p_2+p_3)(-p_1+p_2+p_3)$. The anomalous contribution to the full 3-point function is then

$$\begin{aligned}\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)T^{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\!\rangle_{\text{anomaly}} &= \frac{\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)}{d-1}\langle\!\langle T(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)T^{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\!\rangle_{\text{anomaly}} \\ &\quad [(\boldsymbol{p}_1,\mu_1,\nu_1)\leftrightarrow(\boldsymbol{p}_2,\mu_2,\nu_2)] + [(\boldsymbol{p}_1,\mu_1,\nu_1)\leftrightarrow(\boldsymbol{p}_3,\mu_3,\nu_3)] \\ &- \frac{\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)\pi^{\mu_2\nu_2}(\boldsymbol{p}_2)}{d-1}\left[\Pi^{\mu_3\nu_3}_{\alpha_3\beta_3}(\boldsymbol{p}_3)B_4(p_1,p_2,p_3)+\pi^{\mu_3\nu_3}(\boldsymbol{p}_3)B_5(p_1,p_2,p_3)\right] \\ &\quad [(\boldsymbol{p}_1,\mu_1,\nu_1)\leftrightarrow(\boldsymbol{p}_3,\mu_3,\nu_3)] + [(\boldsymbol{p}_2,\mu_2,\nu_2)\leftrightarrow(\boldsymbol{p}_3,\mu_3,\nu_3)] \\ &+ \frac{\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)\pi^{\mu_2\nu_2}(\boldsymbol{p}_2)\pi^{\mu_3\nu_3}(\boldsymbol{p}_3)}{d-1}B_5. \end{aligned} \tag{R.11.46}$$

and should be added to the right hand side of (R.11.4).

The anomalies are equal

$$a = \frac{1}{2}(16\alpha_1+\alpha_2-c_T), \qquad c = \frac{c_T}{2}, \tag{R.11.47}$$

which allows us to rewrite the solution (R.11.34) - (R.11.38) as

$$A_1 = \alpha_1 I_{7\{222\}}, \tag{R.11.48}$$

$$A_2 = 2\left(a+c-2\alpha_1 p_1\frac{\partial}{\partial p_1}\right)I^{(0)}_{5\{222\}}, \tag{R.11.49}$$

$$A_3 = 2\left(2c-(a+c+\alpha_1)p_3\frac{\partial}{\partial p_3}+\alpha_1 p_3^2\frac{\partial^2}{\partial p_3^2}\right)I^{(0)}_{3\{222\}}, \tag{R.11.50}$$

$$\begin{aligned}A_4 = 4\Bigg(&16\alpha_1 - a - 8\alpha_1 p_1\frac{\partial}{\partial p_1} - 8\alpha_1 p_2\frac{\partial}{\partial p_2} + (a+c-8\alpha_1)p_3\frac{\partial}{\partial p_3} \\ &+ 2\alpha_1 p_1 p_2\frac{\partial^2}{\partial p_1\partial p_2}\Bigg)I^{(0)}_{3\{222\}},\end{aligned} \tag{R.11.51}$$

$$\begin{aligned}A_5 = 4\Bigg[&32(a+c-4\alpha_1) - 8(a+c-4\alpha_1)\left(p_1\frac{\partial}{\partial p_1}+p_2\frac{\partial}{\partial p_2}+p_3\frac{\partial}{\partial p_3}\right) \\ &+ (a+c)\left(p_1p_2\frac{\partial^2}{\partial p_1\partial p_2}+p_1p_3\frac{\partial^2}{\partial p_1\partial p_3}+p_2p_3\frac{\partial^2}{\partial p_2\partial p_3}\right) \\ &- 2\alpha_1 p_1p_2p_3\frac{\partial^3}{\partial p_1\partial p_2\partial p_3}\Bigg]I^{(0)}_{1\{222\}} \\ &+ 4\left[5a-16\alpha_1+3c-8c_gc\right]\times\left[\left(p_1^4\log p_1 - p_2^2p_3^2 - p_1^4(\tfrac{3}{4}-\gamma_E+\log 2)\right)\right. \\ &\left.+(p_1\leftrightarrow p_2)+(p_1\leftrightarrow p_3)\right].\end{aligned} \tag{R.11.52}$$

The scheme dependence of the solution due to the counterterms is

$$A_2 \mapsto A_2 - 16(a_0+c_0), \tag{R.11.53}$$

$$A_3 \mapsto A_3 + 8\left[c_0(p_1^2+p_2^2)-a_0p_3^2\right], \tag{R.11.54}$$

$$A_4 \mapsto A_4 + 8\left[c_0(p_1^2+p_2^2+3p_3^2)-a_0(p_1^2+p_2^2-p_3^2)\right], \tag{R.11.55}$$

$$A_5 \mapsto A_5 - 4\left[a_0\lambda^2 + c_0(p_1^2+p_2^2+p_3^2)^2 + 8c_gc_0(p_1^4+p_2^4+p_3^4)\right], \tag{R.11.56}$$

where a_0 and c_0 are arbitrary constants. See section 8.3 for details.

For $d=5$ we find

$$A_1 = \frac{72\alpha_1}{a_{123}^7}\left[a_{123}^2(a_{123}^4+a_{123}^2b_{123}+b_{123}^2)+a_{123}(a_{123}^2+5b_{123})c_{123}+10c_{123}^2\right], \tag{R.11.57}$$

$$
\begin{aligned}
A_2 = \frac{24\alpha_1}{a_{123}^6}\big[&-12p_3^7 - 72p_3^6a_{12} + 24p_3^5(-8a_{12}^2 + b_{12}) + 24p_3^4a_{12}(-13a_{12}^2 + 6b_{12}) \\
&+ 8p_3^3(-42a_{12}^4 + 33a_{12}^2b_{12} + 8b_{12}^2) + 3p_3^2a_{12}(-77a_{12}^4 + 73a_{12}^2b_{12} + 23b_{12}^2) \\
&+ 30p_3a_{12}^2(-3a_{12}^4 + 3a_{12}^2b_{12} + b_{12}^2) + 5a_{12}^3(-3a_{12}^4 + 3a_{12}^2b_{12} + b_{12}^2)\big] \\
+ \frac{6\alpha_2}{a_{123}^5}\big[&-3a_{123}^6 + 3a_{123}^4b_{123} + a_{123}^2b_{123}^2 + a_{123}^3c_{123} + 3a_{123}b_{123}c_{123} + 4c_{123}^2\big], \qquad \text{(R.11.58)}
\end{aligned}
$$

$$
\begin{aligned}
A_3 = \frac{2\alpha_1p_3^2}{a_{123}^5}\big[&-81p_3^6 - 405p_3^5a_{12} - 3p_3^4(281a_{12}^2 - 22b_{12}) - 15p_3^3a_{12}(65a_{12}^2 - 22b_{12}) \\
&- 8p_3^2(87a_{12}^4 - 63a_{12}^2b_{12} - 13b_{12}^2) - 100p_3a_{12}(3a_{12}^4 - 3a_{12}^2b_{12} - b_{12}^2) \\
&- 20a_{12}^2(3a_{12}^4 - 3a_{12}^2b_{12} - b_{12}^2)\big] \\
+ \frac{a_2p_3^2}{a_{123}^4}\big[&-9p_3^5 - 36p_3^4a_{12} - 3p_3^3(19a_{12}^2 - 2b_{12}) - 24p_3^2a_{12}(2a_{12}^2 - b_{12}) \\
&- 8p_3(3a_{12}^4 - 3a_{12}^2b_{12} - b_{12}^2) - 2a_{12}(3a_{12}^4 - 3a_{12}^2b_{12} - b_{12}^2)\big] \\
- \frac{2c_T}{3a_{123}^3}\big[&3a_{123}^2(a_{123}^4 - 3a_{123}^2b_{123} + b_{123}^2) + 3a_{123}(a_{123}^2 + b_{123})c_{123} + 2c_{123}^2\big], \qquad \text{(R.11.59)}
\end{aligned}
$$

$$
\begin{aligned}
A_4 = \frac{4\alpha_1}{a_{123}^5}\big[&45p_3^8 + 225p_3^7a_{12} + 15p_3^6(29a_{12}^2 + 2b_{12}) + 75p_3^5a_{12}(5a_{12}^2 + 2b_{12}) \\
&+ 8p_3^4(75a_{12}^2 - 23b_{12})b_{12} - 5p_3^3a_{12}(75a_{12}^4 - 255a_{12}^2b_{12} + 79b_{12}^2) \\
&- p_3^2(435a_{12}^6 - 1335a_{12}^4b_{12} + 343a_{12}^2b_{12}^2 - 96b_{12}^3) \\
&- 3a_{12}(5p_3 + a_{12})(15a_{12}^6 - 45a_{12}^4b_{12} + 11a_{12}^2b_{12}^2 - 4b_{12}^3)\big] \\
+ \frac{\alpha_2}{a_{123}^4}\big[&9p_3^7 + 36p_3^6a_{12} + 3p_3^5(17a_{12}^2 + 2b_{12}) + 24p_3^4a_{12}(a_{12}^2 + b_{12}) \\
&- 8p_3^3(3a_{12}^4 - 12a_{12}^2b_{12} + 5b_{12}^2) - p_3^2a_{12}(51a_{12}^4 - 159a_{12}^2b_{12} + 55b_{12}^2) \\
&- 9a_{12}^2(4p_3 + a_{12})(a_{12}^4 - 3a_{12}^2b_{12} + b_{12}^2)\big] \\
- \frac{4c_T}{3a_{123}^3}\big[&3a_{123}^2(a_{123}^4 - 3a_{123}^2b_{123} + b_{123}^2) + 3a_{123}(a_{123}^2 + b_{123})c_{123} + 2c_{123}^2\big], \qquad \text{(R.11.60)}
\end{aligned}
$$

$$
\begin{aligned}
A_5 = \frac{6\alpha_1}{(p_1+p_2+p_3)^4}\big[&5a_{123}^3(a_{123}^2 - 4b_{123})(3a_{123}^4 - 3a_{123}^2b_{123} - b_{123}^2) \\
&+ 5a_{123}^2(23a_{123}^4 - 23a_{123}^2b_{123} + 4b_{123}^2)c_{123} - 4a_{123}(a_{123}^2 - 4b_{123})c_{123}^2 + 8c_{123}^3\big] \\
+ \frac{\alpha_2}{2(p_1+p_2+p_3)^3}\big[&3a_{123}^2(a_{123}^2 - 4b_{123})(3a_{123}^4 - 3a_{123}^2b_{123} - b_{123}^2) \\
&+ 3a_{123}(23a_{123}^4 - 23a_{123}^2b_{123} + 4b_{123}^2)c_{123} - 4(a_{123}^2 - 2b_{123})c_{123}^2\big] \\
+ 2(c_T + 4c_gc_T)&(p_1^5 + p_2^5 + p_3^5). \qquad \text{(R.11.61)}
\end{aligned}
$$

Appendix

A.1 Decomposition of $\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}T^{\mu_3\nu_3}\rangle$ in non-conformal case

In this section we present the decomposition of the stress-energy tensor 3-point function for a general quantum field theory. As the stress-energy tensor in a general theory is no longer traceless, our arguments in the main text need some minor modifications. First, we discuss how to reconstruct the full correlation function from the purely transverse part, making use of the transverse Ward identities in a similar fashion to section 3.3. We then proceed to construct the general tensor decomposition of this transverse part in terms of ten independent form factors.

As in the main text, we will denote the transverse-traceless part of the stress-energy tensor by $t^{\mu\nu} = \Pi^{\mu\nu}_{\alpha\beta}T^{\alpha\beta}$. Here, we will also make use of the purely transverse part, $t_T^{\mu\nu} = \pi^\mu_\alpha\pi^\nu_\beta T^{\alpha\beta}$, which includes a nonvanishing trace part $(t_T)^\mu_\mu$. The difference between the stress-energy tensor and its transverse part can then be written $\tilde{t}^{\mu\nu}_{\text{loc}} = T^{\mu\nu} - t_T^{\mu\nu}$, *i.e.*,

$$\tilde{t}^{\mu\nu}_{\text{loc}} = \left(\frac{p^\mu}{p^2}\delta^\nu_\alpha + \frac{p^\nu}{p^2}\delta^\mu_\alpha - \frac{p^\mu p^\nu p_\alpha}{p^4}\right)p_\beta T^{\alpha\beta}. \tag{A.1.1}$$

To obtain the reconstruction formula, we use the Ward identity (R.11.1) to re-express $p_\beta T^{\alpha\beta}$ in terms of 2-point functions when the expectation value of $\tilde{t}^{\mu\nu}_{\text{loc}}$ with other operators is taken. Defining the operator

$$\begin{aligned}\tilde{\mathcal{L}}^{\mu_1\nu_1\mu_2\nu_2\mu_3\nu_3}(\boldsymbol{p}_1,\boldsymbol{p}_2,\boldsymbol{p}_3) = \frac{1}{p_1^2}\left(2p_1^{(\mu_1}\delta^{\nu_1)}_{\alpha_1} - \frac{p_1^{\mu_1}p_1^{\nu_1}p_{1\alpha_1}}{p_1^2}\right)\times\\ \times\Bigg[2\delta^{\mu_3\alpha_3}\delta^{\nu_3\alpha_3}p_1^{\beta_1}\langle\!\langle\frac{\delta T_{\alpha_1\beta_1}}{\delta g^{\alpha_3\beta_3}}(\boldsymbol{p}_1,\boldsymbol{p}_3)T^{\mu_2\nu_2}(\boldsymbol{p}_2)\rangle\!\rangle\\ + \left(\delta^{\beta_3\alpha_1}(2p_1^{(\mu_3}\delta^{\nu_3)\alpha_3} + p_3^{\alpha_3}\delta^{\mu_3\nu_3}) - p_3^{\alpha_1}\delta^{\alpha_3\mu_3}\delta^{\beta_3\nu_3}\right)\langle\!\langle T_{\alpha_3\beta_3}(\boldsymbol{p}_2)T^{\mu_2\nu_2}(-\boldsymbol{p}_2)\rangle\!\rangle\Bigg],\end{aligned} \tag{A.1.2}$$

the reconstruction formula takes the form

$$\begin{aligned}\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)T^{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\!\rangle &= \langle\!\langle t_T^{\mu_1\nu_1}(\boldsymbol{p}_1)t_T^{\mu_2\nu_2}(\boldsymbol{p}_2)t_T^{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\!\rangle\\ &+ \sum_\sigma \tilde{\mathcal{L}}^{\mu_{\sigma(1)}\nu_{\sigma(1)}\mu_{\sigma(2)}\nu_{\sigma(2)}\mu_{\sigma(3)}\nu_{\sigma(3)}}(\boldsymbol{p}_{\sigma(1)},\boldsymbol{p}_{\sigma(2)},\boldsymbol{p}_{\sigma(3)})\\ &- \frac{1}{p_3^2}\left(2p_3^{(\mu_3}\delta^{\nu_3)}_{\alpha_3} - \frac{p_3^{\mu_3}p_3^{\nu_3}p_{3\alpha_3}}{p_3^2}\right)p_{3\beta_3}\tilde{\mathcal{L}}^{\mu_1\nu_1\mu_2\nu_2\alpha_3\beta_3}(\boldsymbol{p}_1,\boldsymbol{p}_2,\boldsymbol{p}_3)\\ &- [(\mu_1,\nu_1,\boldsymbol{p}_1)\mapsto(\mu_2,\nu_2,\boldsymbol{p}_2)\mapsto(\mu_3,\nu_3,\boldsymbol{p}_3)\mapsto(\mu_1,\nu_1,\boldsymbol{p}_1)]\\ &- [(\mu_1,\nu_1,\boldsymbol{p}_1)\mapsto(\mu_3,\nu_3,\boldsymbol{p}_3)\mapsto(\mu_2,\nu_2,\boldsymbol{p}_2)\mapsto(\mu_1,\nu_1,\boldsymbol{p}_1)],\end{aligned} \tag{A.1.3}$$

where the sum is taken over all six permutations σ of the set $\{1,2,3\}$. Note the similarity between these expression and (R.11.3, R.11.4).

We turn now to the tensor decomposition of the purely transverse part of the 3-point function. The most general form of this is

$$\begin{aligned}&\langle\!\langle t_T^{\mu_1\nu_1}(\boldsymbol{p}_1)t_T^{\mu_2\nu_2}(\boldsymbol{p}_2)t_T^{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\!\rangle =\\ &\quad= \pi^{\mu_1}_{(\alpha_1}(\boldsymbol{p}_1)\pi^{\nu_1}_{\beta_1)}(\boldsymbol{p}_1)\pi^{\mu_2}_{(\alpha_2}(\boldsymbol{p}_2)\pi^{\nu_2}_{\beta_2)}(\boldsymbol{p}_2)\pi^{\mu_3}_{(\alpha_3}(\boldsymbol{p}_3)\pi^{\nu_3}_{\beta_3)}(\boldsymbol{p}_3)X^{\alpha_1\beta_1\alpha_2\beta_2\alpha_3\beta_3},\end{aligned} \tag{A.1.4}$$

where $X^{\alpha_1\beta_1\alpha_2\beta_2\alpha_3\beta_3}$ is a general tensor built from the metric $\delta^{\mu\nu}$ and two independent momenta, with a kinematic dependence on the momentum magnitudes p_1, p_2 and p_3. Note, however, that if $X^{\alpha_1\beta_1\alpha_2\beta_2\alpha_3\beta_3}$ contains $p_j^{\alpha_j}$ or $p_j^{\beta_j}$ for $j\in\{1,2,3\}$ then the contractions with the corresponding transverse projectors vanish. We will assume that $X^{\alpha_1\beta_1\alpha_2\beta_2\alpha_3\beta_3}$ is symmetric under $\alpha_j\leftrightarrow\beta_j$

	$p_2^{\alpha_1} p_2^{\beta_1} p_3^{\alpha_2} p_3^{\beta_2} p_1^{\alpha_3} p_1^{\beta_3}$	
$\delta^{\beta_1\beta_2} p_2^{\alpha_1} p_3^{\alpha_2} p_1^{\alpha_3} p_1^{\beta_3}$	$\delta^{\beta_2\beta_3} p_2^{\alpha_1} p_2^{\beta_1} p_3^{\alpha_2} p_1^{\alpha_3}$	$\delta^{\beta_1\beta_3} p_2^{\alpha_1} p_3^{\alpha_2} p_3^{\beta_2} p_1^{\alpha_3}$
$\delta^{\alpha_1\alpha_2}\delta^{\beta_1\beta_2} p_1^{\alpha_3} p_1^{\beta_3}$	$\delta^{\alpha_2\alpha_3}\delta^{\beta_2\beta_3} p_2^{\alpha_1} p_2^{\beta_1}$	$\delta^{\alpha_1\alpha_3}\delta^{\beta_1\beta_3} p_3^{\alpha_2} p_3^{\beta_2}$
$\delta^{\alpha_1\alpha_3}\delta^{\alpha_2\beta_3} p_2^{\beta_1} p_3^{\beta_2}$	$\delta^{\alpha_1\alpha_3}\delta^{\alpha_2\beta_1} p_3^{\beta_2} p_1^{\beta_3}$	$\delta^{\alpha_1\alpha_2}\delta^{\alpha_3\beta_2} p_2^{\beta_1} p_1^{\beta_3}$
	$\delta^{\alpha_1\beta_2}\delta^{\alpha_2\beta_3}\delta^{\alpha_3\beta_1}$	
$\delta^{\alpha_3\beta_3} p_3^{\alpha_1} p_3^{\beta_1} p_3^{\alpha_2} p_3^{\beta_2}$	$\delta^{\alpha_1\beta_1} p_3^{\alpha_2} p_3^{\beta_2} p_1^{\alpha_3} p_1^{\beta_3}$	$\delta^{\alpha_2\beta_2} p_3^{\alpha_1} p_3^{\beta_1} p_1^{\alpha_3} p_1^{\beta_3}$
$\delta^{\alpha_3\beta_3}\delta^{\beta_1\beta_2} p_2^{\alpha_1} p_3^{\alpha_2}$	$\delta^{\alpha_1\beta_1}\delta^{\beta_2\beta_3} p_3^{\alpha_2} p_1^{\alpha_3}$	$\delta^{\alpha_2\beta_2}\delta^{\beta_1\beta_3} p_2^{\alpha_1} p_1^{\alpha_3}$
$\delta^{\alpha_3\beta_3}\delta^{\alpha_1\alpha_2}\delta^{\beta_1\beta_2}$	$\delta^{\alpha_1\beta_1}\delta^{\alpha_2\alpha_3}\delta^{\beta_2\beta_3}$	$\delta^{\alpha_2\beta_2}\delta^{\alpha_1\alpha_3}\delta^{\beta_1\beta_3}$
$\delta^{\alpha_1\beta_1}\delta^{\alpha_2\beta_2} p_1^{\alpha_3} p_1^{\beta_3}$	$\delta^{\alpha_2\beta_2}\delta^{\alpha_3\beta_3} p_2^{\alpha_1} p_2^{\beta_1}$	$\delta^{\alpha_1\beta_1}\delta^{\alpha_3\beta_3} p_3^{\alpha_2} p_3^{\beta_2}$
	$\delta^{\alpha_1\beta_1}\delta^{\alpha_2\beta_2}\delta^{\alpha_3\beta_3}$	

Table 2: *When contracted with the transverse projectors, this table presents all* 24 *tensor structures in the decomposition of the transverse part of* $\langle\!\langle T^{\mu_1\nu_1} T^{\mu_2\nu_2} T^{\mu_3\nu_3} \rangle\!\rangle$. *Tensors are divided into* 10 *orbits of the action of the symmetry group* S_3, *after the contractions with the transverse projectors are taken.*

and we use the convention (3.7) (explained in detail in section 3.1) for the momenta appearing under the various Lorentz indices:

$$\boldsymbol{p}_1, \boldsymbol{p}_2 \text{ for } \mu_1, \nu_1;\ \boldsymbol{p}_2, \boldsymbol{p}_3 \text{ for } \mu_2, \nu_2 \text{ and } \boldsymbol{p}_3, \boldsymbol{p}_1 \text{ for } \mu_3, \nu_3. \tag{A.1.5}$$

The following table lists all 24 simple tensors from which $X^{\alpha_1\beta_1\alpha_2\beta_2\alpha_3\beta_3}$ may be built.

Contracting each tensor in the table with the transverse projectors we obtain 24 transverse tensors denoted by P_a, $a = 1, 2, \ldots, 24$. Each tensor P_a can then be multiplied by a form factor B_a to obtain the decomposition

$$\langle\!\langle t_T^{\mu_1\nu_1}(\boldsymbol{p}_1) t_T^{\mu_2\nu_2}(\boldsymbol{p}_2) t_T^{\mu_3\nu_3}(\boldsymbol{p}_3) \rangle\!\rangle = \sum_{a=1}^{24} B_a(p_1, p_2, p_3) P_a^{\mu_1\nu_1\mu_2\nu_2\mu_3\nu_3}. \tag{A.1.6}$$

However, the number of independent form factors may be reduced by looking at the symmetry properties. If we denote the permutation group of the set $\{1, 2, 3\}$ by S_3, then the 3-point function is S_3-invariant, *i.e.*, for any $\sigma \in S_3$,

$$\langle\!\langle t_T^{\mu_1\nu_1}(\boldsymbol{p}_1) t_T^{\mu_2\nu_2}(\boldsymbol{p}_2) t_T^{\mu_3\nu_3}(\boldsymbol{p}_3) \rangle\!\rangle = \langle\!\langle t_T^{\mu_{\sigma(1)}\nu_{\sigma(1)}}(\boldsymbol{p}_{\sigma(1)}) t_T^{\mu_{\sigma(2)}\nu_{\sigma(2)}}(\boldsymbol{p}_{\sigma(2)}) t_T^{\mu_{\sigma(3)}\nu_{\sigma(3)}}(\boldsymbol{p}_{\sigma(3)}) \rangle\!\rangle. \tag{A.1.7}$$

When contracted with the transverse projectors, the tensors at the first, fifth and the last row of the table lead to the S_3-invariant tensors. Therefore, corresponding form factors are invariant under any permutation of their arguments, for example

$$B_1(p_1, p_2, p_3) = B_1(p_{\sigma(1)}, p_{\sigma(2)}, p_{\sigma(3)}) \tag{A.1.8}$$

for any $\sigma \in S_3$. The remaining tensors transform non-trivially under the action of S_3. For concreteness, consider the second line of the table, *i.e.*, the part of the decomposition

$$B_2(p_1, p_2, p_3) P_2^{\mu_1\nu_1\mu_2\nu_2\mu_3\nu_3} + B_3(p_1, p_2, p_3) P_3^{\mu_1\nu_1\mu_2\nu_2\mu_3\nu_3} + B_4(p_1, p_2, p_3) P_4^{\mu_1\nu_1\mu_2\nu_2\mu_3\nu_3}. \tag{A.1.9}$$

Under the action of the symmetry group the tensors P_2, P_3, P_4 shuffle among each other. For example, under the action of the transposition $(\boldsymbol{p}_1, \mu_1, \nu_1) \leftrightarrow (\boldsymbol{p}_3, \mu_3, \nu_3)$ we obtain

$$B_2(p_3, p_2, p_1) P_3^{\mu_1\nu_1\mu_2\nu_2\mu_3\nu_3} + B_3(p_3, p_2, p_1) P_2^{\mu_1\nu_1\mu_2\nu_2\mu_3\nu_3} + B_4(p_3, p_2, p_1) P_4^{\mu_1\nu_1\mu_2\nu_2\mu_3\nu_3}. \tag{A.1.10}$$

Since the entire 3-point function is S_3-invariant, this implies that (A.1.9) and (A.1.10) are equal. Since all tensor structures P_a are independent, we find

$$B_3(p_1, p_2, p_3) = B_2(p_3, p_2, p_1), \qquad B_4(p_1, p_2, p_3) = B_4(p_3, p_2, p_1). \tag{A.1.11}$$

By analysing other symmetries we find that (A.1.9) depends on one form factor only, say B_2,

$$B_2(p_1,p_2,p_3)P_2^{\mu_1\nu_1\mu_2\nu_2\mu_3\nu_3} + B_2(p_1 \leftrightarrow p_3)P_3^{\mu_1\nu_1\mu_2\nu_2\mu_3\nu_3} + B_2(p_2 \leftrightarrow p_3)P_4^{\mu_1\nu_1\mu_2\nu_2\mu_3\nu_3}. \quad \text{(A.1.12)}$$

Moreover, $B_2(p_1,p_2,p_3) = B_2(p_1 \leftrightarrow p_2)$.

The described procedure reduces the number of independent form factors from 24 down to 10. The same procedure applied to the transverse-traceless part of the 3-point function reduces the number of independent tensors from 11 down to 5. In this case the decomposition is given by (3.13).

A.2 Degeneracy of the tensor structure of $\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}T^{\mu_3\nu_3}\rangle$ in $d = 3$

In dimension $d = 3$, a special degeneracy occurs which allows the transverse-traceless part of $\langle\!\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}T^{\mu_3\nu_3}\rangle\!\rangle$ to be decomposed in terms of only two form factors rather than five.

To see this, we first define the cross-product

$$\boldsymbol{n} = \boldsymbol{p}_1 \times \boldsymbol{p}_2 = \boldsymbol{p}_2 \times \boldsymbol{p}_3 = \boldsymbol{p}_3 \times \boldsymbol{p}_1 \quad \text{(A.2.1)}$$

and note that $n^2 = \lambda^2/4$, where λ^2 is defined in (6.21). Using (3.5) we find

$$\delta^{\mu\nu} = \frac{4}{\lambda^2}\left[p_i^2 p_j^\mu p_j^\nu + p_j^2 p_i^\mu p_i^\nu - \boldsymbol{p}_i\cdot\boldsymbol{p}_j(p_i^\mu p_j^\nu + p_j^\mu p_i^\nu) + n^\mu n^\nu\right] \quad \text{(A.2.2)}$$

for any $i,j = 1,2,3$ and $i \neq j$. From the fact that $\delta^{\alpha\beta}\Pi^{\mu\nu}_{\alpha\beta}(\boldsymbol{p}_j) = 0$, we find

$$\Pi^{\mu\nu}_{\alpha\beta}(\boldsymbol{p}_j)n^\alpha n^\beta = -p_j^2\Pi^{\mu\nu}_{\alpha\beta}(\boldsymbol{p}_j)\, p^\alpha_{(j+1)\bmod 3}\, p^\beta_{(j+1)\bmod 3}, \qquad j = 1,2,3. \quad \text{(A.2.3)}$$

We can now go back to the decomposition of the transverse-traceless part of $\langle\!\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}T^{\mu_3\nu_3}\rangle\!\rangle$, equation (R.11.5), and exchange all $\delta^{\alpha\beta}$ for (A.2.2). However, if one transverse-traceless projector is contracted with two vectors $\boldsymbol{n}$, then, according to (A.2.3), we can replace such a contraction with a contraction of two momenta with appropriate prefactors. Therefore, the only terms surviving in (R.11.5) are terms with either zero or two vectors $\boldsymbol{n}$. Hence we find only two tensor structures in the decomposition of $\langle\!\langle t^{\mu_1\nu_1}t^{\mu_2\nu_2}t^{\mu_3\nu_3}\rangle\!\rangle$,

$$\begin{aligned}
&\langle\!\langle t^{\mu_1\nu_1}(\boldsymbol{p}_1)t^{\mu_2\nu_2}(\boldsymbol{p}_2)t^{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\!\rangle \\
&\quad= \Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1)\Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)\Pi^{\mu_3\nu_3}_{\alpha_3\beta_3}(\boldsymbol{p}_3)\Big[B_1 p_2^{\alpha_1}p_2^{\beta_1}p_3^{\alpha_2}p_3^{\beta_2}p_1^{\alpha_3}p_1^{\beta_3} \\
&\qquad+\ B_2 n^{\beta_1}n^{\beta_2}p_2^{\alpha_1}p_3^{\alpha_2}p_1^{\alpha_3}p_1^{\beta_3} + B_2(p_1\leftrightarrow p_3)n^{\beta_2}n^{\beta_3}p_2^{\alpha_1}p_2^{\beta_1}p_3^{\alpha_2}p_1^{\alpha_3} \\
&\qquad\quad + B_2(p_2\leftrightarrow p_3)n^{\beta_1}n^{\beta_3}p_2^{\alpha_1}p_3^{\alpha_2}p_3^{\beta_2}p_1^{\alpha_3}\Big].
\end{aligned} \quad \text{(A.2.4)}$$

The new form factors B_j are functions of the momentum magnitudes. As usual, if no arguments are specified then the standard ordering is assumed, $B_j = B_j(p_1,p_2,p_3)$, while by $p_i \leftrightarrow p_j$ we denote the exchange of the two momenta, *e.g.*, $B_1(p_1 \leftrightarrow p_3) = B_2(p_3,p_2,p_1)$.

We can now express the new form factors B_j in terms of the old ones, A_j, defined in (R.11.5). Using equation (A.2.3), we write the explicit form of the contraction of two transverse-traceless projectors with a metric as

$$\Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1)\Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)\delta^{\beta_1\beta_2} = \frac{4}{\lambda^2}\Pi^{\mu_1\nu_1}_{\alpha_1\beta_1}(\boldsymbol{p}_1)\Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)\left[n^{\beta_1}n^{\beta_2} + \frac{1}{2}(p_3^2 - p_1^2 - p_2^2)p_2^{\beta_1}p_3^{\beta_2}\right], \quad \text{(A.2.5)}$$

from which we find

$$\begin{aligned}
B_1 = A_1 &+ \frac{2}{\lambda^2}\left[(p_3^2 - p_1^2 - p_2^2)A_2(p_1,p_2,p_3) + (p_1 \leftrightarrow p_3) + (p_2 \leftrightarrow p_3)\right] \\
&+ \frac{4}{\lambda^4}\left[\left((8p_1^2p_2^2 - \lambda^2)A_3 + (p_3^4 - (p_1^2 - p_2^2)^2)A_4\right) + (p_1 \leftrightarrow p_3) + (p_2 \leftrightarrow p_3)\right] \\
&- \frac{8}{\lambda^4}(p_1^2 + p_2^2 + p_3^2)A_5, \qquad \text{(A.2.6)} \\
B_2 = \frac{4}{\lambda^2}A_2 &+ \frac{16}{\lambda^4}\left[(p_3^2 - p_1^2 - p_2^2)A_3 - p_3^2A_4 + \right. \\
&\left. + \frac{1}{2}(p_2^2 - p_1^2 - p_3^2)A_4(p_1 \leftrightarrow p_3) + \frac{1}{2}(p_1^2 - p_2^2 - p_3^2)A_4(p_2 \leftrightarrow p_3)\right] \\
&+ \frac{16}{\lambda^4}A_5. \qquad \text{(A.2.7)}
\end{aligned}$$

Using the general expressions (R.11.29) - (R.11.33) for the form factors in $d = 3$, we arrive at the final result

$$\begin{aligned}
B_1 = 1920\alpha_1\frac{c_{123}^3}{\lambda^4 a_{123}^4} &- \frac{8c_T}{\lambda^4 a_{123}^2}\left[(3+8c_g)a_{123}^5(a_{123}^2 - 5b_{123}) + 24(1+2c_g)a_{123}^3b_{123}^2\right. \\
&\left. - 8a_{123}b_{123}^3 + \left(3(8c_g-1)a_{123}^4 - 48c_g a_{123}^2 b_{123} - 8b_{123}^2\right)c_{123} + 8a_{123}c_{123}^2\right], \qquad \text{(A.2.8)} \\
B_2 = -1920\alpha_1\frac{c_{123}^2 p_3}{\lambda^4 a_{123}^4} &+ \frac{64c_T}{\lambda^4 a_{123}^2}\left[2(1+c_g)p_3^4(p_3 + 2a_{12})\right. \\
&+ p_3^3\left(2(2+c_g)a_{12}^2 - 4b_{12}\right) + p_3^2 a_{12}\left((3+2c_g)a_{12}^2 - (5+6c_g)b_{12}\right) \\
&+ 2p_3\left((1+2c_g)a_{12}^2(a_{12}^2 - 3b_{12}) + b_{12}^2\right) \\
&\left. - 3(1+2c_g)a_{12}^3b_{12} + a_{12}b_{12}^2 + (1+2c_g)a_{12}^5\right]. \qquad \text{(A.2.9)}
\end{aligned}$$

The variables used in this expression are symmetric polynomials of the momentum magnitudes as defined in (R.1.2). Note that this expression has no dependence on the primary constant α_2. Therefore, in $d = 3$, the most general form of the correlation function $\langle\langle T^{\mu_1\nu_1}T^{\mu_2\nu_2}T^{\mu_3\nu_3}\rangle\rangle$ depends on only one undetermined primary constant and on two 2-point function normalisations c_T and c_g. This is in agreement with [3], noting that the normalisation constant c_g arises through our definition of the 3-point function in (4.47).

Finally, while similar considerations hold for other 3-point correlators in $d = 3$ involving the stress-energy tensor, in these cases it turns out that the use of equation (A.2.2) does not reduce the number of independent primary constants in the final result.

A.3 Triple-K integrals by Fourier transform

Here we present the computation leading from the momentum space integral in (2.4) to the triple-K representation. The method generalises the results of [31], as we consider a general integral

$$\int \frac{\mathrm{d}^d\boldsymbol{k}}{(2\pi)^d}\frac{k^{\mu_1}\dots k^{\mu_r}}{|\boldsymbol{k}|^{2\delta_3}|\boldsymbol{k}-\boldsymbol{p}_1|^{2\delta_2}|\boldsymbol{k}+\boldsymbol{p}_2|^{2\delta_1}}. \qquad \text{(A.3.1)}$$

Using Schwinger parameters

$$\frac{1}{A^\alpha} = \frac{1}{\Gamma(\alpha)}\int_0^\infty \mathrm{d}s\, s^{\alpha-1}e^{-sA}\mathrm{d}s, \qquad \alpha > 0. \qquad \text{(A.3.2)}$$

we can write

$$\begin{aligned}
&\int \frac{\mathrm{d}^d\boldsymbol{k}}{(2\pi)^d}\frac{k^{\mu_1}\dots k^{\mu_r}}{\boldsymbol{k}^{2\delta_3}|\boldsymbol{k}-\boldsymbol{p}_1|^{2\delta_2}|\boldsymbol{k}+\boldsymbol{p}_2|^{2\delta_1}}\\
&=\Gamma^{-3}\int\frac{\mathrm{d}^d\boldsymbol{k}}{(2\pi)^d}k^{\mu_1}\dots k^{\mu_r}\int_{\mathbb{R}^3_+}\mathrm{d}\vec{s}\,s_1^{\delta_1-1}s_2^{\delta_2-1}s_3^{\delta_3-1}\times\\
&\qquad\times\exp\left[-(s_3\boldsymbol{k}^2+s_2|\boldsymbol{k}-\boldsymbol{p}_1|^2+s_1|\boldsymbol{k}+\boldsymbol{p}_2|^2)\right],
\end{aligned}\tag{A.3.3}$$

where we use the following abbreviations

$$\Gamma^3=\Gamma(\delta_1)\Gamma(\delta_2)\Gamma(\delta_3),\qquad\qquad \mathrm{d}\vec{s}=\mathrm{d}s_1\mathrm{d}s_2\mathrm{d}s_3.\tag{A.3.4}$$

Denoting $s_t=s_1+s_2+s_3$, we rewrite the expression in the exponent as

$$s_3\boldsymbol{k}^2+s_2|\boldsymbol{k}-\boldsymbol{p}_1|^2+s_1|\boldsymbol{k}+\boldsymbol{p}_2|^2=s_tl^2+\Delta,\tag{A.3.5}$$

where

$$\boldsymbol{l}=\boldsymbol{k}+\frac{s_1\boldsymbol{p}_2-s_2\boldsymbol{p}_1}{s_t},\qquad\qquad \Delta=\frac{s_1s_2p_3^2+s_1s_3p_2^2+s_2s_3p_1^2}{s_t}.\tag{A.3.6}$$

We can now re-express the integral (A.3.1) as

$$\Gamma^{-3}\int_{\mathbb{R}^3_+}\mathrm{d}\vec{s}\,s_1^{\delta_1-1}s_2^{\delta_2-1}s_3^{\delta_3-1}e^{-\Delta}\int\frac{\mathrm{d}^d\boldsymbol{l}}{(2\pi)^d}e^{-s_tl^2}\prod_{j=1}^{r}\left(l^{\mu_j}+\frac{s_2p_1^{\mu_j}-s_1p_2^{\mu_j}}{s_t}\right).\tag{A.3.7}$$

This expression can be expanded and split up into a sum of integrals. The integral over $\boldsymbol{l}$ gives some moment of a Gaussian random variable. For any a such that $\mathrm{Re}\,a>0$ we can find

$$\int\frac{\mathrm{d}^d\boldsymbol{l}}{(2\pi)^d}l^{2m}e^{-al^2}=\frac{\Gamma\left(\frac{d}{2}+m\right)}{(4\pi)^{\frac{d}{2}}\Gamma\left(\frac{d}{2}\right)}\cdot\frac{1}{a^{\frac{d}{2}+m}},\tag{A.3.8}$$

$$\int\frac{\mathrm{d}^d\boldsymbol{l}}{(2\pi)^d}l^{\mu_1}\dots l^{\mu_{2m}}e^{-al^2}=\frac{S^{\mu_1\dots\mu_{2m}}}{(4\pi)^{\frac{d}{2}}2^ma^{\frac{d}{2}+m}},\tag{A.3.9}$$

and integrals with an odd number of l vanish. $S^{\mu_1\dots\mu_{2m}}$ is a completely symmetric tensor built from metrics only, with each coefficient equal to 1, *e.g.*,

$$S^{\mu_1\mu_2\mu_3\mu_4}=\delta^{\mu_1\mu_2}\delta^{\mu_3\mu_4}+\delta^{\mu_1\mu_3}\delta^{\mu_2\mu_4}+\delta^{\mu_1\mu_4}\delta^{\mu_2\mu_3}.\tag{A.3.10}$$

The calculations of the integral (A.3.1) therefore boil down to the calculation of several integrals of the form

$$i_{d,m,\{\delta_j\}}=\frac{1}{(4\pi)^{\frac{d}{2}}2^m\Gamma^3}\int_{\mathbb{R}^3_+}\mathrm{d}\vec{s}\,s_t^{-\frac{d}{2}-m}s_1^{\delta_1-1}s_2^{\delta_2-1}s_3^{\delta_3-1}e^{-\Delta}.\tag{A.3.11}$$

This expression gives the coefficient of the completely symmetric tensor $S^{\mu_1\dots\mu_{2m}}$ when we evaluate (A.3.7). In fact, the coefficient of all the tensors in (A.3.7) can similarly be expressed in terms of $i_{d,m,\{\delta_j\}}$ for some values of m and δ_j. (In this case, however, the δ_j parameters are no longer equal to those in (A.3.1), since each momentum in (A.3.7) is accompanied by a Schwinger parameter.)

Let us now express the integral (A.3.11) in terms of the triple-K integral (5.1). Defining $\delta_t=\delta_1+\delta_2+\delta_3$, we make the following substitution in (A.3.11)

$$s_j=\frac{v_1v_2+v_1v_3+v_2v_3}{2v_j}=\frac{V}{2v_j},\qquad j=1,2,3,\tag{A.3.12}$$

giving

$$i_{d,m,\{\delta_j\}} = \frac{2^{\frac{d}{2}-\delta_t}}{(4\pi)^{\frac{d}{2}}\Gamma^3}\int_{\mathbb{R}^3_+} \mathrm{d}\vec{v}\, V^{\delta_t-d-2m}\prod_{j=1}^{3} v_j^{\frac{d}{2}+m-\delta_j-1} e^{-\frac{v_j p_j^2}{2}}. \tag{A.3.13}$$

Observing that

$$V = v_1v_2v_3\left(v_1^{-1}+v_2^{-1}+v_3^{-1}\right) \tag{A.3.14}$$

and introducing a new Schwinger parameter t to exponentiate the term in brackets, we find

$$\begin{aligned} i_{d,m,\{\delta_j\}} &= \frac{2^{\frac{d}{2}-\delta_t}}{(4\pi)^{\frac{d}{2}}\Gamma^3\Gamma(d+2m-\delta_t)}\int_0^\infty \mathrm{d}t\, t^{d+2m-\delta_t-1}\times \\ &\quad\times\int_{\mathbb{R}^3_+}\mathrm{d}\vec{v}\prod_{j=1}^{3} v_j^{-\frac{d}{2}-m+\delta_t-\delta_j-1} e^{-\frac{v_j p_j^2}{2}-\frac{t}{v_j}} \\ &= \frac{2^{-d-3m+\delta_t}}{(4\pi)^{\frac{d}{2}}\Gamma^3\Gamma(d+2m-\delta_t)}\int_0^\infty \mathrm{d}t\, t^{d+2m-\delta_t-1}\times \\ &\quad\times\int_{\mathbb{R}^3_+}\mathrm{d}\vec{u}\prod_{j=1}^{3} p_j^{d+2m-2\delta_t+2\delta_j} u_j^{-\frac{d}{2}-m+\delta_t-\delta_j-1} e^{-u_j-\frac{tp_j^2}{2u_j}}. \end{aligned} \tag{A.3.15}$$

Using the standard formula [38]

$$K_\nu(z) = \frac{1}{2}\left(\frac{z}{2}\right)^\nu\int_0^\infty e^{-u-\frac{z^2}{4u}}u^{-\nu-1}\mathrm{d}u, \qquad |\arg z| < \frac{\pi}{4} \tag{A.3.16}$$

we now obtain our final result

$$\begin{aligned} i_{d,m,\{\delta_j\}} &= \frac{2^{-\frac{d}{2}-2m+4}}{(4\pi)^{\frac{d}{2}}\Gamma^3\Gamma(d+2m-\delta_t)}\int_0^\infty \mathrm{d}t(\sqrt{2t})^{\frac{d}{2}+m-2}\prod_{j=1}^{3} p_j^{\frac{d}{2}+m-\delta_t+\delta_j} K_{\frac{d}{2}+m-\delta_t+\delta_j}(\sqrt{2t}p_j) \\ &= \frac{2^{-\frac{d}{2}-2m+4}}{(4\pi)^{\frac{d}{2}}\Gamma^3\Gamma(d+2m-\delta_t)}\int_0^\infty \mathrm{d}x\, x^{\frac{d}{2}+m-1}\prod_{j=1}^{3} p_j^{\frac{d}{2}+m-\delta_t+\delta_j} K_{\frac{d}{2}+m-\delta_t+\delta_j}(p_j x) \\ &= \frac{2^{-\frac{d}{2}-2m+4}}{(4\pi)^{\frac{d}{2}}\Gamma^3\Gamma(d+2m-\delta_t)} I_{\frac{d}{2}+m-1\{\frac{d}{2}+m-\delta_t+\delta_j\}}, \end{aligned} \tag{A.3.17}$$

where $\Gamma^3 = \Gamma(\delta_1)\Gamma(\delta_2)\Gamma(\delta_3)$ and I stands for the triple-K integral (5.1).

A.3.1 Triple-K integrals via Feynman parametrisation

Instead of Schwinger parameters, one can use the more familiar Feynman parametrisation in order to evaluate (A.3.1). Using standard results [53], we can write (A.3.1) as

$$\frac{\Gamma(\delta_t)}{\Gamma^3}\int_{[0,1]^3}\mathrm{d}X\prod_{j=1}^{3} x_j^{\delta_j-1}\int\frac{\mathrm{d}^d\boldsymbol{l}}{(2\pi)^d}\frac{1}{(l^2+D)^{\delta_t}}\prod_{j=1}^{r}\left(l^{\mu_j}+x_2p_1^{\mu_j}-x_1p_2^{\mu_j}\right), \tag{A.3.18}$$

where

$$\mathrm{d}X = \mathrm{d}x_1\mathrm{d}x_2\mathrm{d}x_3\,\delta(x_1+x_2+x_3-1), \tag{A.3.19}$$

$$\boldsymbol{l} = \boldsymbol{k}-x_2\boldsymbol{p}_1+x_1\boldsymbol{p}_2, \tag{A.3.20}$$

$$D = p_1^2x_2x_3+p_2^2x_1x_3+p_3^2x_1x_2. \tag{A.3.21}$$

Looking at the coefficient of the tensor $S^{\mu_1\cdots\mu_{2m}}$ defined in the previous section, we obtain

$$i_{d,m,\{\delta_j\}} = \frac{\Gamma\left(\delta_t - m - \frac{d}{2}\right)}{(4\pi)^{\frac{d}{2}} 2^m \Gamma^3} \int_{[0,1]^3} \mathrm{d}X x_1^{\delta_1-1} x_2^{\delta_2-1} x_3^{\delta_3-1} D^{\frac{d}{2}+m-\delta_t}. \tag{A.3.22}$$

Comparing with (A.3.17), we then find

$$\begin{aligned} I_{\alpha\{\beta_1\beta_2\beta_3\}} &= \int_0^\infty \mathrm{d}x\, x^\alpha \prod_{j=1}^3 p_j^{\beta_j} K_{\beta_j}(p_j x) \\ &= 2^{\alpha-3}\Gamma\left(\frac{\alpha-\beta_t+1}{2}\right)\Gamma\left(\frac{\alpha+\beta_t+1}{2}\right)\times \\ &\quad \times \int_{[0,1]^3} \mathrm{d}X\, D^{\frac{1}{2}(\beta_t-\alpha-1)} \prod_{j=1}^3 x_j^{\frac{1}{2}(\alpha-1-\beta_t)+\beta_j}, \end{aligned} \tag{A.3.23}$$

where $\beta_t = \beta_1+\beta_2+\beta_3$ and I is the triple-K integral (5.1).

A.4 Properties of triple-K integrals

In this appendix we list some properties of modified Bessel functions used in the main text. For further references, see *e.g.*, [38].

The Bessel function I (modified Bessel function of the first kind) is given by the series

$$I_\nu(x) = \sum_{j=0}^\infty \frac{1}{j!\Gamma(\nu+j+1)}\left(\frac{x}{2}\right)^{\nu+2j}, \quad \nu \neq -1,-2,-3,\ldots \tag{A.4.1}$$

The Bessel function K (modified Bessel function of the second kind) is defined by

$$K_\nu(x) = \frac{\pi}{2\sin(\nu\pi)}\left[I_{-\nu}(x) - I_\nu(x)\right], \quad \nu \notin \mathbb{Z}, \tag{A.4.2}$$

$$K_n(x) = \lim_{\epsilon\to 0} K_{n+\epsilon}(x), \quad n \in \mathbb{Z}. \tag{A.4.3}$$

The finite pointwise limit for $x > 0$ exists for any integer n. K_ν is an even function of ν, *i.e.*, $K_{-\nu}(x) = K_\nu(x)$ for any $\nu \in \mathbb{R}$. If $\nu = \frac{1}{2} + n$, for an integer n, the Bessel function reduces to elementary functions

$$K_\nu(x) = \sqrt{\frac{\pi}{2}}\frac{e^{-x}}{\sqrt{x}} \sum_{j=0}^{\lfloor |\nu|-\frac{1}{2}\rfloor} \frac{\left(|\nu|-\frac{1}{2}+j\right)!}{j!\left(|\nu|-\frac{1}{2}-j\right)!}\frac{1}{(2x)^j}, \quad \nu+\frac{1}{2} \in \mathbb{Z}, \tag{A.4.4}$$

and in particular

$$\begin{aligned} K_{\frac{1}{2}}(x) &= \sqrt{\frac{\pi}{2}}\frac{e^{-x}}{x^{\frac{1}{2}}}, & K_{\frac{3}{2}}(x) &= \sqrt{\frac{\pi}{2}}\frac{e^{-x}}{x^{\frac{3}{2}}}(1+x), \\ K_{\frac{5}{2}}(x) &= \sqrt{\frac{\pi}{2}}\frac{e^{-x}}{x^{\frac{5}{2}}}(x^2+3x+3), & K_{\frac{7}{2}}(x) &= \sqrt{\frac{\pi}{2}}\frac{e^{-x}}{x^{\frac{7}{2}}}(x^3+6x^2+15x+5). \end{aligned} \tag{A.4.5}$$

The series expansion of the Bessel function K_ν for $\nu \notin \mathbb{Z}$ is given directly in terms of the expansion (A.4.1) via the definition (A.4.2). In particular

$$K_\nu(x) = \left[\Gamma(-\nu)2^{-\nu-1}x^\nu + O(x^{2-\nu})\right] + \left[\frac{\Gamma(\nu)2^{\nu-1}}{x^\nu} + O(x^{2+\nu})\right], \quad \nu \notin \mathbb{Z}. \tag{A.4.6}$$

For integer index $n \in \mathbb{N}$, the expansion reads

$$\begin{aligned} K_n(x) = \frac{1}{2}\left(\frac{x}{2}\right)^{-n} \sum_{j=0}^{n-1} \frac{(n-j-1)!}{j!}(-1)^j \left(\frac{x}{2}\right)^{2j} \\ + \ (-1)^{n+1} \log\left(\frac{x}{2}\right) I_n(x) \\ + \ (-1)^n \frac{1}{2}\left(\frac{x}{2}\right)^n \sum_{j=0}^{\infty} \frac{\psi(j+1)+\psi(n+j+1)}{j!(n+j)!}\left(\frac{x}{2}\right)^{2j}, \end{aligned} \tag{A.4.7}$$

where ψ is the digamma function. At large x, the Bessel functions have the asymptotic expansions

$$I_\nu(x) = \frac{1}{\sqrt{2\pi}}\frac{e^x}{\sqrt{x}} + \ldots, \qquad K_\nu(x) = \sqrt{\frac{\pi}{2}}\frac{e^{-x}}{\sqrt{x}} + \ldots, \quad \nu \in \mathbb{R}. \tag{A.4.8}$$

For any index $\nu \in \mathbb{R}$, the Bessel function K satisfies the following identities

$$\frac{\partial}{\partial a}\left[a^\nu K_\nu(ax)\right] = -x a^\nu K_{\nu-1}(ax), \tag{A.4.9}$$

$$K_{\nu-1}(x) + \frac{2\nu}{x}K_\nu(x) = K_{\nu+1}(x), \tag{A.4.10}$$

$$K_{-\nu}(x) = K_\nu(x). \tag{A.4.11}$$

A.5 Appell's F_4 function

Appell's F_4 function can be defined by the following double series [39, 40]

$$F_4(\alpha,\beta;\gamma,\gamma';\xi,\eta) = \sum_{i,j=0}^{\infty} \frac{(\alpha)_{i+j}(\beta)_{i+j}}{(\gamma)_i(\gamma')_j i!j!}\xi^i\eta^j, \quad \sqrt{|\xi|}+\sqrt{|\eta|}<1, \tag{A.5.1}$$

where $(\alpha)_i$ is a Pochhammer symbol. Notice that

$$F_4(\alpha,\beta;\gamma,\gamma';\xi,\eta) = F_4(\beta,\alpha;\gamma,\gamma';\xi,\eta) = F_4(\alpha,\beta;\gamma',\gamma;\eta,\xi). \tag{A.5.2}$$

The series representation, however, is not very useful as in our case

$$\xi = \frac{p_1^2}{p_3^2}, \qquad \eta = \frac{p_2^2}{p_3^2} \tag{A.5.3}$$

and the series converges when $p_3 > p_1 + p_2$, which is opposite to the triangle inequality.

As in the case of ordinary hypergeometric functions, the F_4 function satisfies certain differential equations. Let $\alpha, \beta, \gamma, \gamma'$ be fixed numbers. The following system of equations

$$\begin{aligned} 0 = \Bigg[\xi(1-\xi)\frac{\partial^2}{\partial\xi^2} - \eta^2\frac{\partial^2}{\partial\eta^2} - 2\xi\eta\frac{\partial^2}{\partial\xi\partial\eta} \\ + \ (\gamma - (\alpha+\beta+1)\xi)\frac{\partial}{\partial\xi} - (\alpha+\beta+1)\eta\frac{\partial}{\partial\eta} - \alpha\beta\Bigg] F(\xi,\eta), \end{aligned} \tag{A.5.4}$$

$$\begin{aligned} 0 = \Bigg[\eta(1-\eta)\frac{\partial^2}{\partial\eta^2} - \xi^2\frac{\partial^2}{\partial\xi^2} - 2\xi\eta\frac{\partial^2}{\partial\xi\partial\eta} \\ + \ \left(\gamma' - (\alpha+\beta+1)\eta\right)\frac{\partial}{\partial\eta} - (\alpha+\beta+1)\xi\frac{\partial}{\partial\xi} - \alpha\beta\Bigg] F(\xi,\eta), \end{aligned} \tag{A.5.5}$$

has exactly four solutions given by [40, 41]

$$F_4(\alpha,\beta;\gamma,\gamma';\xi,\eta), \tag{A.5.6}$$
$$\xi^{1-\gamma}F_4(\alpha+1-\gamma,\beta+1-\gamma;2-\gamma,\gamma';\xi,\eta), \tag{A.5.7}$$
$$\eta^{1-\gamma'}F_4(\alpha+1-\gamma',\beta+1-\gamma';\gamma,2-\gamma';\xi,\eta), \tag{A.5.8}$$
$$\xi^{1-\gamma}\eta^{1-\gamma'}F_4(\alpha+2-\gamma-\gamma',\beta+2-\gamma-\gamma';2-\gamma,2-\gamma';\xi,\eta). \tag{A.5.9}$$

The following reduction formulae can be found in [42] or [40]

$$F_4\left(\alpha,\beta;\alpha,\beta;-\frac{x}{(1-x)(1-y)},-\frac{y}{(1-x)(1-y)}\right) = \\ = \frac{(1-x)^\beta(1-y)^\alpha}{1-xy}, \tag{A.5.10}$$
$$F_4\left(\alpha,\beta;\beta,\beta;-\frac{x}{(1-x)(1-y)},-\frac{y}{(1-x)(1-y)}\right) = \\ = (1-x)^\alpha(1-y)^\alpha{}_2F_1(\alpha,1+\alpha-\beta;\beta;xy), \tag{A.5.11}$$
$$F_4\left(\alpha,\beta;1+\alpha-\beta,\beta;-\frac{x}{(1-x)(1-y)},-\frac{y}{(1-x)(1-y)}\right) = \\ = (1-y)^\alpha{}_2F_1\left(\alpha,\beta;1+\alpha-\beta;-\frac{x(1-y)}{1-x}\right), \tag{A.5.12}$$
$${}_2F_1(2\nu-1,\nu;\nu;x) = (1-x)^{1-2\nu}. \tag{A.5.13}$$

A.5.1 Integrals

Here we present the list of integrals we use in the paper, which may be found in [42].

(i)

$$\int_0^\infty \mathrm{d}x\, x^{\alpha-1}I_\lambda(ax)I_\mu(bx)K_\nu(cx) = \\ = \frac{2^{\alpha-2}\Gamma\left(\frac{\alpha+\lambda+\mu-\nu}{2}\right)\Gamma\left(\frac{\alpha+\lambda+\mu+\nu}{2}\right)}{\Gamma(\lambda+1)\Gamma(\mu+1)}\cdot\frac{a^\lambda b^\mu}{c^{\alpha+\lambda+\mu}}\times \\ \times F_4\left(\frac{\alpha+\lambda+\mu-\nu}{2},\frac{\alpha+\lambda+\mu+\nu}{2};\lambda+1,\mu+1;\frac{a^2}{c^2},\frac{b^2}{c^2}\right), \tag{A.5.14}$$

valid for

$$\mathrm{Re}(\alpha+\lambda+\mu) > |\,\mathrm{Re}\,\nu|, \qquad |c| > |a|+|b|, \qquad \mathrm{Re}\,c > |\,\mathrm{Re}\,a| + |\,\mathrm{Re}\,b|. \tag{A.5.15}$$

(ii)

$$\int_0^\infty \mathrm{d}x\, x^{\alpha-1}K_\lambda(ax)K_\mu(bx)K_\nu(cx) = \\ = \frac{2^{\alpha-4}}{c^\alpha}\left[A(\lambda,\mu)+A(\lambda,-\mu)+A(-\lambda,\mu)+A(-\lambda,-\mu)\right], \tag{A.5.16}$$

where

$$A(\lambda,\mu) = \left(\frac{a}{c}\right)^\lambda\left(\frac{b}{c}\right)^\mu\Gamma\left(\frac{\alpha+\lambda+\mu-\nu}{2}\right)\Gamma\left(\frac{\alpha+\lambda+\mu+\nu}{2}\right)\Gamma(-\lambda)\Gamma(-\mu)\times \\ \times F_4\left(\frac{\alpha+\lambda+\mu-\nu}{2},\frac{\alpha+\lambda+\mu+\nu}{2};\lambda+1,\mu+1;\frac{a^2}{c^2},\frac{b^2}{c^2}\right), \tag{A.5.17}$$

valid for

$$\mathrm{Re}\,\alpha > |\,\mathrm{Re}\,\lambda| + |\,\mathrm{Re}\,\mu| + |\,\mathrm{Re}\,\nu|, \qquad \mathrm{Re}(a+b+c) > 0. \tag{A.5.18}$$

(iii)

$$\int_0^\infty \mathrm{d}x\, x^{\alpha-1} K_\mu(cx) K_\nu(cx) =$$
$$= \frac{2^{\alpha-3}}{\Gamma(\alpha) c^\alpha} \Gamma\left(\frac{\alpha+\mu+\nu}{2}\right) \Gamma\left(\frac{\alpha+\mu-\nu}{2}\right) \Gamma\left(\frac{\alpha-\mu+\nu}{2}\right) \Gamma\left(\frac{\alpha-\mu-\nu}{2}\right), \quad \text{(A.5.19)}$$

valid for

$$\mathrm{Re}\,\alpha > |\,\mathrm{Re}\,\mu| + |\,\mathrm{Re}\,\nu|, \qquad \mathrm{Re}\,c > 0. \quad \text{(A.5.20)}$$

(iv)

$$\int_0^\infty \mathrm{d}x\, x^{\alpha-1} K_\nu(cx) = \frac{2^{\alpha-2}}{c^\alpha} \Gamma\left(\frac{\alpha+\nu}{2}\right) \Gamma\left(\frac{\alpha-\nu}{2}\right) \quad \text{(A.5.21)}$$

valid for

$$\mathrm{Re}\,\alpha > |\,\mathrm{Re}\,\nu|, \qquad \mathrm{Re}\,c > 0. \quad \text{(A.5.22)}$$

(v)

$$\int_0^\infty \mathrm{d}x\, x^{\alpha-1} \log x K_\nu(cx) = \frac{2^{\alpha-3}}{c^\alpha} \Gamma\left(\frac{\alpha+\nu}{2}\right) \Gamma\left(\frac{\alpha-\nu}{2}\right) \times$$
$$\times \left[\psi\left(\frac{\alpha+\nu}{2}\right) + \psi\left(\frac{\alpha-\nu}{2}\right) - 2\log\frac{c}{2}\right], \quad \text{(A.5.23)}$$

valid for

$$\mathrm{Re}\,\alpha > |\,\mathrm{Re}\,\nu|, \qquad \mathrm{Re}\,c > 0. \quad \text{(A.5.24)}$$

(vi)

$$\int_0^\infty \mathrm{d}x\, x^{\alpha-1} \log^2 x K_\nu(cx) = \frac{2^{\alpha-4}}{c^\alpha} \Gamma\left(\frac{\alpha+\nu}{2}\right) \Gamma\left(\frac{\alpha-\nu}{2}\right) \times$$
$$\times \left[\left(\psi\left(\frac{\alpha+\nu}{2}\right) + \psi\left(\frac{\alpha-\nu}{2}\right)\right) \cdot \left(\psi\left(\frac{\alpha+\nu}{2}\right) + \psi\left(\frac{\alpha-\nu}{2}\right) - 4\log\frac{c}{2}\right)\right.$$
$$\left. + \psi'\left(\frac{\alpha+\nu}{2}\right) + \psi'\left(\frac{\alpha-\nu}{2}\right) + 4\log^2\frac{c}{2}\right], \quad \text{(A.5.25)}$$

valid for

$$\mathrm{Re}\,\alpha > |\,\mathrm{Re}\,\nu|, \qquad \mathrm{Re}\,c > 0. \quad \text{(A.5.26)}$$

A.6 Master integral

The master integral is

$$\begin{aligned}
I_{0+\epsilon\{111\}} &= -\frac{p_1^2+p_2^2+p_3^2}{2\,\epsilon^2} \\
&+ \frac{1}{2\epsilon}\left[-h_{1/2}(p_1^2+p_2^2+p_3^2)+(p_1^2\log p_1+p_2^2\log p_2+p_3^2\log p_3)\right] \\
&+ \left[\frac{p_1p_2Z}{16}\left(\tfrac{1}{2}F^{(2)}(Z^2)+2(h_1+2\log p_3)F^{(1)}(Z^2)+(h_1+2\log^2 p_3)^2+1+\tfrac{1}{2}\pi^2\right)\right. \\
&\quad \left.+\left(p_1\leftrightarrow p_3, Z\mapsto X \text{ but } Z^2\mapsto Z\frac{X}{Y}\right)+\left(p_2\leftrightarrow p_3, Z\mapsto Y \text{ but } Z^2\mapsto Z\frac{Y}{X}\right)\right] \\
&\quad - \frac{\sqrt{-\lambda^2}}{8}\left[\log^2(p_1p_2p_3)+h_2\log(p_1p_2p_3)+\tfrac{1}{4}h_2^2+1+\tfrac{7}{24}\pi^2\right. \\
&\qquad \left.-(h_2+2\log(p_1p_2p_3))\log\sqrt{-\lambda^2}+\log^2\sqrt{-\lambda^2}\right] \\
&\quad + \frac{1}{8}\left[\left(p_3^2(-\tfrac{7}{4}h_{2/7}^2+\tfrac{8}{7}+\tfrac{1}{8}\pi^2)+(h_2(p_1^2+p_2^2)+3h_0p_3^2)\log p_3\right.\right. \\
&\qquad \left.\left.+(p_1^2+p_2^2-p_3^2)(\log^2 p_3-2\log p_1\log p_2)\right)+(p_1\leftrightarrow p_3)+(p_2\leftrightarrow p_3)\right] \\
&+ O(\epsilon),
\end{aligned} \tag{A.6.1}$$

where

$$F^{(1)}(x) = 1-\left(1-\frac{1}{x}\right)\log(1-x), \tag{A.6.2}$$

$$F^{(2)}(x) = 2+\left(1-\frac{1}{x}\right)\left[-\log(1-x)+\log^2(1-x)+\mathrm{Li}_2\,x\right] \tag{A.6.3}$$

are coefficients of the series expansion

$${}_2F_1(1,\epsilon;2-\epsilon;x) = 1+F^{(1)}(x)\epsilon+F^{(2)}(x)\epsilon^2+O(\epsilon^3). \tag{A.6.4}$$

We have also defined

$$\lambda^2 = (p_1+p_2-p_3)(p_1-p_2+p_3)(-p_1+p_2+p_3)(p_1+p_2+p_3), \tag{A.6.5}$$

$$X = \frac{p_1^2-p_2^2-p_3^2+\sqrt{-\lambda^2}}{2p_2p_3}, \qquad Y = \frac{p_2^2-p_1^2-p_3^2+\sqrt{-\lambda^2}}{2p_1p_3}, \tag{A.6.6}$$

$$Z = \frac{p_3^2-p_1^2-p_2^2-\sqrt{-\lambda^2}}{2p_1p_2} \tag{A.6.7}$$

and the constant

$$h_\alpha = \alpha-\gamma_E+\log 2. \tag{A.6.8}$$

The master integral can be evaluated by the method presented in section 6.4. All remaining integrals required for the calculations of 3-point functions of conserved currents and the stress-energy tensor in any even dimension $d \geq 4$ can be obtained from the master integral (A.6.1) by the reduction scheme. For $d = 4$, the reduction scheme is presented in table 1 on page 44.

Below we present the expressions for two other integrals, $I_{1\{222\}}$ and $I_{1\{000\}}$, appearing in table 1 on page 44. Both integrals can be obtained by the method described in section 6.4, but the resulting expressions are simpler than the master integral $I_{0\{111\}}$. Therefore, whenever possible, it is convenient to use them as a starting point in the reduction scheme. In particular the integral $I_{1\{000\}}$ is convergent and known in the literature, *e.g.*, [33, 49]

$$\begin{aligned}
I_{1\{000\}} = \frac{1}{2\sqrt{-\lambda^2}}\left[\frac{\pi^2}{6}-2\log\frac{p_1}{p_3}\log\frac{p_2}{p_3}+\log\left(-X\frac{p_2}{p_3}\right)\log\left(-Y\frac{p_1}{p_3}\right)\right. \\
\left.-\mathrm{Li}_2\left(-X\frac{p_2}{p_3}\right)-\mathrm{Li}_2\left(-Y\frac{p_1}{p_3}\right)\right].
\end{aligned} \tag{A.6.9}$$

From this integral one can find

$$I_{4\{111\}} = -p_1p_2p_3\frac{\partial^3}{\partial p_1\partial p_2\partial p_3}I_{1\{000\}}, \tag{A.6.10}$$

which is the top rightmost entry in table 1, page 44.

The second integral is

$$\begin{aligned} I_{2+\epsilon\{111\}} = \frac{1}{\epsilon} - \Bigg[& \frac{p_3^2 Z}{2p_1p_2}\Big(\tilde{F}^{(1)}(Z^2) + (h_0 + 2\log p_3)\tilde{F}^{(0)}(Z^2) \\ & + (2\log^2 p_3 + 2h_0\log p_3 + \tfrac{1}{2}h_0^2 + \tfrac{1}{4}\pi^2)\tilde{F}^{(-1)}(Z^2)\Big) \\ & + \left(p_1 \leftrightarrow p_3, Z \mapsto X \text{ but } Z^2 \mapsto Z\frac{X}{Y}\right) + \left(p_2 \leftrightarrow p_3, Z \mapsto Y \text{ but } Z^2 \mapsto Z\frac{Y}{X}\right)\Bigg] \\ & + \frac{2p_1^2p_2^2p_3^2}{(\sqrt{-\lambda^2})^3}\left[\log^2(p_1p_2p_3) + h_2\log(p_1p_2p_3) + \tfrac{1}{4}h_2^2 - 1 + \tfrac{7}{24}\pi^2\right. \\ & \left. - (h_2 + 2\log(p_1p_2p_3))\log\sqrt{-\lambda^2} + \log^2\sqrt{-\lambda^2}\right] + \frac{3}{2}h_0 \\ & + O(\epsilon), \end{aligned} \tag{A.6.11}$$

where

$$\tilde{F}^{(-1)}(x) = \frac{2x}{(x-1)^3}, \tag{A.6.12}$$

$$\tilde{F}^{(0)}(x) = \frac{1}{(x-1)^3}\left[-1 + x(4+x) - 4x\log(1-x)\right], \tag{A.6.13}$$

$$\tilde{F}^{(1)}(x) = \frac{4x}{(x-1)^3}\left[-2\log(1-x) + \log^2(1-x) + \mathrm{Li}_2\, x\right] \tag{A.6.14}$$

are coefficients of the series expansion

$${}_2F_1(1, 2+\epsilon; -\epsilon; x) = \frac{\tilde{F}^{(-1)}(x)}{\epsilon} + \tilde{F}^{(0)}(x) + \tilde{F}^{(1)}(x)\epsilon + O(\epsilon^2). \tag{A.6.15}$$

Note that both $I_{1\{000\}}$ and $I_{1\{222\}}$ can be obtained from $I_{0\{111\}}$,

$$I_{1\{000\}} = \frac{1}{2p_1p_2p_3}\left[p_1\frac{\partial^2}{\partial p_2\partial p_3} + p_2\frac{\partial^2}{\partial p_1\partial p_3} + p_3\frac{\partial^2}{\partial p_1\partial p_2}\right] I_{0+\epsilon\{111\}}, \tag{A.6.16}$$

$$I_{2+\epsilon\{111\}} = \left[\frac{\partial^2}{\partial p_1^2} - \frac{1}{p_1}\frac{\partial}{\partial p_1}\right] I_{0+\epsilon\{111\}}. \tag{A.6.17}$$

A.7 Triviality of $\langle T^{\mu_1\nu_1} J^{\mu_2}\mathcal{O}\rangle$

As our analysis shows, for any $d \geq 3$ and Δ_3 satisfying unitarity bound the correlation functions $\langle\!\langle T^{\mu_1\nu_1} J^{\mu_2}\mathcal{O}\rangle\!\rangle$ and $\langle\!\langle T^{\mu_1\nu_1} T^{\mu_2\nu_2} J^{\mu_3}\rangle\!\rangle$ are trivial, *i.e.*, they are at most semi-local. The triviality of $\langle\!\langle T^{\mu_1\nu_1} T^{\mu_2\nu_2} J^{\mu_3}\rangle\!\rangle$ was proved in [6] through a position space analysis. Our results independently confirm this through calculations in momentum space. In this section we discuss the triviality of $\langle\!\langle T^{\mu_1\nu_1} J^{\mu_2}\mathcal{O}\rangle\!\rangle$ as an example.

The tensor decomposition of the transverse-traceless part of the $\langle\!\langle T^{\mu_1\nu_1} J^{\mu_2}\mathcal{O}\rangle\!\rangle$ correlator, the primary and secondary CWIs, and the transverse Ward identities are presented along with all 3-point functions in part II of the paper. Let us rewrite here the important data. The solution in terms of triple-K integrals is

$$A_1^{aI} = \alpha_1^{aI} J_{3\{000\}}, \tag{A.7.1}$$

$$A_2^{aI} = 2\alpha_1^{aI} J_{2\{001\}} + \alpha_2^{aI} J_{1\{000\}}. \tag{A.7.2}$$

The independent secondary CWIs are

$$\mathrm{L}_{2,0}\, A_1^{aI} + \mathrm{R}_2\, A_2^{aI} = 2d \cdot \text{coefficient of } p_2^{\mu_1} p_3^{\mu_2} \text{ in } p_{1\nu_1} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2 a}(\boldsymbol{p}_2) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle, \tag{A.7.3}$$

$$\begin{aligned}&\mathrm{L}'_{1,0}\, A_1^{aI} + 2\,\mathrm{R}'_1\, A_2^{aI} \\ &\quad = -2(d-2) \cdot \text{coefficient of } p_2^{\mu_1} p_2^{\nu_1} \text{ in } p_{2\mu_2} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2 a}(\boldsymbol{p}_2) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle,\end{aligned} \tag{A.7.4}$$

$$\mathrm{L}_{2,0}\, A_2^{aI} = 4d \cdot \text{coefficient of } \delta^{\mu_1\mu_2} \text{ in } p_{1\nu_1} \langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2 a}(\boldsymbol{p}_2) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle \tag{A.7.5}$$

and the transverse Ward identities are

$$p_1^{\nu_1} \langle\!\langle T_{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2 a}(\boldsymbol{p}_2) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle = p_1^{\nu_1} \langle\!\langle \frac{\delta T_{\mu_1\nu_1}}{\delta A^a_{\mu_2}}(\boldsymbol{p}_1, \boldsymbol{p}_2) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle, \tag{A.7.6}$$

$$p_{2\mu_2} \langle\!\langle T_{\mu_1\nu_1}(\boldsymbol{p}_1) J^{\mu_2 a}(\boldsymbol{p}_2) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle = 2p_{2\mu_2} \langle\!\langle \frac{\delta J^{\mu_2 a}}{\delta g^{\mu_1\nu_1}}(\boldsymbol{p}_2, \boldsymbol{p}_1) \mathcal{O}^I(\boldsymbol{p}_3) \rangle\!\rangle. \tag{A.7.7}$$

If $\beta_3 = \Delta_3 - \frac{d}{2} > 0$, the same reasoning as in section 7.2 shows that the right-hand sides of (A.7.6, A.7.7) vanish in the zero-momentum limit. Then, by the results of section 5.3.5, in the remaining cases the coefficient of p_3^0 in the series expansion of the right-hand sides of (A.7.6, A.7.7) is at most ultralocal. The secondary CWIs lead to the following equations

$$\begin{aligned}0 = {} & \frac{(\Delta_3 - 2d - \epsilon)(\Delta_3 - d - 2 - \epsilon)}{2(\Delta_3 - d - 1 - \epsilon)} l_{\frac{d}{2}+\epsilon\{\frac{d}{2}, \frac{d}{2}-1, \Delta_3 - \frac{d}{2}\}} \times \\ & \times \left[\left(\Delta_3^2 - \Delta_3(d - 2 + 2\epsilon) + \epsilon(2 + d + \epsilon) \right) \alpha_1^{aI} + \alpha_2^{aI} \right],\end{aligned} \tag{A.7.8}$$

$$\begin{aligned}0 = {} & \frac{(\Delta_3 - 2d - \epsilon)(\Delta_3 - d - \epsilon)}{2(\Delta_3 - d - 1 - \epsilon)} l_{\frac{d}{2}+\epsilon\{\frac{d}{2}, \frac{d}{2}-1, \Delta_3 - \frac{d}{2}\}} \times \\ & \times \left[\left((\Delta_3 - \epsilon)^2 - d(\Delta_3 + 2 - \epsilon) + 4(\Delta_3 + 1 + \epsilon) \right) \alpha_1^{aI} + 2\alpha_2^{aI} \right],\end{aligned} \tag{A.7.9}$$

$$0 = (\Delta_3 - d - \epsilon) l_{\frac{d}{2}+\epsilon\{\frac{d}{2}, \frac{d}{2}-1, \Delta_3 - \frac{d}{2}\}} \left[2(2\Delta_3 - d)\alpha_1^{aI} + \alpha_2^{aI} \right]. \tag{A.7.10}$$

If the $\epsilon \to 0$ limit exists, we find three equations whose only solutions are either $\alpha_1^{aI} = \alpha_2^{aI} = 0$, or else $\alpha_2^{aI} = 2(d-4)\alpha_1^{aI}$ and $\Delta_3 = 2$. The second solution, however, is not valid since $\Delta_3 = 2$ is a special case where the regulator cannot be removed. Instead one must analyse all the special cases when the regulator cannot be removed from $l_{\frac{d}{2}+\epsilon\{\frac{d}{2}, \frac{d}{2}-1, \Delta_3 - \frac{d}{2}\}}$. The analysis is identical to that of section 5.3.6 and leads to the conclusion that $\alpha_1^{aI} = \alpha_2^{aI} = 0$.

Unlike in section 5.3.6, there are no additional conditions following from the coefficients of $p_3^{2\beta_3}$ or $p_3^{2\beta_3} \log p_3$ in the series expansion in p_3 of the secondary CWIs (A.7.8) - (A.7.10). Recall that such additional constraints arise when the equations following from the coefficients of $p_3^{2\beta_3}$ or $p_3^{2\beta_3} \log p_3$ are more singular than the equations following from the zero-momentum limit. In our case, it turns out that all coefficients of $p_3^{2\beta_3}$ or $p_3^{2\beta_3} \log p_3$ on the left-hand sides of (A.7.8) - (A.7.10) can be written in terms of $l_{\frac{d}{2}+\epsilon\{\frac{d}{2}, \frac{d}{2}-1, -\Delta_3 + \frac{d}{2}\}}$, accounting for all possible singularities. One can check that $l_{\frac{d}{2}+\epsilon\{\frac{d}{2}, \frac{d}{2}-1, -\Delta_3 + \frac{d}{2}\}}$ cannot be more singular than $l_{\frac{d}{2}+\epsilon\{\frac{d}{2}, \frac{d}{2}-1, \Delta_3 - \frac{d}{2}\}}$ assuming the unitarity bound $\Delta_3 \geq \frac{d}{2} - 1$.

A.8 Form of scaling anomalies

In this appendix we list the most most general form for the anomalous part of the trace Ward identities, and evaluate the resulting contribution to the reconstruction formulae. Following the discussion at the end of section 8.2, the most general anomalous contribution to the trace Ward

identities takes the form

$$\langle\!\langle T(\boldsymbol{p}_1)\mathcal{O}^I(\boldsymbol{p}_2)\mathcal{O}^J(\boldsymbol{p}_3)\rangle\!\rangle_{\text{anomaly}} = B_1^{IJ}, \tag{A.8.1}$$

$$\langle\!\langle T(\boldsymbol{p}_1)J^{\mu_2 a}(\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle_{\text{anomaly}} = \pi^{\mu_2}_{\alpha_2}(\boldsymbol{p}_2)p_3^{\alpha_2}\cdot B_1^{aI}, \tag{A.8.2}$$

$$\langle\!\langle T(\boldsymbol{p}_1)J^{\mu_2 a_2}(\boldsymbol{p}_2)J^{\mu_3 a_3}(\boldsymbol{p}_3)\rangle\!\rangle_{\text{anomaly}} = \pi^{\mu_2}_{\alpha_2}(\boldsymbol{p}_2)\pi^{\mu_3}_{\alpha_3}(\boldsymbol{p}_3)\left[B_1^{a_2a_3}p_3^{\alpha_2}p_1^{\alpha_3} + B_2^{a_2a_3}\delta^{\alpha_2\alpha_3}\right], \tag{A.8.3}$$

$$\langle\!\langle T(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle_{\text{anomaly}} = \Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)p_3^{\alpha_2}p_3^{\beta_2}B_1^I + \pi^{\mu_2\nu_2}(\boldsymbol{p}_2)B_2^I, \tag{A.8.4}$$

$$\begin{aligned}\langle\!\langle T(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)J^{\mu_3 a}(\boldsymbol{p}_3)\rangle\!\rangle_{\text{anomaly}} &= \Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)\pi^{\mu_3}_{\alpha_3}(\boldsymbol{p}_3)\left[B_1^a p_3^{\alpha_2}p_3^{\beta_2}p_1^{\alpha_3} + B_2^a\delta^{\alpha_2\alpha_3}p_3^{\beta_2}\right]\\ &+ \pi^{\mu_2\nu_2}(\boldsymbol{p}_2)\pi^{\mu_3}_{\alpha_3}(\boldsymbol{p}_3)p_1^{\alpha_3}B_3^a,\end{aligned} \tag{A.8.5}$$

$$\begin{aligned}\langle\!\langle T(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)T^{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\!\rangle_{\text{anomaly}} &= \Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)\Pi^{\mu_3\nu_3}_{\alpha_3\beta_3}(\boldsymbol{p}_3)\left[B_1 p_3^{\alpha_2}p_3^{\beta_2}p_1^{\alpha_3}p_1^{\beta_3}\right.\\ &\left.+ B_2\delta^{\beta_2\beta_3}p_3^{\alpha_2}p_1^{\alpha_3} + B_3\delta^{\alpha_2\alpha_3}\delta^{\beta_2\beta_3}\right]\\ &+ \pi^{\mu_2\nu_2}(\boldsymbol{p}_2)\Pi^{\mu_3\nu_3}_{\alpha_3\beta_3}(\boldsymbol{p}_3)p_1^{\alpha_3}p_1^{\beta_3}B_4 + \Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)\pi^{\mu_3\nu_3}(\boldsymbol{p}_3)p_3^{\alpha_2}p_3^{\beta_2}B_4(p_2\leftrightarrow p_3)\\ &+ \pi^{\mu_2\nu_2}(\boldsymbol{p}_2)\pi^{\mu_3\nu_3}(\boldsymbol{p}_3)B_5,\end{aligned} \tag{A.8.6}$$

where the form factors B_j are polynomials in momenta squared p_j^2, $j = 1, 2, 3$. In particular B_1^{IJ} can be non-zero only if $2\Delta = d + 2n$ for some non-negative integer n.

The anomaly in the trace of the stress-energy tensor leads to specific anomalous contributions to the full correlation functions. These anomalous contributions can be recovered from the expressions above by the reconstruction formulae using the results of the section 3.3 and equation (3.21). In the presence of the trace anomaly, the following expressions should be added to the right-hand sides of the corresponding reconstruction equations, such as (R.11.4), listed in part II of the paper.

$$\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)\mathcal{O}^I(\boldsymbol{p}_2)\mathcal{O}^J(\boldsymbol{p}_3)\rangle\!\rangle_{\text{anomaly}} = \frac{\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)}{d-1}B_1^{IJ}, \tag{A.8.7}$$

$$\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)J^a(\boldsymbol{p}_1)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle_{\text{anomaly}} = \frac{\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)}{d-1}\pi^{\mu_2}_{\alpha_2}(\boldsymbol{p}_2)p_3^{\alpha_2}B_1^{aI}, \tag{A.8.8}$$

$$\begin{aligned}&\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)J^{\mu_2 a_2}(\boldsymbol{p}_2)J^{\mu_3 a_3}(\boldsymbol{p}_3)\rangle\!\rangle_{\text{anomaly}} =\\ &\qquad= \frac{\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)}{d-1}\pi^{\mu_2}_{\alpha_2}(\boldsymbol{p}_2)\pi^{\mu_3}_{\alpha_3}(\boldsymbol{p}_3)\left[B_1^{a_2a_3}p_3^{\alpha_2}p_1^{\alpha_3} + B_2^{a_2a_3}\delta^{\alpha_2\alpha_3}\right],\end{aligned} \tag{A.8.9}$$

$$\begin{aligned}\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)\mathcal{O}^I(\boldsymbol{p}_3)\rangle\!\rangle_{\text{anomaly}} &= \frac{\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)}{d-1}\left[\Pi^{\mu_2\nu_2}_{\alpha_2\beta_2}(\boldsymbol{p}_2)p_3^{\alpha_2}p_3^{\beta_2}B_1^I + \pi^{\mu_2\nu_2}(\boldsymbol{p}_2)B_2^I\right]\\ &+ [(\boldsymbol{p}_1,\mu_1,\nu_1)\leftrightarrow(\boldsymbol{p}_2,\mu_2,\nu_2)] - \frac{\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)\pi^{\mu_2\nu_2}(\boldsymbol{p}_2)}{d-1}B_2^I,\end{aligned} \tag{A.8.10}$$

$$\begin{aligned}\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)J^{\mu_3 a}(\boldsymbol{p}_3)\rangle\!\rangle_{\text{anomaly}} &= \frac{\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)}{d-1}\langle\!\langle T(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)J^a(\boldsymbol{p}_3)\rangle\!\rangle_{\text{anomaly}}\\ &+ [(\boldsymbol{p}_1,\mu_1,\nu_1)\leftrightarrow(\boldsymbol{p}_2,\mu_2,\nu_2)]\\ &- \frac{\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)\pi^{\mu_2\nu_2}(\boldsymbol{p}_2)}{d-1}\pi^{\mu_3}_{\alpha_3}(\boldsymbol{p}_3)p_1^{\alpha_3}B_3^a,\end{aligned} \tag{A.8.11}$$

$$\begin{aligned}\langle\!\langle T^{\mu_1\nu_1}(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)T^{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\!\rangle_{\text{anomaly}} &= \frac{\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)}{d-1}\langle\!\langle T(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)T^{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\!\rangle_{\text{anomaly}}\\ &[(\boldsymbol{p}_1,\mu_1,\nu_1)\leftrightarrow(\boldsymbol{p}_2,\mu_2,\nu_2)] + [(\boldsymbol{p}_1,\mu_1,\nu_1)\leftrightarrow(\boldsymbol{p}_3,\mu_3,\nu_3)]\\ &- \frac{\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)\pi^{\mu_2\nu_2}(\boldsymbol{p}_2)}{d-1}\left[\Pi^{\mu_3\nu_3}_{\alpha_3\beta_3}(\boldsymbol{p}_3)B_4(p_1,p_2,p_3) + \pi^{\mu_3\nu_3}(\boldsymbol{p}_3)B_5(p_1,p_2,p_3)\right]\\ &[(\boldsymbol{p}_1,\mu_1,\nu_1)\leftrightarrow(\boldsymbol{p}_3,\mu_3,\nu_3)] + [(\boldsymbol{p}_2,\mu_2,\nu_2)\leftrightarrow(\boldsymbol{p}_3,\mu_3,\nu_3)]\\ &+ \frac{\pi^{\mu_1\nu_1}(\boldsymbol{p}_1)\pi^{\mu_2\nu_2}(\boldsymbol{p}_2)\pi^{\mu_3\nu_3}(\boldsymbol{p}_3)}{d-1}B_5.\end{aligned} \tag{A.8.12}$$

If present, such contributions should be added to the reconstruction formulae for the corresponding 3-point functions presented in part II.

In particular, in $d = 4$ the anomalous trace Ward identity is

$$\langle T\rangle = \frac{1}{2}P^{IJ}\phi_0^I\Box^{\Delta-2}\phi_0^J + \frac{1}{4}\kappa F^a_{\mu\nu}F^{\mu\nu a} + aE_4 + cW^2, \tag{A.8.13}$$

where W^2 is the square of the Weyl tensor and E_4 is the Euler density, (8.3). From this we find

$$B_1^{IJ} = (-p_1^2)^{\Delta-2}P^{IJ}, \tag{A.8.14}$$
$$B_1^{aI} = 0, \tag{A.8.15}$$
$$B_1^{a_2a_3} = -\kappa\delta^{a_2a_3}, \qquad B_2^{a_2a_3} = \tfrac{1}{2}(p_2^2 + p_3^2 - p_1^2)\kappa\delta^{a_2a_3}, \tag{A.8.16}$$
$$B_1^I = 0, \tag{A.8.17}$$
$$B_1^a = B_2^a = B_3^a = 0. \tag{A.8.18}$$

For $\langle\!\langle T(\boldsymbol{p}_1)T^{\mu_2\nu_2}(\boldsymbol{p}_2)T^{\mu_3\nu_3}(\boldsymbol{p}_3)\rangle\!\rangle$,

$$B_1 = 8(a+c), \tag{A.8.19}$$
$$B_2 = 8(p_1^2 - p_2^2 - p_3^2)(a+c), \tag{A.8.20}$$
$$B_3 = -2\lambda^2(a+c) + 4p_2^2p_3^2c, \tag{A.8.21}$$
$$B_4 = -\frac{8}{3}ap_3^2, \tag{A.8.22}$$
$$B_5 = \frac{4}{9}a\lambda^2. \tag{A.8.23}$$

These results are in agreement with the results of sections 8.2 and 8.3.

A.9 Identities with projectors

The projectors are defined as

$$\pi^\mu_\alpha = \delta^\mu_\alpha - \frac{p^\mu p_\alpha}{p^2}, \tag{A.9.1}$$
$$\Pi^{\mu\nu}_{\alpha\beta} = \frac{1}{2}\left(\pi^\mu_\alpha\pi^\nu_\beta + \pi^\mu_\beta\pi^\nu_\alpha\right) - \frac{1}{d-1}\pi^{\mu\nu}\pi_{\alpha\beta}, \tag{A.9.2}$$
$$\Pi^{\mu\nu\rho\sigma} = \delta^{\rho\alpha}\delta^{\sigma\beta}\Pi^{\mu\nu}_{\alpha\beta}, \tag{A.9.3}$$

One can find the following identities:

$$p_\mu\pi^{\mu\nu} = p_\mu\Pi^{\mu\nu\rho\sigma} = 0, \tag{A.9.4}$$
$$\delta_{\mu\nu}\pi^{\mu\nu} = \pi^\mu_\mu = d-1, \tag{A.9.5}$$
$$\Pi^{\mu\nu\rho}{}_\rho = \delta_{\rho\sigma}\Pi^{\mu\nu\rho\sigma} = \pi_{\rho\sigma}\Pi^{\mu\nu\rho\sigma} = 0, \tag{A.9.6}$$
$$\Pi^{\mu\rho\nu}_\rho = \delta_{\rho\sigma}\Pi^{\mu\rho\nu\sigma} = \pi_{\rho\sigma}\Pi^{\mu\rho\nu\sigma} = \frac{(d+1)(d-2)}{2(d-1)}\pi^{\mu\nu}, \tag{A.9.7}$$
$$\pi^\mu_\alpha\pi^\alpha_\nu = \pi^\mu_\nu, \tag{A.9.8}$$
$$\Pi^{\mu\nu}_{\alpha\beta}\Pi^{\alpha\beta}_{\rho\sigma} = \Pi^{\mu\nu}_{\rho\sigma}, \tag{A.9.9}$$
$$\Pi^{\mu\nu\rho}{}_\alpha\pi^{\alpha\sigma} = \Pi^{\mu\nu\rho\sigma}, \tag{A.9.10}$$
$$\Pi^{\mu\nu}_{\alpha\beta}\Pi^{\alpha\rho\beta\sigma} = \frac{d-3}{2(d-1)}\Pi^{\mu\nu\rho\sigma}. \tag{A.9.11}$$

Basic derivatives can be calculated directly. Denoting

$$\partial_\mu = \frac{\partial}{\partial p^\mu}. \tag{A.9.12}$$

we find

$$\partial_\kappa \pi_{\mu\nu} = -\frac{p_\mu}{p^2}\pi_{\nu\kappa} - \frac{p_\nu}{p^2}\pi_{\mu\kappa}, \tag{A.9.13}$$

$$\partial_\kappa \Pi_{\mu\nu\rho\sigma} = -\frac{p_\mu}{p^2}\Pi_{\kappa\nu\rho\sigma} - \frac{p_\nu}{p^2}\Pi_{\mu\kappa\rho\sigma} - \frac{p_\rho}{p^2}\Pi_{\mu\nu\kappa\sigma} - \frac{p_\sigma}{p^2}\Pi_{\mu\nu\rho\kappa}, \tag{A.9.14}$$

$$\pi^\mu_\alpha \partial_\kappa \pi^\alpha_\nu = -\frac{p_\nu}{p^2}\pi^\mu_\kappa, \tag{A.9.15}$$

$$\pi^{\mu\kappa}\partial_\alpha \pi^\alpha_\nu - \pi^{\mu\alpha}\partial_\alpha \pi^\kappa_\nu = -(d-2)\frac{p_\nu}{p^2}\pi^{\mu\kappa} + \frac{p^\kappa}{p^2}\pi^\mu_\nu, \tag{A.9.16}$$

$$\Pi^{\mu\nu}_{\alpha\beta}\partial_\kappa \Pi^{\alpha\beta}_{\rho\sigma} = -\frac{p_\rho}{p^2}\Pi^{\mu\nu}_{\kappa\sigma} - \frac{p_\sigma}{p^2}\Pi^{\mu\nu}_{\rho\kappa}, \tag{A.9.17}$$

$$\Pi^{\mu\nu}_{\kappa\beta}\partial_\alpha \Pi^{\alpha\beta}_{\rho\sigma} - \Pi^{\mu\nu\alpha}{}_{\beta}\partial_\alpha \Pi^{\kappa\beta}_{\rho\sigma} = -\frac{1}{2}\frac{d-1}{p^2}\left[p_\rho \Pi^{\mu\nu}_{\kappa\sigma} + p_\sigma \Pi^{\mu\nu}_{\rho\kappa}\right] + \frac{p_\kappa}{p^2}\Pi^{\mu\nu}_{\rho\sigma}. \tag{A.9.18}$$

Analogous expressions with two derivatives are

$$\pi^\mu_\alpha \partial^2 \pi^\alpha_\nu = -\frac{2}{p^2}\pi^\mu_\nu, \tag{A.9.19}$$

$$p^\alpha \pi^\mu_\beta \partial_\alpha \partial_\kappa \pi^\beta_\nu = \frac{p_\nu}{p^2}\pi^\mu_\kappa, \tag{A.9.20}$$

$$\Pi^{\mu\nu}_{\alpha\beta}\partial^2 \Pi^{\alpha\beta}_{\rho\sigma} = -\frac{4}{p^2}\Pi^{\mu\nu}_{\rho\sigma}, \tag{A.9.21}$$

$$p^\gamma \Pi^{\mu\nu}_{\alpha\beta}\partial_\gamma \partial_\kappa \Pi^{\alpha\beta}_{\rho\sigma} = \frac{p_\rho}{p^2}\Pi^{\mu\nu}_{\kappa\sigma} + \frac{p_\sigma}{p^2}\Pi^{\mu\nu}_{\rho\kappa}. \tag{A.9.22}$$

For the semi-local operators defined in (3.17) and (3.18) we find

$$\pi^\mu_\alpha \partial_\kappa j^\alpha_{\rm loc} = \frac{1}{p^2}\pi^\mu_\kappa r, \tag{A.9.23}$$

$$\pi^{\mu\kappa}\partial_\alpha j^\alpha_{\rm loc} - \pi^{\mu\alpha}\partial_\alpha j^\kappa_{\rm loc} = \frac{d-3}{p^2}\pi^{\mu\kappa} r + \frac{1}{p^2}\pi^{\mu\kappa} p^\alpha \partial_\alpha r - \frac{p^\kappa}{p^2}\pi^{\mu\alpha}\partial_\alpha r, \tag{A.9.24}$$

$$\pi^\mu_\alpha \partial^2 j^\alpha_{\rm loc} = \frac{2}{p^2}\pi^{\mu\alpha}\partial_\alpha r, \tag{A.9.25}$$

$$p^\alpha \pi^\mu_\beta \partial_\alpha \partial_\kappa j^\beta_{\rm loc} = -\frac{2}{p^2}\pi^\mu_\kappa r + \frac{1}{p^2}\pi^\mu_\kappa p^\alpha \partial_\alpha r \tag{A.9.26}$$

and

$$\Pi^{\mu\nu}_{\alpha\beta}\partial_\kappa t^{\alpha\beta}_{\rm loc} = \frac{2}{p^2}\Pi^{\mu\nu}_{\alpha\kappa} R^\alpha, \tag{A.9.27}$$

$$\Pi^{\mu\nu\kappa}{}_{\beta}\partial_\alpha t^{\alpha\beta}_{\rm loc} - \Pi^{\mu\nu\alpha}{}_{\beta}\partial_\alpha t^{\kappa\beta}_{\rm loc} = \frac{d-2}{p^2}\Pi^{\mu\nu\kappa}{}_{\alpha} R^\alpha + \frac{p^\beta}{p^2}\Pi^{\mu\nu\kappa}{}_{\alpha}\partial_\beta R^\alpha - \frac{p^\kappa}{p^2}\Pi^{\mu\nu\alpha}{}_{\beta}\partial_\alpha R^\beta, \tag{A.9.28}$$

$$\Pi^{\mu\nu}_{\alpha\beta}\partial^2 t^{\alpha\beta}_{\rm loc} = \frac{4}{p^2}\Pi^{\mu\nu\alpha}{}_{\beta}\partial_\alpha R^\beta, \tag{A.9.29}$$

$$p^\gamma \Pi^{\mu\nu}_{\alpha\beta}\partial_\gamma \partial_\kappa t^{\alpha\beta}_{\rm loc} = -\frac{4}{p^2}\Pi^{\mu\nu}_{\alpha\kappa} R^\alpha + \frac{2}{p^2}\Pi^{\mu\nu}_{\alpha\kappa} p^\beta \partial_\beta R^\alpha. \tag{A.9.30}$$

Acknowledgments: We would like to thank Anatoly Dymarsky for discussions. Research at the Perimeter Institute is supported by the Government of Canada through Industry Canada and by the Province of Ontario through the Ministry of Research & Innovation. K.S. acknowledges support from NWO via a VICI grant and from a grant of the John Templeton Foundation. The opinions expressed in this publication are those of the authors and do not necessarily reflect the views of the John Templeton Foundation. A.B. is supported through NWO via the VICI grant of K.S.

References

[1] A. M. Polyakov, *Conformal symmetry of critical fluctuations*, *JETP Lett.* **12** (1970) 381–383.

[2] E. Schreier, *Conformal symmetry and three-point functions*, *Phys.Rev.* **D3** (1971) 980–988.

[3] H. Osborn and A. Petkou, *Implications of conformal invariance in field theories for general dimensions*, *Annals Phys.* **231** (1994) 311–362, [hep-th/9307010].

[4] J. Erdmenger and H. Osborn, *Conserved currents and the energy momentum tensor in conformally invariant theories for general dimensions*, *Nucl.Phys.* **B483** (1997) 431–474, [hep-th/9605009].

[5] S. Giombi, S. Prakash, and X. Yin, *A note on CFT correlators in three dimensions*, arXiv:1104.4317.

[6] M. S. Costa, J. Penedones, D. Poland, and S. Rychkov, *Spinning conformal correlators*, *JHEP* **1111** (2011) 071, [arXiv:1107.3554].

[7] M. S. Costa, J. Penedones, D. Poland, and S. Rychkov, *Spinning conformal blocks*, *JHEP* **1111** (2011) 154, [arXiv:1109.6321].

[8] J. Maldacena and A. Zhiboedov, *Constraining conformal field theories with a higher spin symmetry*, arXiv:1112.1016.

[9] J. Maldacena and A. Zhiboedov, *Constraining conformal field theories with a slightly broken higher spin symmetry*, arXiv:1204.3882.

[10] Y. S. Stanev, *Correlation functions of conserved currents in four dimensional conformal field theory*, *Nucl.Phys.* **B865** (2012) 200–215, [arXiv:1206.5639].

[11] A. Zhiboedov, *A note on three-point functions of conserved currents*, arXiv:1206.6370.

[12] D. Z. Freedman, K. Johnson, and J. I. Latorre, *Differential regularization and renormalization: A new method of calculation in quantum field theory*, *Nucl.Phys.* **B371** (1992) 353–414.

[13] C. Coriano, L. Delle Rose, E. Mottola, and M. Serino, *Graviton vertices and the mapping of anomalous correlators to momentum space for a general conformal field theory*, *JHEP* **1208** (2012) 147, [arXiv:1203.1339].

[14] P. McFadden and K. Skenderis, *Holography for cosmology*, *Phys. Rev.* **D81** (2010) 021301, [arXiv:0907.5542].

[15] P. McFadden and K. Skenderis, *The Holographic Universe*, *J. Phys. Conf. Ser.* **222** (2010) 012007, [arXiv:1001.2007].

[16] P. McFadden and K. Skenderis, *Holographic non-Gaussianity*, *JCAP* **1105** (2011) 013, [arXiv:1011.0452].

[17] A. Bzowski, P. McFadden, and K. Skenderis, *Holographic predictions for cosmological 3-point functions*, *JHEP* **1203** (2012) 091, [arXiv:1112.1967].

[18] A. Bzowski, P. McFadden, and K. Skenderis, *Holography for inflation using conformal perturbation theory*, arXiv:1211.4550.

[19] I. Antoniadis, P. O. Mazur, and E. Mottola, *Conformal invariance, dark energy, and CMB non-Gaussianity*, *JCAP* **1209** (2012) 024, [arXiv:1103.4164].

[20] J. M. Maldacena and G. L. Pimentel, *On graviton non-Gaussianities during inflation*, arXiv:1104.2846.

[21] P. Creminelli, J. Norena, and M. Simonovic, *Conformal consistency relations for single-field inflation*, *JCAP* **1207** (2012) 052, [arXiv:1203.4595].

[22] A. Kehagias and A. Riotto, *Operator product expansion of inflationary correlators and conformal symmetry of de Sitter*, *Nucl.Phys.* **B864** (2012) 492–529, [arXiv:1205.1523].

[23] A. Kehagias and A. Riotto, *The four-point correlator in multifield inflation, the operator product expansion and the symmetries of de Sitter*, arXiv:1210.1918.

[24] C. Coriano, L. Delle Rose, and M. Serino, *Three and four point functions of stress energy tensors in d=3 for the analysis of cosmological non-gaussianities*, arXiv:1210.0136.

[25] I. Mata, S. Raju, and S. Trivedi, *CMB from CFT*, arXiv:1211.5482.

[26] M. Giannotti and E. Mottola, *The trace anomaly and massless scalar degrees of freedom in gravity*, *Phys.Rev.* **D79** (2009) 045014, [arXiv:0812.0351].

[27] R. Armillis, C. Coriano, L. Delle Rose, and M. Guzzi, *Anomalous U(1) models in four and five dimensions and their anomaly poles*, *JHEP* **0912** (2009) 029, [arXiv:0905.0865].

[28] R. Armillis, C. Coriano, and L. Delle Rose, *Conformal anomalies and the gravitational effective action: The TJJ correlator for a Dirac fermion*, *Phys.Rev.* **D81** (2010) 085001, [arXiv:0910.3381].

[29] D. Chowdhury, S. Raju, S. Sachdev, A. Singh, and P. Strack, *Multipoint correlators of conformal field theories: implications for quantum critical transport*, arXiv:1210.5247.

[30] A. Cappelli, R. Guida, and N. Magnoli, *Exact consequences of the trace anomaly in four-dimensions*, *Nucl.Phys.* **B618** (2001) 371–406, [hep-th/0103237].

[31] E. Barnes, D. Vaman, C. Wu, and P. Arnold, *Real-time finite-temperature correlators from AdS/CFT*, *Phys.Rev.* **D82** (2010) 025019, [arXiv:1004.1179].

[32] A. Petkou and K. Skenderis, *A non-renormalization theorem for conformal anomalies*, *Nucl.Phys.* **B561** (1999) 100–116, [hep-th/9906030].

[33] A. I. Davydychev, *Recursive algorithm of evaluating vertex type Feynman integrals*, *J.Phys.* **A25** (1992) 5587–5596.

[34] C. Coriano, L. D. Rose, E. Mottola, and M. Serino, *Solving the conformal constraints for scalar operators in momentum space and the evaluation of Feynman's master integrals*, arXiv:1304.6944.

[35] A. Dymarsky, *Comments on the four point function of the stress-energy tensors in a conformal field theory*, to appear.

[36] P. Di Francesco, P. Mathieu, and D. Senechal, *Conformal field theory*. Springer, New York, 1997.

[37] D. Z. Freedman, S. D. Mathur, A. Matusis, and L. Rastelli, *Correlation functions in the CFT(d) / AdS(d+1) correspondence*, *Nucl.Phys.* **B546** (1999) 96–118, [hep-th/9804058].

[38] M. Abramowitz and e. Stegun, Irene A., *Handbook of mathematical functions with formulas, graphs, and mathematical tables*. New York, Dover Publications, 1972.

[39] P. Appell and M. Kampé de Fériet, *Fonctions hypegeométriques et hypersphériques. Polynomes d' Hermite.* Gauthier-Villars, 1926.

[40] H. Bateman, A. Erdélyi, and U. S. O. of Naval Research, *Bateman Manuscript Project, Higher transcendental functions.* No. v. 1 in Higher Transcendental Functions. McGraw-Hill, 1953.

[41] H. Exton, *On the system of partial differential equations associated with Appell's function F_4*, *J. Phys. A: Math. Gen.* **28** (1995) 631–641.

[42] A. Prudnikov, Y. Brychkov, and O. Marichev, *Integrals and series, vol. 2.* Gordon and Breach, 1992.

[43] L. Rosenberg, *Electromagnetic interactions of neutrinos*, *Phys.Rev.* **129** (1963) 2786–2788.

[44] A. Denner and S. Dittmaier, *Reduction schemes for one-loop tensor integrals*, *Nucl.Phys.* **B734** (2006) 62–115, [hep-ph/0509141].

[45] T. Mehen, I. W. Stewart, and M. B. Wise, *Conformal invariance for nonrelativistic field theory*, *Phys.Lett.* **B474** (2000) 145–152, [hep-th/9910025].

[46] S. Raju, *New recursion relations and a flat space limit for AdS/CFT correlators*, *Phys.Rev.* **D85** (2012) 126009, [arXiv:1201.6449].

[47] S. Raju, *Four point functions of the stress tensor and conserved currents in AdS_4/CFT_3*, *Phys.Rev.* **D85** (2012) 126008, [arXiv:1201.6452].

[48] S. Weinberg, *Minimal fields of canonical dimensionality are free*, *Phys.Rev.* **D86** (2012) 105015, [arXiv:1210.3864].

[49] G. 't Hooft and M. Veltman, *Scalar one loop integrals*, *Nucl.Phys.* **B153** (1979) 365–401.

[50] G. Passarino and M. Veltman, *One loop corrections for e^+e^- annihilation into $\mu^+\mu^-$ in the Weinberg Model*, *Nucl.Phys.* **B160** (1979) 151.

[51] R. M. Wald, *General Relativity.* Chicago, Univ. Pr., 1984.

[52] N. Birrell and P. Davies, *Quantum fields in curved space.* Cambridge, Univ. Pr., 1982.

[53] M. E. Peskin and D. V. Schroeder, *An introduction to quantum field theory.* Westview Press Inc., Boulder, 1995.